

6-19-00

**CONTINUING PATENT APPLICATION TRANSMITTAL**  
 (for Continuing Applications  
 under 37 C.F.R. §1.53(b))

Attorney Docket No. 69176

First Named Inventor or  
 Application Identifier: Biedermann et

Box PATENT APPLICATION  
 Commissioner of Patents and Trademarks  
 ATTENTION: Assistant Commissioner  
 for Patents  
 Washington, D.C. 20231

Sir:

This is a request under 37 C.F.R.  
 §1.53(b) for filing a:

- (X) Continuation application,  
 ( ) Divisional application,  
 ( ) Continuation-in-Part application,

pending prior application number PCT/EP98/08268,

filed on 16 December 1998 of Biedermann et al.

(Date) (Inventor(s))  
 or NEW PIPERAZINYL-SUBSTITUTED PYRIDYLALKANE, ALKENE AND ALKINE  
 (Title)  
CARBOXAMIDES

1. ( ) This is a continuation or divisional application. Enclosed is a copy of the prior application as originally filed, including specification, claims, drawings, and oath or declaration.

- or -

- (X) Enclosed is a patent application (for continuation, divisional, or continuation-in-part applications) containing:

(X) 219 pages of the specification (including claims).

( ) \_\_\_\_\_ sheets of drawings ( ) Formal ( ) Informal.

2. (X) Amend the specification by inserting before the first line the sentence: --This is a [X] continuation, [ ] division, [ ] continuation-in-part, of prior application number PCT/EP98/08268, filed 16 December 1998 and designating the U.S., which is hereby incorporated herein by reference in its entirety.-- The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under paragraph 3 below, is considered as being part of the disclosure of the accompanying application, and is hereby incorporated by reference therein.

06/16/00



06/16/00

13844 U.S. PTO

05596001



06/16/00

) CERTIFICATE OF MAILING BY "EXPRESS MAIL"

) "Express Mail" Mailing Label Number

) EL 485 057 857 US

) Date of Deposit June 16, 2000

) I hereby certify that this paper or fee is being  
 deposited with the United States Postal Service  
 "Express Mail Post Office to Addressee" Service  
 under 37 CFR §1.10 on the date indicated above and  
 is addressed to the Commissioner of Patents and  
 Trademarks, Washington, D.C. 20231.

) ED PRICE

) (Typed or printed name of person mailing)

) (Signature of person mailing)

3. ( ) A copy of the executed oath or declaration filed in the prior nonprovisional application is enclosed.

4. (X) Inventorship:

( ) A newly-executed oath or declaration and power of attorney is enclosed (for continuation-in-part applications, or for continuation or divisional applications naming an inventor not named in the prior application) (§1.63(a), (d)(5) and (e)).

( ) Because this application is being filed by fewer than all of the inventors named in the prior application, delete the following inventor(s) named in the prior nonprovisional application (37 C.F.R. §1.63(d)(1)(2)):

---



---

(X) The names of persons believed to be the actual inventors are set forth in the enclosed unexecuted oath or declaration and power of attorney (§1.41(a) and §1.53(b)).

( ) Assignment(s) of the invention to \_\_\_\_\_, and cover sheet are enclosed.

( ) A check in the amount of \$\_\_\_\_\_ to cover the fee for recording the assignment(s) is enclosed.

( ) The prior application is assigned of record to \_\_\_\_\_.

( ) Small Entity Status (37 C.F.R. §1.28(a)(2)):

( ) A statement of status as a small entity is enclosed.

( ) A statement of status as a small entity was filed in the prior application, and small entity status is still proper and desired in this new nonprovisional application.

( ) Status as a small entity is no longer claimed.

8. ( ) A 37 C.F.R. §3.73(b) statement is enclosed (where an assignee seeks to take action in a matter before the Patent Office).

9. (X) A preliminary amendment is enclosed.

10. ( ) Drawings:

( ) Transfer the drawings from the prior application to this application and abandon said prior application as of the filing date accorded this application. A duplicate copy of this sheet is enclosed for filing in the prior application file. (May be used only if signed by person authorized by §1.138 and before payment of base issue fee.)

( ) New formal drawings are enclosed.

( ) Informal drawings are enclosed.

11. (X) A separate written request under 37 C.F.R. §1.136(a)(3), which is a general authorization to treat any concurrent or future reply requiring a petition for an extension of time under 37 C.F.R. §1.136(a) for its timely submission as incorporating a petition for an extension of time for the appropriate length of time, is enclosed.

12. ( ) An Information Disclosure Statement is enclosed.

( ) A Form PTO-1449 is enclosed.

( ) \_\_\_\_\_ References (copies) listed on the Form PTO-1449 are enclosed.

3. ( ) A MicroFiche Computer Program (Appendix) is enclosed.

4. (X) A Return Receipt Postcard is enclosed (MPEP §503).

5. ( ) A Nucleotide and/or Amino Acid Sequence Submission is enclosed.

( ) A Computer Readable Copy is enclosed.

( ) A Paper Copy (Identical to Computer Copy) is enclosed.

( ) A Statement Verifying Identity of above Copies is enclosed.

6. (X) Priority of application number DE/19756236.1 filed on 17 December 1997 in Germany is claimed under 35 U.S.C. §119.

( ) The certified copy of the priority document has been filed in prior application number \_\_\_\_/\_\_\_\_, filed \_\_\_\_\_.

(X) A certified copy of the priority document is enclosed.

17. ( ) Power of Attorney:

( ) The power of attorney in the prior application is to:

( ) \_\_\_\_\_ Reg. No. \_\_\_\_\_,  
FITCH, EVEN, TABIN, & FLANNERY  
Suite 1600  
120 South LaSalle Street  
Chicago, Illinois 60603-3406  
and other members of the firm.

( ) Customer Number 22242.

( ) The power appears in the original papers in the prior application.

( ) Since the power does not appear in the original papers in the prior application, a copy of the power in the prior application is enclosed.

18. ( ) Cancel in this application original claims \_\_\_\_\_ of the prior application before calculating the filing fee. (At least one original independent claim must be retained for filing purposes.)

19. (X) The filing fee is calculated below:

Fee Calculation for Claims as Filed in the Prior Application, Less Any Claims Cancelled by Amendment			
( ) Basic Utility Fee		\$ 690.00	\$ _____
• ( ) Independent Claims _____ - 3 = _____	x \$ 78.00 =	\$ _____	
• ( ) Total Claims _____ - 20 = _____	x \$ 18.00 =	\$ _____	
• ( ) Fee for Multiply Dependent Claims	\$260.00	\$ _____	
or			
( ) Basic Design Fee		\$ 310.00	\$ _____
Total of above Calculations			\$ _____
Reduction by 50% for Filing by Small Entity			\$ _____
Total			\$ <u>DEFER</u>

20. ( ) A check in the amount of \$ \_\_\_\_\_ is enclosed.

20a. (X) **THE PAYMENT OF THE FILING FEE IS TO BE DEFERRED UNTIL THE DECLARATION IS FILED. DO NOT CHARGE OUR DEPOSIT ACCOUNT.**

21. (X) The Commissioner is hereby authorized to charge any fees which may be required under 37 C.F.R. §§1.16 and 1.17 and are not paid herewith, or credit any overpayment, to Deposit Account Number 06-1135. A duplicate copy of this request is enclosed.

22. ( ) Also enclosed:

23. (X) Address all future communications to Customer Number 22242.



FITCH, EVEN, TABIN & FLANNERY  
Suite 1600  
120 South LaSalle Street  
Chicago, Illinois 60603-3406  
Telephone: (312) 577-7000  
Facsimile: (312) 577-7007

June 16, 2000  
(Date)

James P. Krueger  
James P. Krueger  
Registration No. 35,234  
( ) Attorney or agent of record  
( ) Filed under \$1.34(a)



In the second line of claim 8, please delete "claims 1-7" and insert --claim 7--.

In the third line of claim 8, please delete "Characterized in that" and insert --wherein--.

In the first line of claim 9, please delete "Claims 1 to 7" and insert --claim 7--.

In the second line of claim 10, please delete "characterized in that" and insert --wherein--.

In the second line of claim 11, please delete "claims 1 to 7" and insert --claim 7--.

In the second line of claim 12, please delete "claim 1 to 7" and insert --claim 7--.

In the second line of claim 13, please delete "characterized in that" and insert --wherein--.

In the second line of claim 14, please delete "characterized in that" and insert --wherein--.

In the second line of claim 15, please delete "characterized in that" and insert --wherein--.

In the second line of claim 16, please delete "characterized in that" and insert --wherein--.

In the second line of claim 17, please delete "characterized in that" and insert --wherein--.

In the second line of claim 18, please delete "characterized in that" and insert --wherein--.

In the second line of claim 19, please delete "characterized in that" and insert --wherein--.

In the first line of claim 21, please delete "characterized in that" and insert --wherein--.

In the first line of claim 22, please delete "characterized in that" and insert --wherein--.

In the second line of claim 23, please delete "characterized in that" and insert --wherein--.

In the first line of claim 24, please delete "one of the claims 12 and 14 to 16, characterized in that" and insert --claim 16, wherein--.

In the first line of claim 25, please delete "characterized in that" and insert --wherein--.

In the first line of claim 26, please delete "characterized in that" and insert --wherein--.

In the first line of claim 27, please delete "one of the claims 16, 25 or 26, characterized in that" and insert --claim 26, wherein--.

In the first line of claim 28, please delete "characterized in that" and insert --wherein--.

In the second line of claim 29, please delete "characterized in that" and insert --wherein--.

In the first line of claim 30, please delete "one of the claims 12 and 14 to 28, characterized in that" and insert --claim 28, wherein--.

In the second line of claim 31, please delete "claims 1 to 6 and according to claim 7" and insert --claim 7, wherein--.


REMARKS

Upon entry of the present amendment, claims 4 through 19 and claims 21 through 31 have been amended and claim 1-31 are pending.

The Commissioner is hereby authorized to charge any additional fees which may be required in this application to Deposit Account No. 06-1135.

Respectfully submitted

FITCH, EVEN, TABIN & FLANNERY

By   
James P. Krueger  
Registration No. 35,234

June 16, 2000

120 S. LaSalle St., Suite 1600  
Chicago, Illinois 60603  
(312) 577-7000

New Piperaziny1-Substituted Pyridylalkane, Alkene and Alkine  
Carboxamides

0355001-051500

The invention relates to new piperaziny1-substituted pyridyl-alkane, alkene and alkine carboxamides with a saturated, one or several-fold unsaturated hydrocarbon residue in the carboxylic acid portion, methods for the synthesis of these compounds, medicaments containing these and their production as well as their therapeutic use especially as cytostatic agents and immunosuppressive agents, for example, in the treatment or prevention of various types of tumors and control of immune reactions, for example of autoimmune diseases.

A pressing need exists for new pharmaceuticals and/or medicaments for cytostatic therapy which not only possess a strong activity, but also exert diminished side effects in comparison to many classical cancerostatic agents, whereby treatment of a broad as possible spectrum of tumors should be made accessible. Furthermore, effective cytostatic agents for an efficient therapy should be made available. Active ingredients of this type should also be exceptionally suitable in the mentioned indications for a combination therapy, be it in connection with other cytostatic agents or with radiation (for example X-rays, radioactive elements, such as cobalt, or linear accelerator, etc.), with operative procedures, heat treatment, etc.

Additionally, from another point of view, there exists a strong need in the field of tumor therapy for new compounds, for example for overcoming or avoiding resistances, which enrich the pallet of cancerostatics based on new modes of action in the ideal case.

090106Z



2007

3

The synthesis and pharmacological evaluation of heterocyclic carboxamides which can be substituted at an end of the molecule by completely different heterocycles such as thiophene, guanine, indole, benzimidazole or indazole as well as pyridine are described in J. Med. Chem., 1996, pages 4692-4706. As opposed to the compounds according to the

invention, those published carboxamides possess an activity directed against psychoses. A few particularly named piperazine-substituted pyridyl carboxamides and/or their formula group therein are not encompassed by the present protective scope of the compounds of the invention according to the meanings given in the present claims because they have a direct bond instead of the structural element A. Based on the completely different therapeutic possibility of the known compounds in the field of psychiatry as compared to the indications according to the invention, it could not be expected that the compounds of the present invention would have the named cancerostatic and immunosuppressive effects.

Pyridyl ureas, pyridyl thioureas and pyridyl carbonamides, wherein the amide portion is bound over an aryl-substituted alkyl chain with a piperidine ring or piperidine ring or piperazine ring, are described for example in EP-A-0 428 434 or in EP-A-0 512 902 as antagonists of the neurokinin receptor and substance P. Furthermore, pyridyl(alkyl)carbonamides, pyridyl(alkyl)sulfonamides and analogous ureas, wherein the amide portion is bound to piperidine ring over an alkyl chain, are disclosed in EP-A-0 479 601 as active ingredients with anti-arrhythmic properties.

Further structurally closely related compounds are represented by the piperidine compounds described in EP-A-0 330 026. However, no 3-pyridyl derivatives were concretely described and no concrete examples were disclosed in this publication, aside from a single compound which is described below. These known compounds are distinguished by an anti-cholinesterase activity, an anti-amnesia activity as well as activities directed against hyperkinesia, senile dementia, mania and Alzheimer's disease.

In WO 91/15 485, the production of pyridine-3,5-dicarboxylic acid esters and amides as well as their use for the treatment of tumor conditions is described. These compounds differ from

the compounds according to the invention described below in very important structural features, for example by the dicarboxyl grouping on the pyridine ring or the absence of the hydrocarbon chain between the pyridine ring and the amide grouping. The compounds disclosed in WO 89/07 443 in the form of optically pure R(-)-niguldipin and further analogous dihydropyridines with cytotoxic activity have larger structural differences. However, as compared to these known compounds, the compounds according to the invention unexpectedly possess a better activity and a wider spectrum of action despite the large structural difference.

In the international PCT patent applications WO 95/10516, WO 96/31477, WO 96/31478 or for example in WO 95/10515, tricyclic amide compounds are described which possess an anti-proliferative activity. All of these compounds described therein are distinguished in that they must imperatively possess a tricyclic anellated ring system with at least one nitrogen atom, for example 6, 11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridinyl ring system as a pharmacophoric group, whereby the molecule portion opposite this tricyclic anellated system is uncommonly variable and represents one of many variation possibilities among numerous substitution possibilities of the given pyridyl substitution. A further meaningful difference in the substitution of these molecules in comparison to the compounds according to the invention is to be seen in the lack of the present structural element D, i.e. both heterocycles found in opposition are directly bound over the carboxy group without being bound over an alkaline chain found there.

A further essential difference of the compounds according to the invention in comparison to these tricyclic compounds is to be recognized in the presence of the terminal 3-pyridyl-substitution which must be present. The presence of this heterocyclic ring required according to the invention as well as the particular bond in the 3-position according to the substitution of the invention in comparison to the above

mentioned anti-proliferative compounds of the state of the art indicates that this 3-pyridyl group is an important factor for the anti-tumor action.

In fact, the compounds according to the invention cover a different tumor spectrum from those named in the PCT/WO publications with this necessarily present tricyclic anellated ring system. In the mentioned PCT/WO publications of the state of the art, a treatment possibility in tumors is merely mentioned which is made in connection with a potential inhibition of the farnesyl protein transferase, whereby this mechanism relates to the expression of the activated ras-oncogene. In contrast to this, the compounds according to the invention with the 3-pyridyl-substitution required according to the invention are not limited to the therapy of tumor cells of this type with abnormal production of the ras-oncogene; rather, the therapy possibilities with the new compounds according to the invention extend to the combat of numerous other types of tumors with different causal mechanisms as well as immunosuppressive treatment possibilities such as autoimmune diseases.

In view of this art, the finding that the compounds according to the general formula (I) with the particular substitutions defined below have superior pharmacological activities which make them particularly suitable in an excellent manner for the therapy of abnormal cell growth such as tumor illnesses over a broad anti-proliferative spectrum, was completely unexpected. The pharmacological finding that, aside from the cytostatic effectiveness, especially with different tumor spectra, the compounds according to the invention also possess immunosuppressive properties and additionally favorable abortive properties without harmful mutagenic effects is to be considered as equally surprising.

Structurally close pyridyl compounds wherein instead of the piperazine ring, a cyclic non-aromatic heterocyclic ring residue with merely one ring nitrogen atom and optionally an


ORIGINAL RECEIVED




additional ring oxygen atom, preferably a piperidinyl residue, is incorporated as well as their use especially as cytostatic agents are subject matter of the older patent applications P 196 24 704.7-44, P 196 24 668.7-41 as well as P 196 24 659.8-44 which have not yet been published.

The most important differentiating feature of the new compounds according to the invention with respect to these older, non-prepublished compounds is therefore the structural feature E which in the present application always represents especially piperazine or hexahydro-1,4-diazepine.

These new piperazinyl-substituted pyridyl carboxamides correspond to the following general formula:



wherein the structural element E has the following meaning:



whereby  $q$  is 1, 2 or 3 and therewith can represent the N-heterocyclic ring piperazine, hexahydro-1,4-diazepine or octahydro-1,4-azocine.

The meaning of the remaining substituents and the preferred embodiments of the compound groups according to the invention falling under the general formula as well as particularly preferred end products are defined in claims 1 to 7 in detail.

The compounds of Formula (I), which represent the end products can optionally exist as cis- and trans-isomers, E- and Z-isomers, for example when A is a cyclopropane ring or D contains one or more double bonds. Subject matter of the invention is the pure isomers as well as their mixtures. Furthermore, the compounds of Formula (I) can contain one or more asymmetric carbon atoms and, as a result, exist in the form of different optical isomers (enantiomers, diastereomers). The invention includes all optical isomers and their racemic or non-racemic mixtures. Finally, compounds of Formula (I) can exist as endo/exo-isomers in case the ring system E is bicyclic. The pure endo- and exo-isomers as well as their mixtures are also comprised by the invention.

Compounds of Formula (I), in which G is a heterocyclic aromatic ring or contains such in an anellated ring system can optionally be present as tautomers when this heterocyclic ring is substituted by free hydroxy-, mercapto- or amino groups. In this case, the invention includes all tautomeric forms.

Subject matter of the invention are further pharmacologically acceptable acid addition salts of the compounds of Formula (I) with inorganic or organic acids. Preferable examples for addition salts with suitable inorganic acids are hydrochlorides, hydrobromides, hydroiodides, sulfates and phosphates. Addition salts of organic acids are preferably acetates, benzoates, citrates, fumarates, gluconates, malates, maleates, methanesulfonates, lactates, oxalates, succinates, tartrates and tosylates.

00596001.061600

Compounds of Formula (I) as well as their acid addition salts can also be optionally present as hydrates or other solvates. The invention includes such hydrates and solvates.

In the compounds of Formula (I), the definitions for the atoms or atomic groups preferably have the following meanings:

Halogen means fluorine, chlorine, bromine or iodine;

Alkyl can be straight chained or branched and preferably signifies a  $C_1$ - $C_6$ -alkyl residue, especially a methyl-, ethyl-, propyl-, isopropyl-, butyl-, isobutyl-, sec-butyl-, tert-butyl-, cyclopropylmethyl-, pentyl-, isopentyl-, tert-pentyl-, neopentyl-, cyclopropylethyl-, cyclobutylmethyl- or hexyl group.

Alkylene signifies for example methylene, ethylene, propylene, tetramethylene, pentamethylene, hexamethylene, heptamethylene, octamethylene, nonamethylene or decamethylene.

Alkenyl preferably signifies  $C_3$ - $C_6$ -alkenyl and can be straight chained or branched and preferably signifies an allyl-, 2-butenyl-, 3-butenyl-, 2-methyl-2-propenyl-, 2-pentenyl-, 4-pentenyl-, 2-methyl-2-butenyl-, 3-methyl-2-butenyl-, 2-hexenyl-, 5-hexenyl-, 4-methyl-3-pentenyl- or 2,2-dimethyl-3-butenyl-group.

Alkenylene signifies for example ethenylene, propenylene, butenylene, pentenylene, hexenylene, hexadienylene, heptenylene, octenylene, nonenylene or decenylene.

Alkynyl preferably signifies  $C_2$ - $C_6$ -alkynyl which can be straight chained or branched and can preferably signify an ethynyl-, propargyl-, 2-butylnyl-, 3-butylnyl-, 4-pentylnyl-, 5-hexynyl- or 4-methyl-2-pentylnyl group.

0596001-067870

Alkinylene signifies for example propinylene, butinylene, pentinylene, hexinylene, hexeninylene, heptinylene, octinylene, noninylene or decinylene.

Cycloalkyl is preferably a C<sub>3</sub>-C<sub>8</sub>-cycloalkyl residue, especially a cyclopropyl-, cyclobutyl-, cyclopentyl-, cyclohexyl-, cycloheptyl- or cyclooctyl group.

Hydroxyalkyl contains a hydroxyl group in one of the above mentioned alkyl residues, especially in a C<sub>1</sub>-C<sub>6</sub>-alkyl residue, whereby among the C<sub>1</sub>-C<sub>6</sub>-hydroxyalkyl residues, the hydroxymethyl- and the hydroxyethyl residue are preferred.

Aside from the oxygen atom, alkoxy, alkenyloxy, alkinyloxy contain one of the above mentioned preferred C<sub>1</sub>-C<sub>6</sub>-alkyl-, C<sub>3</sub>-C<sub>6</sub>-alkenyl- and/or C<sub>3</sub>-C<sub>6</sub>-Alkynyl groups. Particularly preferred groups for this are the methoxy-, ethoxy-, isopropoxy-, tert-butoxy-, allyloxy- and propargyloxy groups.

Alkoxy, especially C<sub>1</sub>-C<sub>6</sub>-alkoxy, entirely or partially replaced by fluorine is for example difluoromethoxy, trifluoromethoxy or 2,2,2-trifluoroethoxy.

Aside from the sulphur atom, alkylthio, alkenylthio, alkynylthio contain one of the above mentioned preferred C<sub>1</sub>-C<sub>6</sub>-alkyl-, C<sub>3</sub>-C<sub>6</sub>-alkenyl- or C<sub>3</sub>-C<sub>6</sub>-alkynyl groups. Preferred groups among these are the methylthio-, ethylthio-, isopropylthio- and tert-butylthio groups.

Cyclopentyloxy- and cyclopentylthio- and/or cyclohexyloxy- and cyclohexylthio residues represent preferred C<sub>3</sub>-C<sub>8</sub>-cycloalkyloxy and C<sub>3</sub>-C<sub>8</sub>-cycloalkylthio.

Aside from the oxygen atom, alkanoyloxy groups preferably contain an aliphatic acyl group with 1 to 7 carbon atoms.

Among preferred alkanoyloxy groups are the acetoxy-, propionyloxy- and pivaloyloxy groups.

Alkoxy carbonyl groups, preferably C<sub>2</sub>-C<sub>7</sub>-alkoxy carbonyl groups contain, aside from the carbonyl group, one of the above mentioned alkoxy groups, especially C<sub>1</sub>-C<sub>6</sub>-alkoxy groups. Preferred alkoxy carbonyl groups are the methoxy carbonyl-, ethoxy carbonyl-, isopropoxy carbonyl-, isobutoxy carbonyl- and tert-butoxy carbonyl groups.

Aside from the oxygen atom, alkoxy carbonyloxy groups preferably contain one of the above mentioned C<sub>2</sub>-C<sub>7</sub>-alkoxy carbonyl residues. Among preferred alkoxy carbonyl groups are the methoxy carbonyloxy-, ethoxy carbonyloxy-, isopropoxy carbonyloxy-, isobutoxy carbonyloxy- and tert-butoxy carbonyl groups as well as allyloxy carbonyloxy groups.

Aside from the carbonyl group, alkylaminocarbonyl, especially C<sub>2</sub>-C<sub>7</sub>-alkylaminocarbonyl and dialkylaminocarbonyl groups, preferably C<sub>3</sub>-C<sub>13</sub>-dialkylaminocarbonyl groups, contain an alkylamino- and/or dialkylamino residue whose alkyl groups correspond especially to the C<sub>1</sub>-C<sub>6</sub>-alkyl groups of the above description. Preferred groups are the dimethylaminocarbonyl-, diethylaminocarbonyl- and diisopropylamino-carbonyl groups.

Aside from the unsubstituted amino group, the amino group of the Formula  $NR^5R^6$  is one of the below mentioned alkylamino groups, especially C<sub>1</sub>-C<sub>6</sub>-alkylamino groups and/or dialkyl-amino groups, especially di-(C<sub>1</sub>-C<sub>6</sub>-alkyl)amino groups.

Alkylamino especially contains one of the above mentioned C<sub>1</sub>-C<sub>6</sub>-alkyl groups. Preferred groups are the methylamino-, ethylamino-, propylamino-, isopropylamino-, butylamino-, and the tert-butylamino groups.

The preferred di-(C<sub>1</sub>-C<sub>6</sub>-alkyl)amino residue carries two of the same or different of the above mentioned C<sub>1</sub>-C<sub>6</sub>-alkyl groups on the nitrogen atom. Preferred groups are the dimethylamino-, diethylamino-, dipropylamino-, diisopropylamino-, isopropyl-, methylamino-, dibutylamino- or tert-butylmethylamino groups.

Acyl, especially C<sub>1</sub>-C<sub>6</sub>-acyl, signifies the residue of an aliphatic saturated or unsaturated, straight chained, branched or cyclic carboxylic acid. Preferred acyl residues are formyl-, acetyl-, propionyl-, acryloyl-, butyryl-, isobutyryl-, methacryloyl-, cyclopropylcarbonyl-, pentanoyl-, pivaloyl-, cyclobutylcarbonyl-, hexanoyl- and dimethylacryloyl groups.

Alkanesulfonyl, especially C<sub>1</sub>-C<sub>6</sub>-alkanesulfonyl is preferably the methanesulfonyl-, ethanesulfonyl-, propanesulfonyl-, butanesulfonyl-, pentanesulfonyl- or hexanesulfonyl groups.

Saturated or unsaturated, preferably four- to eight-membered heterocycles with one or two hetero-atoms, are for example azetidine, tetrahydrofuran, tetrahydrothiophene, pyrrolidine, tetrahydropyran, tetrahydropyridine, piperidine, tetrahydroazepine, hexahydroazepine, octahydroazocine, pyrazolidine, piperazine, morpholine, thiomorpholine, thiomorpholin-1,1-dioxide, hexahydrodiazepine or hexahydrooxazepine.

Preferred monocyclic aromatic five- or six-membered heterocycles with one to three hetero-atoms are for example furyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, pyrazolyl, oxadiazolyl, thiadiazolyl, triazolyl, pyridyl, pyrazinyl, pyridazinyl, pyrimidinyl or triazinyl.

Anellated bi- and tricyclic aromatic or partially hydrated carbocyclic ring systems with 8 to 16 ring atoms and at least

one aromatic ring are preferably benzocyclobutyl, indanyl, indenyl, naphthyl, dihydronaphthyl, tetrahydronaphthyl, biphenylenyl, fluoroenyl, anthryl, dihydroanthryl, phenanthryl, dihydrophenanthryl, dibenzocycloheptenyl, dihydrodibenzocycloheptenyl, dihydrodibenzocyclooctenyl or tetrahydrodibenzocyclooctenyl. Their mono- or dioxo-derivates, i.e. for example the residues of indanone, tetralone, anthrone, anthraquinone, fluoroenone, phenanthrone, dibenzocycloheptenone, dihydrodibenzocycloheptenone or tetrahydrodibenzocyclooctenone are also to be understood as partially hydrated carbocyclic ring systems.

Anellated bi- and tricyclic aromatic or partially hydrated heterocyclic ring systems with 8 to 16 ring atoms and at least one aromatic ring are, for example, imidazothiazolyl, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl, indolinyl, isoindolinyl, benzimidazolyl, indazolyl, benzoxazolyl, benzoisoxazolyl, benzothiazolyl, benzoisothiazolyl, benzofurazanyl, benzothiadiazolyl, benzotriazolyl, oxazolopyridyl, thiazolopyridyl, isothiazolopyridyl, imidazopyridyl, pyrazolopyridyl, thienopyrimidinyl, chromanyl, benzopyranyl, quinolyl, isoquinolyl, dihydroquinolyl, tetrahydroquinolyl, benzodioxanyl, quinoxalanyl, quinazolinyl, naphthyridinyl, carbazolyl, tetrahydrocarbazolyl, pyridoindolyl, acridinyl, phenanthridinyl, dihydrophenanthridinyl, dibenzoisoquinolinyl, dihydrodibenzoisquinolinyl, phenothiazinyl, dihydrodibenzoxepinyl, benzocycloheptathienyl, dihydrothienobenzothiepinyl, dihydrodibenzothiepinyl, octahydrodibenzothiepinyl, dibenzoazepinyl, dihydrodibenzazepinyl, octahydrodibenzazepinyl, benzocycloheptapyridyl, dihydrobenzocycloheptapyridyl, pyridobenzoazepinyl, dihydropyridobenzoazepinyl, dihydropyridobenzodiazepinyl, dihydrodibenzoxazepinyl, dihydropyridobenzoxazepinyl, dihydrodibenzothiazepinyl or dihydropyridobenzothiazepinyl.

Furthermore, their mono- or dioxo-derivates and/or optionally their possible tautomers are also to be understood as partially hydrated heterocyclic ring systems, i.e. for example the residues of indolinone, isatin, of benzooxazolone and/or its tautomer hydroxybenzooxazole, of benzisoxazolone, benzothiazolone, benzoisothiazolone and benzimidazolone and/or their corresponding tautomers, hydroxybenzoisoxazole, hydroxybenzothiazole, hydroxybenzoisothiazole and hydroxybenzimidazole, as well as indazolinone, of oxazolopyridinones, thiazolopyridinones, pyrazolopyridinones and imidazopyridinones and/or their corresponding tautomers hydroxyoxazolopyridine, hydroxythiazolopyridine, hydroxypyrazolopyridine and hydroxyimidazopyridine, the residues from the series chromanone, chromone, quinolinone, dihydroquinolinone, tetrahydrocarbazonone, acridone, phenanthridone, benzoisoquinolinone, dihydrodibenzoxepinones, benzocycloheptathiophenones, dihydrothienobenzothiepinones, dihydrodibenzothiepinones, dihydrodibenzoazepinones, benzocycloheptapyridinones, dihydropyridobenzoazepinones, dihydropyridobenzodiazepinones, dihydropyridobenzoazepinones, dihydrodibenzothiazepinones and of dihydropyridobenzothiazepinones.

Saturated and unsaturated monocyclic, four- to eight-membered heterocycles (as the group  $\text{—NR}^{13}\text{R}^{15}$ ) which, aside from the essential nitrogen atom, can optionally contain one or two further hetero-atoms selected from N and/or S and/or O, are for example azetidine, pyrrolidine, piperidine, (1H)-tetrahydropyridine, hexahydroazepine, (1H)-tetrahydroazepine, octahydroazocine, pyrazolidine, piperazine, hexahydrodiazepine, morpholine, hexahydroazepine, thiomorpholine or thiomorpholin-1,1-dioxide.


Saturated or unsaturated bi- or tricyclic, anellated or bridged heterocycles with 8 to 16 ring atoms (as the group  $\text{—NR}^{13}\text{R}^{15}$ ) which, aside from the essential nitrogen atom, can optionally contain one or two further hetero-atoms, selected

09596001-061600



from N and/or S and/or O, are for example 5-aza-bicyclo[2.1.1]hexane, 2-aza-bicyclo[2.2.1]heptane, 7-aza-bicyclo[2.2.1]heptane, 2,5-diaza-bicyclo[2.2.1]heptane, 2-aza-bicyclo[2.2.2]octane, 8-aza-bicyclo[3.2.1]octane, 2,5-diaza-bicyclo[2.2.2]octane, 9-aza-bicyclo[3.3.1]nonane, indoline, isoindoline, (1H)-dihydroquinoline, (1H)-tetrahydroquinoline, (2H)-tetrahydroisoquinoline, (1H)-tetrahydroquinoxaline, (4H)-dihydrobenzoxazine, (4H)-dihydrobenothiazine, (1H)-tetrahydrobenzo[b]azepine, (1H)-tetrahydrobenzo[c]azepine, (1H)-tetrahydrobenzo[d]azepine, (5H)-tetrahydrobenzo[b]oxazepine, (5H)-tetrahydrobenzo[b]thiazepine, 1,2,3,4-tetrahydro-9H-pyrido[3,4-b]indole, (10H)-dihydroacridine, (10H)-dihydrophenanthridine, 1,2,3,4-tetrahydroacridanone, (10H)-phenoxazine, (10H)-phenothiazine, (5H)-dibenzazepine, (5H)-dihydrodibenzazepine, (5H)-octahydrodibenzazepine, dihydrobenzo[d,e]isoquinoline, (5H)-dihydrodibenzodiazepine, (5H)-benzo[b]pyrido[f]azepine, (5H)-dihydrobenzo[b]pyrido-[f]azepine (11H)-dihydrodibenzo[b,e]oxazepine, (11H)-dihydrodibenzo[b,e]thiazepine, (10H)-dihydrodibenzo[b,f]oxazepine, (10H)-dihydrodibenzo[b,f]thiazepine, (5H)-tetrahydrodibenzazocine, (11H)-dihydrobenzo[e]pyrido[b]-1,4-diazepin-6-one, (11H)-di-hydrobenzo[b]pyrido[e]-1,4-diazepin-5-one.

Concretely, the invention relates to new piperazinyl-substituted compounds of the general Formula (I)



(I)

wherein  
 $R^1$  is selected from  
 hydrogen, hydroxy, halogen, cyano, aminocarbonyl,  
 carboxy,

saturated, single or several-fold unsaturated, branched  
 or straight chained or cyclic hydrocarbon residues such  
 as alkyl, alkenyl, alkynyl or cycloalkyl,

aryl such as phenyl or heteroaryl such as pyridyl,

alkoxy, cycloalkyloxy, alkenyloxy or alkinyloxy or  
 aralkyloxy such as the benzyloxy group, alkoxycarbonyl,  
 alkylaminocarbonyl, dialkylaminocarbonyl, alkanoyloxy,  
 alkoxycarbonyloxy, alkylthio, cycloalkylthio,  
 alkenylthio, alkynylthio, aryloxy such as phenoxy,  
 heteroaryloxy such as pyridyloxy, arylthio such as  
 phenylthio, heteroarylthio such as pyridylthio,

trifluoromethyl,

hydroxyalkyl,

NR<sup>5</sup>R<sup>6</sup>, wherein

R<sup>5</sup> and R<sup>6</sup> are selected independent of each other from hydrogen, saturated or unsaturated hydrocarbon residues such as alkyl, alkenyl, alkynyl, or aryl such as phenyl and aralkyl such as benzyl;

R<sup>2</sup> is selected from hydrogen, halogen, cyano, saturated hydrocarbon residues such as alkyl, or halogenated hydrocarbon residues such as trifluoromethyl, hydroxy, alkoxy, aralkyloxy residues such as benzyloxy, as well as alkanoyloxy,

whereby R<sup>1</sup> and R<sup>2</sup>, in the case that they are immediately adjacent to each other, optionally form a bridge which is selected from

-(CH<sub>2</sub>)<sub>4</sub>- and -(CH=CH)<sub>2</sub>- and -CH<sub>2</sub>O-CR<sup>7</sup>R<sup>8</sup>-O-, wherein

R<sup>7</sup> and R<sup>8</sup> are selected independently of each other from hydrogen and alkyl residues;

R<sup>3</sup> is selected from hydrogen, halogen, saturated hydrocarbon residues such as alkyl, or halogenated hydrocarbon residues such as trifluoromethyl, or hydroxyalkyl;

R<sup>4</sup> is selected from hydrogen, hydroxy, or single or several-fold unsaturated, branched or straight chained or cyclic hydrocarbon residues such as alkyl, alkenyl, alkynyl or cycloalkyl, alkoxy and aralkyloxy such as benzyloxy;

k is 0 or 1;

A is selected from

Alkylene which is optionally substituted one to three-fold by straight chained or branched chained hydrocarbon residues such as alkyl, hydroxy, alkoxy, halogen such as fluorine, or aryl such as phenyl,

Alkylene, wherein a methylene unit is isosterically replaced by O, S,  $\text{NR}^9$ , CO, SO or  $\text{SO}_2$  whereby, with the exception of CO, the isosteric substitution cannot be adjacent to the amide group and, in  $\text{NR}^9$ , the residue  $\text{R}^9$  is selected from hydrogen, alkyl, alkenyl, alkynyl, acyl or alkanesulfonyl;

Cycloalkylene such as 1,2-cyclopropylene;

Alkenylene which is optionally substituted one to three-fold by alkyl, hydroxy, alkoxy, fluorine, cyano or aryl such as phenyl;

Alkadienylene, which is optionally substituted once or twice by alkyl, fluorine, cyano or aryl such as phenyl, 1,3,5-hexatrienylene, which is optionally substituted by alkyl, fluorine, cyano or aryl such as phenyl, and

ethynylene;

D is selected from alkylene, which is optionally substituted once or twice by alkyl, hydroxy, or alkoxy;

alkenylene, which is optionally substituted once or twice by alkyl, hydroxy, or alkoxy;


alkynylene, which is optionally substituted once or twice by alkyl, hydroxy, or alkoxy, as well as

0056607.061600

alkylene, alkenylene or alkynylene, wherein one to three methylene units is each isosterically replaced by O, S, NR<sup>10</sup>, CO, SO or SO<sub>2</sub>, wherein

R<sup>10</sup> has the same meaning as R<sup>9</sup> but is selected independently thereof;

E



whereby

q is 1, 2 or 3;


R<sup>11</sup> is selected from hydrogen, alkyl, hydroxy, hydroxymethyl, carboxy, or alkoxycarbonyl and

R<sup>12</sup> is selected from hydrogen, alkyl or an oxo group immediately adjacent to a nitrogen atom or

R<sup>11</sup> and R<sup>12</sup>, optionally together, form an alkylene bridge under formation of a bicyclic ring system;

G is selected from G<sup>1</sup>, G<sup>2</sup>, G<sup>3</sup>, G<sup>4</sup> or G<sup>5</sup>, wherein

G<sup>1</sup> is



(G<sup>1</sup>)

whereby

r has the meaning 0 to 3,

s is 0 or 1;

R<sup>13</sup> is selected from  
hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl;  
saturated or unsaturated, four to eight-membered  
heterocycles which can contain one or two hetero-atoms  
that are selected from N and/or S and/or O;  
benzyl, phenyl;  
monocyclic aromatic five- or six-membered heterocycles  
which can contain 1 to 3 hetero-atoms that are selected  
from N and/or S and/or O and are either directly bound  
or bound over a methylene group;

anellated bi- and tricyclic aromatic or partially  
hydrated carbocyclic ring systems with 8 to 16 ring  
atoms and at least one aromatic ring, whereby the bond  
can occur either over an aromatic or a hydrated ring and  
either directly or over a methylene group;

anellated bi- and tricyclic aromatic or partially  
hydrated heterocyclic ring systems with 8 to 16 ring  
atoms and at least one aromatic ring, whereby one to  
three ring atoms can be selected from N and/or S and/or  
O and the bond can occur either over an aromatic or a  
hydrated ring and either directly or over a methylene  
group;

R<sup>14</sup> has the same meaning as R<sup>13</sup> but is selected  
independently thereof;


R<sup>15</sup> is selected from  
hydrogen, hydroxy, C<sub>1</sub>-C<sub>3</sub>-alkyl, aralkyl such as benzyl  
or aryl such as phenyl,

monocyclic aromatic five or six-membered heterocycles, which can contain one to three hetero-atoms selected from the group N and/or S and/or O and are either bound directly or over a methylene group,

annelated bi- and tricyclic aromatic or partially hydrated carbocyclic ring systems with 8 to 16 ring atoms and at least one aromatic ring, wherein the linkage can occur either over an aromatic or a hydrated ring and either directly or over a methylene group,

annelated bi- and tricyclic aromatic or partially hydrated heterocyclic ring systems with 8 to 16 ring atoms and at least one aromatic ring, wherein one to three ring atoms can be selected from N and/or S and/or O and the linkage can occur either over an aromatic ring or a hydrated ring and either directly or over a methylene group,

with the exception of compounds in which G has the meaning




in the case that the following substituents are simultaneously signify


- |                 |  |
|-----------------|--|
| R <sup>13</sup> | pyridyl or (optionally halogen-, alkyl-, alkoxy- or Trifluoromethyl-substituted) phenyl,       |
| R <sup>14</sup> | hydrogen or (phenyl optionally substituted with halogen-, alkyl-, alkoxy- or Trifluoro methyl, |
| R <sup>15</sup> | is hydrogen, and   |
| A               | represents alkylene, optionally substituted ethenylene or butadienylene,                       |

D       alkylene or alkenylene as well as  
 E       piperazine or homopiperazine and  
 S = 1;

G<sup>2</sup>   is selected from



or



whereby r and s as well as the substituents R<sup>13</sup> to R<sup>15</sup> can have the above meaning, or the grouping



can also be a nitrogen heterocycle bound over the nitrogen atom selected from

saturated or unsaturated monocyclic, four to eight-membered heterocycles, which, aside from the essential nitrogen atom, can still optionally contain one or two further hetero-atoms selected from N and/or S and/or O, or

saturated or unsaturated bi- or tricyclic, anellated or bridged heterocycles with 8 to 16 ring atoms, that aside from the essential nitrogen atom, can optionally still contain one or two further hetero-atoms that are selected from N and/or S and/or O;

G<sup>3</sup>   has the meaning  $\text{---} \text{SO}_2 \text{---} (\text{CH}_2)_r \text{---} \text{R}^{13}$  (G3)

wherein r and R<sup>13</sup> have the above definition,



G<sup>4</sup> has the meaning



(G4)

whereby

Ar<sup>1</sup> and Ar<sup>2</sup> can be selected independently from each other from phenyl, pyridyl or naphthyl;

G<sup>5</sup> has the meaning



(G5)

whereby

R<sup>16</sup> is selected from trifluoromethyl, alkoxy, alkenyloxy, and aralkyloxy such as benzyloxy,

whereby aromatic ring systems in the substituents R<sup>1</sup>, R<sup>2</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>13</sup>, R<sup>14</sup>, R<sup>15</sup>, R<sup>16</sup>, Ar<sup>1</sup> and Ar<sup>2</sup> and/or in the ring system  $\text{---NR}^{13}\text{R}^{15}$  can be substituted independently from each other by one to three of the same or different groups selected from

halogen, cyano, alkyl, halogen alkyl such as trifluoromethyl, cycloalkyl, aryl such as phenyl, arylalkyl such as benzyl; hydroxy, hydroxy alkyl, alkoxy, alkoxy entirely or partially substituted by fluorine, aralkyloxy such as benzyloxy, aryloxy such as phenoxy; mercapto, alkylthio, carboxy, carboxyalkyl, carboxyalkenyl, alkoxycarbonyl, aralkyloxycarbonyl such as benzyloxycarbonyl, nitro, amino, monoalkylamino, dialkylamino and in the case of two adjacent residues on the aromatic ring, also methylenedioxy, and


whereby alkyl-, alkenyl- and cycloalkyl residues in the groups  $G^1$ ,  $G^2$  and  $G^3$  can be substituted by one or two of the same or different groups which are selected from hydroxy, carboxy, alkoxycarbonyl, aralkyloxycarbonyl such as benzyloxycarbonyl, amino, monoalkylamino and dialkylamino;

their cis- and trans-isomers, E- and Z-isomers, especially in case that A is a cyclopropane ring or D contains one or more double bonds, including the enantiomers, diastereomers and other isomers as well as their racemic or non-racemic mixtures and the corresponding endo- and exo-isomers for the case that the ring system E is bicyclic;

their tautomers;

their acid addition salts including their hydrates and solvates.

Furthermore, the invention relates to preferred pyridylalkane, pyridylalkene and pyridylalkine carboxamides of the Formula (I)



(I)

wherein the substituents have the following meanings:

R<sup>1</sup> is selected from hydrogen, halogen, cyano, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-alkinyl, trifluoromethyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl, C<sub>1</sub>-C<sub>6</sub>-hydroxyalkyl, hydroxy, C<sub>1</sub>-C<sub>6</sub>-alkoxy, C<sub>3</sub>-C<sub>6</sub>-alkenyloxy, C<sub>3</sub>-C<sub>6</sub>-alkinyloxy, benzyloxy, C<sub>1</sub>-C<sub>7</sub>-alkanoyloxy, C<sub>2</sub>-C<sub>7</sub>-alkoxycarbonyloxy, C<sub>1</sub>-C<sub>6</sub>-alkylthio, C<sub>3</sub>-C<sub>6</sub>-alkenylthio, C<sub>3</sub>-C<sub>6</sub>-alkinylthio, C<sub>3</sub>-C<sub>8</sub>-cycloalkyloxy, C<sub>3</sub>-C<sub>8</sub>-cycloalkylthio, C<sub>2</sub>-C<sub>7</sub>-alkoxycarbonyl, aminocarbonyl, C<sub>2</sub>-C<sub>7</sub>-alkylaminocarbonyl, C<sub>3</sub>-C<sub>13</sub>-dialkylaminocarbonyl, carboxy, phenyl, phenoxy, phenylthio, pyridyloxy, pyridylthio, and NR<sup>5</sup>R<sup>6</sup>, wherein

R<sup>5</sup> and R<sup>6</sup> are selected independently of each other from hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-alkenyl, C<sub>3</sub>-C<sub>6</sub>-alkinyl, benzyl and phenyl;

R<sup>2</sup> is selected from hydrogen, halogen, cyano, C<sub>1</sub>-C<sub>6</sub>-alkyl, trifluoromethyl, hydroxy, C<sub>1</sub>-C<sub>6</sub>-alkoxy, benzyloxy and C<sub>1</sub>-C<sub>7</sub>-alkanoyloxy; R<sup>1</sup> and R<sup>2</sup>, if adjacent, optionally form a bridge selected from - (CH<sub>2</sub>)<sub>4</sub> - and - (CH=CH)<sub>2</sub> - or -CH<sub>2</sub>O-CR<sup>7</sup>R<sup>8</sup>-O-, wherein

R<sup>7</sup> and R<sup>8</sup> are selected independently from each other from hydrogen and C<sub>1</sub>-C<sub>6</sub>-alkyl;

R<sup>3</sup> is selected from hydrogen, halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, trifluoromethyl and C<sub>1</sub>-C<sub>6</sub>-hydroxyalkyl;

R<sup>4</sup> is selected from hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-alkenyl, C<sub>3</sub>-C<sub>6</sub>-alkinyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, hydroxy, C<sub>1</sub>-C<sub>6</sub>-alkoxy and benzyloxy;

A is selected from  
C<sub>1</sub>-C<sub>6</sub>-alkylene, optionally substituted one to three-fold  
by C<sub>1</sub>-C<sub>3</sub>-alkyl, hydroxy, C<sub>1</sub>-C<sub>3</sub>-alkoxy, fluorine, or  
phenyl,

C<sub>2</sub>-C<sub>6</sub>-alkylene, in which a methylene unit is isosterically replaced by O, S, NR<sup>9</sup>, CO, SO or SO<sub>2</sub>, whereby, with the exception of CO, the isosteric substitution cannot be adjacent to the amide group and, in NR<sup>9</sup>, the residue R<sup>9</sup> is selected from hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-alkenyl, C<sub>3</sub>-C<sub>6</sub>-alkinyl, C<sub>1</sub>-C<sub>6</sub>-acyl or C<sub>1</sub>-C<sub>6</sub>-alkanesulfonyl,

1,2-cyclopropylene,

C<sub>2</sub>-C<sub>6</sub>-Alkenylene, optionally substituted once to three-fold by C<sub>1</sub>-C<sub>3</sub>-alkyl, hydroxy, C<sub>1</sub>-C<sub>3</sub>-alkoxy, fluorine, cyano or phenyl,

C<sub>4</sub>-C<sub>6</sub>-alkadienylene, optionally substituted once or twice by C<sub>1</sub>-C<sub>3</sub>-alkyl, fluorine, cyano or phenyl;

1,3,5-hexatrienylene, optionally substituted by C<sub>1</sub>-C<sub>3</sub>-alkyl, fluorine, cyano or phenyl, and

ethinylene


D is selected from  
C<sub>2</sub>-C<sub>10</sub>-alkylene, optionally substituted once or twice by  
C<sub>1</sub>-C<sub>6</sub>-alkyl, hydroxy, or C<sub>1</sub>-C<sub>6</sub>-alkoxy;

C<sub>4</sub>-C<sub>10</sub>-alkenylene, optionally substituted once or twice by C<sub>1</sub>-C<sub>6</sub>-alkyl, hydroxy, or C<sub>1</sub>-C<sub>6</sub>-alkoxy;

C<sub>4</sub>-C<sub>10</sub>-alkynylene, optionally substituted once or twice by C<sub>1</sub>-C<sub>6</sub>-alkyl, hydroxy, or C<sub>1</sub>-C<sub>6</sub>-alkoxy; as well as

C<sub>2</sub>-C<sub>10</sub>-alkylene, C<sub>4</sub>-C<sub>10</sub>-alkenylene or C<sub>4</sub>-C<sub>10</sub>-alkynylene, in which one to three methylene units are isosterically replaced by O, S, NR<sup>10</sup>, CO, SO or SO<sub>2</sub>, whereby R<sup>10</sup> has the same meaning as R<sup>9</sup>, but is selected independently thereof;

E signifies



whereby

q has the meaning 1, 2 or 3;

R<sup>11</sup> is selected from hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, hydroxy, hydroxymethyl, carboxy, or C<sub>2</sub>-C<sub>7</sub>-alkoxycarbonyl and

R<sup>12</sup> is selected from hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl or an oxo group adjacent to a nitrogen atom, or R<sup>11</sup> and R<sup>12</sup> optionally together, form a C<sub>1</sub>-C<sub>3</sub>-alkylene bridge under formation of a bicyclic ring system;

G is selected from G<sub>1</sub>, G<sub>2</sub>, G<sub>3</sub>, G<sub>4</sub> or G<sub>5</sub>, whereby

G<sup>1</sup> represents  $\text{---}(\text{CH}_2)_r\text{---}(\text{CR}^{14}\text{R}^{15})_s\text{---R}^{13}$  (G1)

r has the meaning 0 to 3,

s is 0 or 1 and

R<sup>13</sup> is selected from

hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-alkenyl, C<sub>3</sub>-C<sub>6</sub>-alkinyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl;

saturated or unsaturated, four to eight-membered heterocycles, which can contain one or two hetero-atoms that are selected from N and/or S and/or O; benzyl, phenyl;

monocyclic aromatic five or six-membered heterocycles, which can contain one to three hetero-atoms selected from the group N and/or S and/or O and are either bound directly or over a methylene group,

annelated bi- and tricyclic aromatic or partially hydrated carbocyclic ring systems with 8 to 16 ring atoms and at least one aromatic ring, wherein the linkage can occur either over an aromatic or a hydrated ring and either directly or over a methylene group,

annelated bi- and tricyclic aromatic or partially hydrated heterocyclic ring systems with 8 to 16 ring atoms and at least one aromatic ring, wherein one to three ring atoms can be selected from N and/or S and/or O and the linkage can occur either over an aromatic ring or a hydrated ring and either directly or over a methylene group,

R<sup>14</sup> has the same meaning as R<sup>13</sup>, but is selected independently thereof;


R<sup>15</sup> is selected from  
hydrogen, hydroxy, methyl, benzyl, phenyl,

monocyclic aromatic five or six-membered heterocycles, which can contain one to three hetero-atoms from the group N and/or S and/or O and are either bound directly or over a methylene group,

annelated bi- and tricyclic aromatic or partially hydrated carbocyclic ring systems with 8 to 16 ring atoms and at least one aromatic ring, wherein the linkage can occur either over an aromatic or a hydrated ring and either directly or over a methylene group,

annelated bi- and tricyclic aromatic or partially hydrated heterocyclic ring systems with 8 to 16 ring atoms and at least one aromatic ring, wherein one to three ring atoms can be selected from N and/or S and/or O and the linkage can occur either over an aromatic ring or a hydrated ring and either directly or over a methylene group,


whereby G in the form of G<sup>1</sup> cannot have the meaning




in the case that the following substitutions simultaneously signify

- R<sup>13</sup> pyridyl or (optionally halogen-, alkyl-, alkoxy- or Trifluoromethyl- substituted) phenyl,
- R<sup>14</sup> hydrogen or (phenyl optionally substituted with halogen-, alkyl-, alkoxy- or trifluoromethyl,
- R<sup>15</sup> is hydrogen, and
- A represents alkylene, optionally substituted ethenylene or butadienylene,
- D alkylene or alkenylene as well as
- E piperazine or homopiperazine and
- s = 1;

G<sup>2</sup> is selected from



or



whereby r and s as well as the substituents R<sup>13</sup> to R<sup>15</sup> can have the above meaning, or the grouping



can also be a nitrogen heterocycle bound over the nitrogen atom selected from

saturated or unsaturated monocyclic, four to eight-membered heterocycles, which, aside from the essential nitrogen atom, can optionally still contain one or two further hetero-atoms selected from N and/or S and/or O, or



saturated or unsaturated bi- or tricyclic, anellated or bridged heterocycles with 8 to 16 ring atoms, that aside from the essential nitrogen atom, can optionally still contain one or two further hetero-atoms that are selected from N and/or S and/or O;

G<sup>3</sup> has the meaning  $\text{—SO}_2\text{—(CH}_2\text{)}_r\text{—R}^{13}$  (G3)

wherein r and R<sup>13</sup> have the above definition,

G<sup>4</sup> has the meaning



whereby

Ar<sup>1</sup> and Ar<sup>2</sup> can be selected independently from each other from phenyl, pyridyl or naphthyl;

G<sup>5</sup> has the meaning



R<sup>16</sup> is selected from trifluoromethyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, C<sub>3</sub>-C<sub>6</sub>-alkenyloxy, and benzyloxy,

whereby aromatic ring systems in the substituents are R<sup>1</sup>, R<sup>2</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>13</sup>, R<sup>14</sup>, R<sup>15</sup>, R<sup>16</sup>, Ar<sup>1</sup> and Ar<sup>2</sup> and/or in the ring system  $\text{—NR}^{13}\text{R}^{15}$  can be substituted independently from each other by one to three of the same or different groups selected from

halogen, cyano, C<sub>1</sub>-C<sub>6</sub>-alkyl, trifluoromethyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl, phenyl, benzyl, hydroxy, C<sub>1</sub>-C<sub>6</sub>-hydroxyalkyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, C<sub>1</sub>-C<sub>6</sub>-alkoxy entirely or partially substituted by fluorine, benzyloxy, phenoxy, mercapto, C<sub>1</sub>-C<sub>6</sub>-alkylthio,

carboxy, C<sub>2</sub>-C<sub>7</sub>-carboxyalkyl, C<sub>2</sub>-C<sub>7</sub>-carboxyalkenyl, C<sub>2</sub>-C<sub>7</sub>-alkoxycarbonyl, benzyloxycarbonyl, nitro, amino, mono-C<sub>1</sub>-C<sub>6</sub>-alkylamino, di-(C<sub>1</sub>-C<sub>6</sub>-alkyl)-amino and, methylene dioxide for two adjacent residues on the aromatic ring, and whereby

alkyl-, alkenyl- and cycloalkyl residues in the groups G<sup>1</sup>, G<sup>2</sup> and G<sup>3</sup> can be substituted by one or two of the same or different groups which are selected from hydroxy, carboxy, C<sub>2</sub>-C<sub>7</sub>-alkoxycarbonyl, benzyloxycarbonyl, amino, mono-C<sub>1</sub>-C<sub>6</sub>-alkylamino and di-(C<sub>1</sub>-C<sub>6</sub>-alkyl-amino);

their cis- and trans-isomers, E- and Z-isomers, especially in case that A is a cyclopropane ring or D contains one or more double bonds, including the enantiomers, diastereomers and other isomers as well as their racemic or non-racemic mixtures and the corresponding endo- and exo-isomers for the case that the ring system E is bicyclic;

their tautomers;

their acid addition salts including their hydrates and solvates.

According to a further preferred embodiment, the invention relates to compounds of the general Formula (I), whereby

R<sup>1</sup> is selected from hydrogen, halogen, cyano, C<sub>1</sub>-C<sub>6</sub>-alkyl, trifluoromethyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl, C<sub>1</sub>-C<sub>6</sub>-hydroxyalkyl, hydroxy, C<sub>1</sub>-C<sub>4</sub>-alkoxy, benzyloxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>5</sub>-alkanoyloxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>2</sub>-C<sub>5</sub>-alkoxycarbonyl, aminocarbonyl, C<sub>2</sub>-C<sub>5</sub>-alkylaminocarbonyl, C<sub>3</sub>-C<sub>9</sub>-dialkylaminocarbonyl, carboxy, phenyl, phenoxy, phenylthio, pyridyloxy, and NR<sup>5</sup>R<sup>6</sup>, whereby

- $R^5$  and  $R^6$  are selected independently from each other from hydrogen and  $C_1-C_6$ -alkyl;
- $R^2$  is selected from hydrogen, halogen, cyano,  $C_1-C_6$ -alkyl, trifluoromethyl, hydroxy,  $C_1-C_4$ -alkoxy;
- $R^3$  is selected from hydrogen, halogen and  $C_1-C_6$ -alkyl;
- $R^4$  is selected from hydrogen,  $C_1-C_6$ -alkyl,  $C_3-C_6$ -alkenyl,  $C_3-C_6$ -cycloalkyl, hydroxy,  $C_1-C_6$ -alkoxy and benzyloxy;
- $k$  has the meaning 0 or 1,
- $A$  is selected from  $C_1-C_6$ -alkylene, optionally substituted one to three-fold by  $C_1-C_3$ -alkyl, hydroxy, fluorine, or phenyl,
- $C_2-C_6$ -alkylene, in which a methylene unit is isosterically replaced by O, S,  $NR^9$ , CO, SO or  $SO_2$ , whereby, with the exception of CO, the isosteric substitution cannot be adjacent to the amide group and, in  $NR^9$ , the residue  $R^9$  is selected from hydrogen,  $C_1-C_6$ -alkyl,  $C_1-C_6$ -acyl or methanesulfonyl;
- 1,2-cyclopropylene,
- $C_2-C_6$ -Alkenylene, optionally substituted once to three-fold by  $C_1-C_3$ -alkyl, hydroxy, fluorine, cyano or phenyl,
- $C_4-C_6$ -alkadienylene, optionally substituted once to two-fold by  $C_1-C_3$ -alkyl, fluorine, cyano or phenyl;

1,3,5-hexatrienylene, optionally substituted by  $C_1$ - $C_3$ -alkyl, fluorine, cyano, and

ethynylene

D is selected from  $C_2$ - $C_{10}$ -alkylene, optionally substituted once or twice by  $C_1$ - $C_3$ -alkyl or hydroxy;


$C_4$ - $C_{10}$ -alkenylene, optionally substituted once or twice by  $C_1$ - $C_3$ -alkyl or hydroxy;

$C_4$ - $C_{10}$ -alkynylene, optionally substituted once or twice by  $C_1$ - $C_3$ -alkyl or hydroxy; as well as

$C_2$ - $C_{10}$ -alkylene,  $C_4$ - $C_{10}$ -alkenylene or  $C_4$ - $C_{10}$ -alkynylene, in which one to three methylene units is each isosterically replaced by O, S,  $NR^{10}$ , CO, SO or  $SO_2$ , whereby

$R^{10}$  has the same meaning as  $R^9$ , but is selected independently thereof;

E signifies



whereby

q has the meaning 1, 2 or 3;

- R<sup>11</sup> is selected from  
hydrogen, C<sub>1</sub>-C<sub>3</sub>-alkyl, hydroxy, hydroxymethyl, carboxy,  
or C<sub>2</sub>-C<sub>7</sub>-alkoxycarbonyl and
- R<sup>12</sup> is selected from  
hydrogen or an oxo group adjacent to a nitrogen atom or
- R<sup>11</sup> and R<sup>12</sup>, optionally together, form a C<sub>1</sub>-C<sub>3</sub>-alkylene  
bridge under formation of a bicyclic ring system;
- G is selected from G<sub>1</sub>, G<sub>2</sub>, G<sub>3</sub>, G<sub>4</sub> or G<sub>5</sub>, whereby
- G<sub>1</sub> represents  $\text{---}(\text{CH}_2)_r\text{---}(\text{CR}^{14}\text{R}^{15})_s\text{---R}^{13}$  (G<sub>1</sub>)
- r has the meaning 0 to 2,  
s is 0 or 1 and
- R<sup>13</sup> is selected from  
hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-alkenyl, C<sub>3</sub>-C<sub>6</sub>-alkinyl, C<sub>3</sub>-  
C<sub>8</sub>-cycloalkyl; benzyl, phenyl;  
  
monocyclic aromatic five or six-membered heterocycles,  
which can contain one to three hetero-atoms selected  
from the group N and/or S and/or O and are either bound  
directly or over a methylene group,  
  
anellated bi- and tricyclic aromatic or partially  
hydrated carbocyclic ring systems with 8 to 16 ring  
atoms and at least one aromatic ring, wherein the  
linkage can occur either over an aromatic or a hydrated  
ring and either directly or over a methylene group,  
  
anellated bi- and tricyclic aromatic or partially  
hydrated heterocyclic ring systems with 8 to 16 ring

atoms and at least one aromatic ring, wherein one to three ring atoms can be selected from N and/or S and/or O and the linkage can occur either over an aromatic ring or a hydrated ring and either directly or over a methylene group,

R<sup>14</sup> has the same meaning as R<sup>13</sup>, but is selected independently thereof;


R<sup>15</sup> is selected from  
hydrogen, hydroxy, methyl, benzyl, phenyl;

monocyclic aromatic five or six-membered heterocycles, which can contain one to three hetero-atoms selected from the group N and/or S and/or O and are either bound directly or over a methylene group,

anellated bi- and tricyclic aromatic or partially hydrated carbocyclic ring systems with 8 to 16 ring atoms and at least one aromatic ring, wherein the linkage can occur either over an aromatic or a hydrated ring and either directly or over a methylene group,


anellated bi- and tricyclic aromatic or partially hydrated heterocyclic ring systems with 8 to 16 ring atoms and at least one aromatic ring, wherein one to three ring atoms can be selected from N and/or S and/or O and the linkage can occur either over an aromatic ring or a hydrated ring and either directly or over a methylene group;

G<sup>2</sup> is selected from



(G2a)

or



(G2b)

whereby  $r$  and  $s$  as well as the substituents  $\text{R}^{13}$  to  $\text{R}^{15}$  can have the above meaning, or the grouping



can also be a nitrogen heterocycle bound over the nitrogen atom selected from

saturated or unsaturated monocyclic, four to eight-membered heterocycles, which, aside from the essential nitrogen atom, can optionally still contain one or two further hetero-atoms selected from N and/or S and/or O, or

saturated or unsaturated bi- or tricyclic, anellated or bridged heterocycles with 8 to 16 ring atoms, that aside from the essential nitrogen atom, can optionally still contain one or two further hetero-atoms that are selected from N and/or S and/or O;

$\text{G}^3$  has the meaning  $\text{---SO}_2\text{---}(\text{CH}_2)_r\text{---R}^{13}$  (G3)

wherein  $r$  and  $\text{R}^{13}$  have the above definition,

$\text{G}^4$  has the meaning



(G4)

whereby

$\text{Ar}^1$  and  $\text{Ar}^2$  can be selected independently from each other from phenyl, pyridyl or naphthyl;

G<sup>5</sup> has the meaning

—COR<sup>16</sup>

(G<sup>5</sup>)

R<sup>16</sup> is selected from  
trifluoromethyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, C<sub>3</sub>-C<sub>6</sub>-alkenyloxy, and  
benzyloxy,

whereby aromatic ring systems in the substituents are R<sup>1</sup>,  
R<sup>2</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>13</sup>, R<sup>14</sup>, R<sup>15</sup>, R<sup>16</sup>, Ar<sup>1</sup> and Ar<sup>2</sup> and/or in the  
ring system —NR<sup>13</sup>R<sup>15</sup> can be substituted independently from  
each other by one to three of the same or different groups  
selected from

halogen, cyano, C<sub>1</sub>-C<sub>6</sub>-alkyl, trifluoromethyl, C<sub>3</sub>-C<sub>8</sub>-  
cycloalkyl, phenyl, benzyl, hydroxy, C<sub>1</sub>-C<sub>6</sub>-hydroxyalkyl, C<sub>1</sub>-  
C<sub>6</sub>-alkoxy, C<sub>1</sub>-C<sub>6</sub>-alkoxy entirely or partially substituted by  
fluorine, benzyloxy, phenoxy, mercapto, C<sub>1</sub>-C<sub>6</sub>-alkylthio,  
carboxy, C<sub>2</sub>-C<sub>7</sub>-carboxyalkyl, C<sub>2</sub>-C<sub>7</sub>-carboxyalkenyl, C<sub>2</sub>-C<sub>7</sub>-  
alkoxycarbonyl, benzyloxycarbonyl, nitro, amino, mono-C<sub>1</sub>-C<sub>6</sub>-  
alkylamino, di-(C<sub>1</sub>-C<sub>6</sub>-alkyl)-amino and, methylene dioxide in  
the case of two adjacent residues on the aromatic ring,

whereby alkyl-, alkenyl- and cycloalkyl residues in the  
groups G<sup>1</sup>, G<sup>2</sup> and G<sup>3</sup> can be substituted by one or two of the  
same or different groups which are selected from  
hydroxy, carboxy, C<sub>2</sub>-C<sub>7</sub>-alkoxycarbonyl, benzyloxycarbonyl,  
amino, mono-C<sub>1</sub>-C<sub>6</sub>-alkylamino and di-(C<sub>1</sub>-C<sub>6</sub>-alkyl-amino).

According to a further embodiment, the invention especially  
relates to compounds of the general Formula (1), wherein

R<sup>1</sup> is selected from  
hydrogen, halogen, cyano, methyl, ethyl,  
trifluoromethyl, hydroxy, C<sub>1</sub>-C<sub>4</sub>-alkoxy, benzyloxy, C<sub>1</sub>-  
C<sub>5</sub>-alkanoyloxy, methylthio, ethylthio, methoxycarbonyl,



tert-butoxycarbonyl aminocarbonyl, carboxy, phenoxy, and phenylthio


- R<sup>2</sup> is selected from  
hydrogen, halogen, trifluoromethyl, hydroxy;
- R<sup>3</sup> is selected from  
hydrogen, halogen;
- R<sup>4</sup> is selected from  
hydrogen, C<sub>1</sub>-C<sub>3</sub>-alkyl, allyl, hydroxy and C<sub>1</sub>-C<sub>3</sub>-alkoxy;
- k is 0 or 1,
- A is selected from  
C<sub>1</sub>-C<sub>6</sub>-alkylene, optionally substituted once or twice by  
C<sub>1</sub>-C<sub>3</sub>-alkyl, hydroxy or fluorine;
- C<sub>2</sub>-C<sub>6</sub>-alkylene, in which a methylene unit is  
isosterically replaced by O, S, NR<sup>9</sup>, CO, SO or SO<sub>2</sub>,  
whereby, with the exception of CO, the isosteric  
substitution cannot be adjacent to the amide group,
- C<sub>2</sub>-C<sub>6</sub>-alkylene, optionally substituted once or twice by  
C<sub>1</sub>-C<sub>3</sub> alkyl, hydroxy and/or fluorine;
- C<sub>4</sub>-C<sub>6</sub>-alkadienylene, optionally substituted by C<sub>1</sub>-C<sub>3</sub>-  
alkyl or one or two fluorine atoms;
- 1,3,5-hexatrienylene, optionally substituted by  
fluorine;
- D is selected from  
C<sub>2</sub>-C<sub>8</sub>-alkylene, optionally substituted once or twice by  
methyl or hydroxy;

C<sub>4</sub>-C<sub>8</sub>-alkenylene, optionally substituted once or twice by methyl or hydroxy;

C<sub>4</sub>-C<sub>8</sub>-alkinylene, optionally substituted once or twice by methyl or hydroxy; and

C<sub>2</sub>-C<sub>8</sub>-alkylene, C<sub>4</sub>-C<sub>8</sub>-alkenylene or C<sub>4</sub>-C<sub>8</sub>-alkinylene, wherein one to three methylene units are each isosterically replaced by O, S, NH, N(CH<sub>3</sub>) N(COCH<sub>3</sub>), N(SO<sub>2</sub>CH<sub>3</sub>), CO, SO or SO<sub>2</sub>, whereby

E has the meaning



whereby

q is 1 or 2;

R<sub>11</sub> is selected from hydrogen, C<sub>1</sub>-C<sub>3</sub>-alkyl, hydroxymethyl, or carboxy, and

R<sub>12</sub> is selected from hydrogen or an oxo group adjacent to a nitrogen atom

G is selected from G<sub>1</sub>, G<sub>2</sub>, G<sub>3</sub>, G<sub>4</sub> or G<sub>5</sub>, whereby

G<sub>1</sub>  $\text{---} (\text{CH}_2)_r \text{---} (\text{CR}^{14}\text{R}^{15})_s \text{---} \text{R}^{13}$  (G<sub>1</sub>)  
represents

r is 0 to 2 and,

$R^{13}$  is selected from

benzocyclobutyl, indanyl, indenyl, oxoindanyl, naphthyl, dihydronaphthyl, tetrahydronaphthyl, oxotetrahydronaphthyl, biphenylenyl, fluorenyl, oxofluorenyl, anthryl, dihydroanthryl, oxodihydroanthryl, dioxodihydroanthryl, phenanthryl, dihydrophenanthryl, oxodihydrophenanthryl, dibenzocycloheptenyl, oxodibenzocycloheptenyl, dihydrodibenzocycloheptenyl, oxodihydrodibenzocycloheptenyl, dihydrodibenzocyclooctenyl, tetrahydrodibenzocyclooctenyl or oxotetrahydrodibenzocyclooctenyl bound directly or over a methylene group;

furyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, iso-thiazolyl, pyrazolyl, imidazolyl, oxadiazolyl, thiadiazolyl, triazolyl, pyridyl, pyrazinyl, pyridazinyl, pyrimidinyl, triazinyl, imidazothiazolyl, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl, indolinyl, isoindolinyl, oxoindolinyl, dioxoindolinyl, benzoxazolyl, oxobenzooxazolyl, benzoisoxazolyl, oxobenzoisoxazolyl, benzothiazolyl, oxobenz-thiazolyl, benzoisothiazolyl, oxobenzoisothiazolyl, benzimidazolyl, oxobenzimidazolyl, indazolyl, oxoindazolyl, benzofurazanyl, benzothiadiazolyl, benzotriazolyl, oxazolopyridyl, oxodihydrooxazolo-pyridyl, thiazolopyridyl, oxodihydrothiazolopyridyl, isothiazolopyridyl, imidazopyridyl, oxodihydroimidazo-pyridyl, pyrazolopyridyl, oxodihydropyrazolopyridyl, thienopyrimidinyl, chromanyl, chromanonyl, benzopyranlyl, chromonyl, quinoloyl, isoquinoloyl, dihydroquinolyl, oxodihydroquinolyl, tetrahydroquinolyl, oxotetrahydroquinolyl, benzodioxanyl, quinoxalyl,

quinazolinyl, naphthyridinyl, carbazolyl, tetrahydrocarbazolyl, oxotetrahydrocarbazolyl, pyridoindolyl, acridinyl, oxodihydroacridinyl, phenanthridinyl, dihydrophenanthridinyl, oxodihydrophenanthridinyl, dibenzoisoquinolinyl, dihydrodibenzoisoquinolinyl, oxodihydrodibenzoisoquinolinyl, phenothiazinyl, dihydrodibenzooxepinyl, oxodihydrodibenzooxepinyl, benzocycloheptathienyl, oxobenzocycloheptathienyl, dihydrothienobenzothiepinyl, oxodihydrothienobenzothiepinyl, dihydrodibenzothiepinyl, oxodihydrodibenzothiepinyl, octahydrodibenzothiepinyl, dibenzoazepinyl, dihydrodibenzoazepinyl, oxodihydrodibenzoazepinyl, octahydrodibenzoazepinyl, benzocycloheptapyridyl, oxobenzocycloheptapyridyl, pyridobenzoazepinyl, dihydropyridobenzoazepinyl, oxodihydropyridobenzoazepinyl, dihydropyridobenzo-diazepinyl, dihydrodibenzooxazepinyl, dihydropyridobenzooxepinyl, dihydropyridobenzoazepinyl, oxodihydropyridobenzoazepinyl, dihydrodibenzothiazepinyl, oxodihydrodibenzothiazepinyl, dihydropyridobenzothiazepinyl or oxodihydropyridobenzothiazepinyl bound directly or over a methylene group;


R<sup>14</sup> is synonymous with R<sup>13</sup> but is selected independent thereof;

R<sup>15</sup> is selected from


hydroxy, methyl, benzyl, phenyl, indanyl, indenyl, naphthyl, dihydronaphthyl, tetrahydronaphthyl, furyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyrazolyl, imidazolyl, oxadiazolyl, thiadiazolyl, triazolyl, pyridyl, pyrazinyl, pyridazinyl, pyrimidinyl, triazinyl, benzofuryl, benzothienyl, indolyl, indolinyl, benzoazazolyl, benzothiazolyl, benzoimidazolyl, chromanyl, quinolyl or

tetrahydroquinolyl bound directly or over a methylene group;

G<sup>2</sup> is selected from



or




whereby r and s and the substituents R<sup>13</sup> to R<sup>15</sup> can have the above meaning, or the grouping




represents the ring of azetidine bound over the nitrogen or one of the following residues: pyrrolidine, piperidine, (1H)-tetrahydropyridine, hexahydroazepine, (1H)-tetrahydroazepine, octahydroazocine, pyrazolidine, piperazine, hexahydrodiazepine, morpholine, hexahydro-oxazepine, thiomorpholine, thiomorpholin-1,1-dioxide, of 5-aza-bicyclo-[2.1.1]hexane, 2-aza-bicyclo[2.2.1]heptane, 7-aza-bicyclo-[2.2.1]heptane, 2,5-diaza-bicyclo[2.2.1]heptane, 2-aza-bicyclo[2.2.2]octane, 8-aza-bicyclo[3.2.1]octane, 2,5-diaza-bicyclo[2.2.2]octane, 9-aza-bicyclo[3.3.1]nonane, indoline, isoindoline, (1H)-dihydroquinoline, (1H)-tetrahydroquinolin, (2H)-tetrahydroisoquinoline, (1H)-tetrahydroquinoxaline, (4H)-dihydrobenzooxazine, (4H)-dihydrobenzothiazine, (1H)-tetrahydrobenzo[b]azepine, (1H)-tetrahydrobenzo[c]azepine, (1H)-tetrahydrobenzo[d]azepine, (5H)-tetrahydrobenzo[b]ox-azepine, (5H)-tetrahydrobenzo[b]thiazepine, 1,2,3,4-tetrahydro-9H-pyrido[3,4-b]indole, (10H)-dihydroacridine, (10H)-dihydrophenanthridine, 1,2,3,4-tetrahydroacridanone,

(10H)-phenoxazine, (10H)-phenothiazine, (5H)-dibenzoazepine, (5H)-dihydrodibenzoazepine, (5H)-octahydrodibenzoazepine, dihydrobenzo[d,e]isoquinoline, (5H)-dihydrodibenzodiazepine, (5H)-benzo[b]pyrido-[f]azepine, (5H)-Dihydrobenzo[b]pyrido-[f]azepine, (11H)-Dihydrodibenzo[b,e]oxazepine, (11H)-dihydrodibenzo[b,e]thiazepine, (10H)-dihydrodibenzo[b,f]-oxazepine, (10H)-dihydrodibenzo[b,f]thiazepine, (5H)-tetra-hydrodibenzoazocine, (11H)-dihydrobenzo[e]pyrido[b]-1,4-diazepin-6-one or (11H)-Dihydrobenzo[b]pyrido[e]-1,4-diazepin-5-one.



wherein  $r$  and  $R^{13}$  have the above definition,

$G^4$  has the meaning



whereby

$\text{Ar}^1$  and  $\text{Ar}^2$  are selected independently from each other from phenyl, pyridyl or naphthyl;

$G^5$  has the meaning



$R^{16}$  is selected from trifluoromethyl,  $\text{C}_1\text{--C}_6\text{-alkoxy}$ ,  $\text{C}_3\text{--C}_6\text{-alkenyloxy}$ , and benzyloxy,

whereby aromatic ring systems in the substituents can be substituted independently from each other by one to three of the same or different groups selected from

halogen, cyano, C<sub>1</sub>-C<sub>6</sub>-alkyl, trifluoromethyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl, phenyl, benzyl, hydroxy, C<sub>1</sub>-C<sub>6</sub>-hydroxyalkyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, C<sub>1</sub>-C<sub>6</sub>-alkoxy entirely or partially substituted by fluorine, benzyloxy, phenoxy, mercapto, C<sub>1</sub>-C<sub>6</sub>-alkylthio, carboxy, C<sub>2</sub>-C<sub>7</sub>-carboxyalkyl, C<sub>2</sub>-C<sub>7</sub>-carboxyalkenyl, C<sub>2</sub>-C<sub>7</sub>-alkoxycarbonyl, benzyloxycarbonyl, nitro, amino, mono-C<sub>1</sub>-C<sub>6</sub>-alkylamino, di-(C<sub>1</sub>-C<sub>6</sub>-alkyl)-amino and, methylene dioxide in the case of two adjacent residues on the aromatic ring, and

whereby alkyl-, alkenyl- and cycloalkyl residues in the groups G<sup>1</sup>, G<sup>2</sup> and G<sup>3</sup> can be substituted by one or two of the same or different groups which are selected from hydroxy, carboxy, C<sub>2</sub>-C<sub>7</sub>-alkoxycarbonyl, benzyloxycarbonyl, amino, mono-C<sub>1</sub>-C<sub>6</sub>-alkylamino and di-(C<sub>1</sub>-C<sub>6</sub>-alkyl-amino).

According to a further very preferred embodiment, the invention relates to compounds of Formula (I), wherein

R<sup>1</sup> is selected from hydrogen, fluorine, chlorine, bromine, methyl, ethyl, trifluoromethyl, hydroxy, C<sub>1</sub>-C<sub>4</sub>-alkoxy, methylthio, ethylthio, carboxy and phenoxy;

R<sup>2</sup> is selected from hydrogen, chlorine and methyl;

R<sup>3</sup> is hydrogen;

R<sup>4</sup> is selected from hydrogen, C<sub>1</sub>-C<sub>3</sub>-alkyl and hydroxy,

k is 0

A is selected from C<sub>2</sub>-C<sub>6</sub>-alkylene, which is optionally substituted once or twice by hydroxy or fluorine;

C<sub>2</sub>-C<sub>6</sub>-alkylene, wherein a methylene unit is isosterically replaced by O, S or CO, whereby, with the

exception of CO, the isosteric substitution cannot be adjacent to the amide group;

C<sub>2</sub>-C<sub>8</sub>-alkenylene which is optionally substituted by C<sub>1</sub>-C<sub>3</sub>-alkyl and/or fluorine;

C<sub>4</sub>-C<sub>6</sub>-alkadienylene;

- D is selected from  
C<sub>2</sub>-C<sub>8</sub>-alkylene which is optionally substituted by methyl or hydroxy;


C<sub>4</sub>-C<sub>8</sub>-alkenylene, which is optionally substituted by hydroxy;

C<sub>4</sub>-C<sub>8</sub>-alkinylene, which is optionally substituted by hydroxy;

C<sub>2</sub>-C<sub>8</sub>-alkylene, C<sub>4</sub>-C<sub>8</sub>-alkenylene, C<sub>4</sub>-C<sub>8</sub>-alkinylene wherein a methylene unit is respectively isosterically replaced by O, NH, N(CH<sub>3</sub>), CO or SO<sub>2</sub> or an ethylene group is isosterically replaced by a group NH-CO and/or CO-NH or a propylene group is isosterically replaced by a group NH-CO-O and/or O-CO-NH;


- E is selected from piperazine or hexahydro-1,4-diazepine (homopiperazine), wherein the ring can be optionally substituted by one or two methylene groups and/or by an oxo group adjacent to a nitrogen atom;

- G is selected from hydrogen, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl, methoxy-carbonyl, tert-butoxycarbonyl, benzyloxycarbonyl, trifluoroacetyl, diphenylphosphinoyl, or a group







and



and



and



wherein

r has the meaning 0 or 1

s is 0 or 1,

R<sup>13</sup> is selected from hydrogen, methyl, benzyl, phenyl,

indanyl, indenyl, oxoindanyl, naphthyl, tetrahydronaphthyl, fluorenyl, oxofluorenyl, anthryl, dihydroanthryl, oxodihydroanthryl, dioxodihydroanthryl, phenanthryl, dihydrophenanthryl, oxydihydrophenanthryl, dibenzocycloheptenyl, dihydrodibenzocycloheptenyl, oxodihydrodibenzocycloheptyl bound directly or over a methylene group,

furyl, thienyl, oxazolyl, isooxazolyl, thiazolyl, imidazolyl, oxadiazolyl, pyridyl, pyrazinyl, pyrimidinyl, imidazothiazolyl, benzofuryl, benzothienyl, indolyl, indolinyl, oxoindolinyl, dioxoindolinyl, benzoxazolyl, oxobenzoxazolyl, benzothiazolyl, oxobenzthiazolyl, benzimidazolyl, oxobenzimidazolyl, indazolyl, benzofurazanyl, benzotriazolyl, oxazolo-pyridyl, oxodihydrooxazolopyridyl, thiazolopyridyl,

oxodihydrothiazolopyridyl, imidazopyridyl, oxodihydroimidazopyridyl, chromanyl, chromanonyl, benzopyranyl, chromonyl, quinolyl, isoquinolyl, oxodihydroquinolinyl, tetrahydroquinolyl, oxotetrahydroquinolinyl, benzodioxanyl, quinazolinyl, carbazolyl, acridinyl, dihydroacridinyl, oxodihydroacridinyl, dihydrophenanthridinyl dihydrobenzoisoquinolinyl, phenothiazinyl, dihydrodibenzoxepinyl, benzocycloheptathienyl, dihydrothienobenzothiepinyl, dihydrodibenzothiepinyl, oxodihydrodibenzothiepinyl, dihydrodibenzoazepinyl, oxodihydrodibenzoazepinyl, octahydrodibenzoazepinyl, benzocycloheptapyridyl, dihydropyridobenzodiazepinyl, dihydrodibenzothiazepinyl bound directly or over a methylene group,

R<sup>14</sup> is selected from hydrogen, methyl, benzyl, phenyl;

R<sup>15</sup> is selected from hydrogen, hydroxy, methyl, benzyl, phenyl;

naphthyl, tetrahydronaphthyl, furyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridyl, benzofuryl, benzothienyl, indolyl, indolinyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, chromanyl, quinolyl or tetrahydroquinolyl bound directly or over a methylene group;

whereby the grouping -NR<sup>13</sup>R<sup>15</sup> represents a ring bound over the nitrogen atom of a residue from the series

pyrrolidine, piperidine, hexahydroazepine, piperazine, hexahydrodiazepine, morpholine, hexahydroazepine, thiomorpholine, 7-aza-bicyclo[2.2.1]heptane, 2,5-diazabicyclo[2.2.1]heptane, indoline, isoindoline, (1H)-dihydroquinoline, (1H)-tetrahydroquinoline, (2H)-tetrahydroisoquinoline, (4H)-dihydrobenzoxazine, (4H)-dihydrobenzothiazine, (1H)-tetrahydrobenzo[b]azepine, (1H)-tetrahydrobenzo[d]azepine, (5H)-tetrahydroben-

zo[b]oxazepine, (5H)-tetrahydrobenzo[b]thiazepine, (10H)-dihydroacridine, 1,2,3,4-tetrahydroacridanone, (10H)-dihydrophenanthridine, (1H)-dihydrobenzo-[d,e]isoquinoline, (10H)-phenothiazine, (5H)-dibenzo[b,f]azepine, (5H)-dihydrodibenzo[b,f]azepine, (5H)-dihydrodibenzo[c,e]azepine, (5H)-dihydrodibenzo-diazepine, (11H)-dihydrodibenzo[b,e]oxazepine (11H)-dihydrodibenzo[b,e]thiazepine, (5H)-dihydrobenzo[b]pyrido[3,2-f]azepine and (11H)-6-oxodihydrobenzo[e]pyrido[3,2-b][1,4]diazepine, and whereby

aromatic ring systems in the substituents can be substituted, independently of each other, by one to three of the same or different groups selected from halogen, cyano, C<sub>1</sub>-C<sub>6</sub>-alkyl, trifluoromethyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl, phenyl, benzyl, hydroxy, C<sub>1</sub>-C<sub>6</sub>-hydroxyalkyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, C<sub>1</sub>-C<sub>6</sub>-alkoxy which can be entirely or partially substituted by fluorine, benzyloxy, phenoxy, mercapto, C<sub>1</sub>-C<sub>6</sub>-alkylthio, carboxy, C<sub>2</sub>-C<sub>7</sub>-carboxyalkyl, C<sub>2</sub>-C<sub>7</sub>-carbocycloalkenyl, C<sub>2</sub>-C<sub>7</sub>-alkoxycarbonyl, benzyloxycarbonyl, nitro, amino, mono-C<sub>1</sub>-C<sub>6</sub>-alkylamino or di-(C<sub>1</sub>-C<sub>6</sub>-alkyl)-amino, and in the case of two adjacent residues on the aromatic ring, methylenedioxy, and

whereby alkyl, alkenyl and cycloalkyl residues in the group G can be substituted by one or two of the same or different groups which are selected from hydroxy, carboxy, C<sub>2</sub>-C<sub>7</sub>-alkoxycarbonyl, benzyloxycarbonyl, amino, mono-C<sub>1</sub>-C<sub>6</sub>-alkylamino and di-(C<sub>1</sub>-C<sub>6</sub>-alkyl) amino.

Furthermore, according to a particularly preferred embodiment, the invention relates to compounds of formula (I), wherein

R<sup>1</sup> is selected from hydrogen, fluorine, methyl, trifluoromethyl ethylthio;

R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are each hydrogen;

k has the meaning 0,

A is selected from

ethylene, propylene or butylene which are each optionally substituted by hydroxy or one or two fluorine atoms; or OCH<sub>2</sub>,SCH<sub>2</sub>;

ethenylene, or 1,3-butadienylene;

D is selected from

C<sub>2</sub>-C<sub>6</sub>-alkylene which is optionally substituted by hydroxy;

C<sub>4</sub>-C<sub>6</sub> alkenylene;

C<sub>4</sub>-C<sub>6</sub> alkynylene; or

C<sub>2</sub>-C<sub>6</sub> alkylene, C<sub>4</sub>-C<sub>6</sub> alkenylene or C<sub>4</sub>-C<sub>6</sub> alkynylene, wherein one or two methylene units is isosterically replaced by O, NH, CO or SO<sub>2</sub>;

E is selected from piperazine or hexahydro-1,4-diazezepine;

G is selected from

phenyl, benzyl, phenethyl, diphenylmethyl, naphthyl, tetrahydronaphtyl, naphthylmethyl, fluorenyl fluorenylmethyl, anthrylmethyl, dihydrodibenzo-cycloheptenyl;

furylmethyl, thienylmethyl, thiazolylmethyl, pyridylmethyl, benzothienylmethyl, quinolylmethyl, phenylthienylmethyl, phenylpyridylmethyl, benzocycloheptapyridinyl, dihydrobenzocycloheptapyridinyl, dihydrodibenzooxepinyl, dihydrodibenzothiepinyl, dihydrodibenzoazepinyl, dihydrobenzopyridodiazepinyl;

formyl, acetyl, pivaloyl, phenylacetyl, diphenylacetyl, diphenylpropionyl, naphthylacetyl, benzoyl, naphthoyl, oxofluorenylcarbonyl, oxodihydroanthrylcarbonyl, dioxodihydroanthrylcarbonyl,

furoyl, pyridylacetyl, pyridylcarbonyl, chromonylcarbonyl, quinolylcarbonyl,

phenylaminocarbonyl, naphthylaminocarbonyl, tetrahydronaphthylaminocarbonyl, dibenzylaminocarbonyl, benzylphenylaminocarbonyl, diphenylaminocarbonyl, indolinyl-N-carbonyl, isoindolin-N-carbonyl, tetrahydroquinolinyl-N-carbonyl, carbazolyl-N-carbonyl, tetrahydrobenzoazepinyl-N-carbonyl, dihydrodibenzoazepin-N-carbonyl, dihydrobenzopyridoazepinyl-N-carbonyl, oxodihydrobenzopyridoazepinyl-N-carbonyl;

methanesulfonyl, toluenesulfonyl, naphthylsulfonyl, quinolinsulfonyl and

diphenylphosphinoyl,

wherein aromatic ring systems can be substituted independently of each other by one to three of the same or different groups which are selected from halogen, cyano, C<sub>1</sub>-C<sub>6</sub>-alkyl, trifluoromethyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl, phenyl, benzyl, hydroxy, C<sub>1</sub>-C<sub>6</sub>-hydroxyalkyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, C<sub>1</sub>-C<sub>6</sub>-alkoxy entirely or partially substituted by fluorine, benzyloxy,

phenoxy, mercapto, C<sub>1</sub>-C<sub>6</sub>-alkylthio, carboxy, C<sub>2</sub>-C<sub>7</sub>-carboxyalkyl, C<sub>2</sub>-C<sub>7</sub>-carboxyalkenyl, C<sub>2</sub>-C<sub>7</sub>-alkoxycarbonyl, benzyloxycarbonyl, nitro, amino, mono-C<sub>1</sub>-C<sub>6</sub>-alkylamino or di-(C<sub>1</sub>-C<sub>6</sub>-alkyl)-amino, and in the case of two adjacent residues in the aromatic ring methylenedioxy, and whereby

alkyl, alkenyl and cycloalkyl residues in the group G can be substituted by one or two of the same or different groups which are selected from

hydroxy, carboxy, C<sub>2</sub>-C<sub>7</sub>-alkoxycarbonyl, benzyloxycarbonyl, amino, mono-C<sub>1</sub>-C<sub>6</sub>-alkylamino or di-(C<sub>1</sub>-C<sub>6</sub>-alkyl)-amino.

The following end products represent very particular concrete preferred embodiments of the invention:

a) N-[4-(4-diphenylmethylpiperazin-1-yl)-3-hydroxybutyl]-3-pyridin-3-yl-acrylamide;  
 N-[3-(4-diphenylmethylpiperazin-1-yl)-propoxy]-3-pyridin-3-yl-acrylamide;  
 N-[4-(4-diphenylmethylpiperazin-1-yl)-4-oxo-butyl]-3-pyridin-3-yl-acrylamide;  
 N-[3-(4-diphenylmethylpiperazin-1-sulfonyl)-propyl]-3-pyridin-3-yl-acrylamide;  
 N-[2-[2-(4-diphenylmethylpiperazin-1-yl)-ethoxy]-ethyl]-3-pyridin-3-yl-acrylamide;  
 N-(4-{4-[bis-(4-fluorophenyl)-methyl]-piperazin-1-yl}-but-2-in-yl)-3-pyridin-3-yl-acrylamide;  
 N-{4-[4-(4-carboxyphenyl-phenylmethyl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-acrylamide and  
 N-(4-{4-[(4-aminophenyl)-phenylmethyl]-piperazin-1-yl}-butyl)-3-pyridin-3-yl-acrylamide.

b) N-{4-[4-(9H-fluoren-9-yl)-piperazin-1-yl]-butyl}-2-(pyridin-3-yloxy)-acetamide;  
 N-{5-[4-(9H-fluoren-9-yl)-piperazin-1-yl]-pentyl}-3-pyridin-3-yl-acrylamide;

N-{6-[4-(9H-fluoren-9-yl)-piperazin-1-yl]-hexyl}-3-pyridin-3-yl-acrylamide;  
 3-pyridin-3-yl-N-{4-[4-(1,2,3,4-tetrahydronaphthalin-1-yl)-piperazin-1-yl]-butyl}-acrylamide;  
 3-pyridin-3-yl-N-{4-[4-(5,6,7,8-tetrahydronaphthalin-1-yl)-piperazin-1-yl]-butyl}-acrylamide and  
 N-{4-[4-(naphthalin-1-yl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-acrylamide.

c) N-[4-(4-biphenyl-2-yl-piperazin-1-yl)-butyl]-3-pyridin-3-yl-propionamide;  
 N-[5-(4-biphenyl-2-yl-piperazin-1-yl)-pentyl]-3-pyridin-3-yl-acrylamide;  
 N-[6-(4-biphenyl-2-yl-piperazin-1-yl)-hexyl]-3-pyridin-3-yl-acrylamide;  
 N-[4-(4-biphenyl-2-yl-piperazin-1-yl)-butyl]-2-(pyridin-3-yloxy)-acetamide as well as  
 N-[4-(4-biphenyl-2-yl-piperazin-1-yl)-butyl]-5-(pyridin-3-yl)-penta-2,4-diensäureamide.

d) N-{4-[4-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-yl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-propionamide;  
 N-{5-[4-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-yl)-piperazin-1-yl]-pentyl}-3-pyridin-3-yl-acrylamide;  
 N-{6-[4-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-yl)-piperazin-1-yl]-hexyl}-3-pyridin-3-yl-acrylamide;  
 N-{4-[4-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-yl)-piperazin-1-yl]-butyl}-5-(pyridin-3-yl)-penta-2,4-diensäureamide;  
 N-{4-[4-(6,11-dihydro-dibenzo[b,e]oxepin-11-yl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-propionamide and  
 N-{2-[4-(6,11-dihydrodibenzo[b,e]thiepin-11-yl)-piperazin-1-yl]-ethyl}-3-pyridin-3-yl-acrylamide.

e) N-[4-(4-diphenylacetyl-piperazin-1-yl)-butyl]-3-pyridin-3-yl-acrylamide;  
 N-[4-(4-benzoylpiperazin-1-yl)-butyl]-3-pyridin-3-yl-acrylamide;

N-{4-[4-(2-aminobenzoyl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-acrylamide;

N-{4-[4-(4-carboxybenzoyl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-acrylamide;

N-{4-[4-(biphenyl-2-carbonyl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-acrylamide;

N-{4-[4-(9-oxo-9H-fluoren-4-carbonyl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-acrylamide and

N-{4-[4-(furan-2-carbonyl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-acrylamide.

f) N-{4-[4-(naphthalin-1-yl-aminocarbonyl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-propionamide;

N-{4-[4-(diphenylaminocarbonyl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-acrylamide;

N-{4-[4-(naphthalin-2-sulfonyl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-acrylamide as well as

N-[4-(4-diphenylphosphinonyl-piperazin-1-yl)-butyl]-3-pyridin-3-yl-acrylamide.

g) N-[4-(4-biphenyl-2-yl-piperazin-1-yl)-butyl]-3-pyridin-3-yl-acrylamide;

N-[4-[4-(9H-fluoren-9-yl)-piperazin-1-yl]-butyl]-3-pyridin-3-yl-acrylamide and


N-{4-[4-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-yl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-acrylamide.

In the following, a series of compounds with the respective specific substituent definitions are listed in Table 1 without any limitation for further illustration of the compounds according to the invention.



Table 1

Exemplifying  
compounds of formula  
(I) according to the  
invention















Nr	R <sup>1</sup> - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
1	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N 
2	H	0	CH=CH	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N 
3	H	0	CH=CH	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N  -CH <sub>2</sub> -C≡CH
4	H	0	CH=CH	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N  -CH <sub>2</sub> -CH <sub>2</sub> -COOH
5	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N  -CH <sub>2</sub> -CH <sub>2</sub> -C(=O)-O-CH <sub>2</sub> - 
6	H	0	CH=CH	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N  -CH <sub>2</sub> -CH <sub>2</sub> -N(H)-CH(CH <sub>3</sub> ) <sub>2</sub>
7	H	0	CH=CH	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N  -CH <sub>2</sub> - 
8	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N  -N 

Table 1 (continuation)

Nr	R' - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
9	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N<img alt="piperazine ring" data-bbox="665 135 715 165"/>-N-CH<img alt="morpholine ring" data-bbox="755 135 845 175"/>
10	H	0	CH=CH	H	CH <sub>2</sub> C≡CCH <sub>2</sub> -N<img alt="piperazine ring" data-bbox="665 185 715 215"/>-N-CH<img alt="phenyl ring" data-bbox="765 185 815 215"/>
11	H	0	OCH <sub>2</sub>	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N<img alt="piperazine ring with a methyl group (H3C) on one nitrogen" data-bbox="665 225 715 275"/>-N-CH<img alt="phenyl ring" data-bbox="765 225 815 255"/>
12	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	CH <sub>2</sub> CH <sub>2</sub> OCH <sub>2</sub> CH <sub>2</sub> -N<img alt="piperazine ring" data-bbox="665 275 715 305"/>-N-CH<img alt="phenyl ring" data-bbox="765 275 815 305"/>
13	H	0		H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N<img alt="piperazine ring" data-bbox="665 315 715 345"/>-N-CH<img alt="phenyl ring" data-bbox="765 315 815 345"/>
14	H	0	CH=CH	H	CH <sub>2</sub> CH <sub>2</sub> -N<img alt="piperazine ring" data-bbox="665 365 715 395"/>-N-CH<img alt="2-phenylpropan-1-ol side chain" data-bbox="715 355 835 395"/>
15	H	0	CH=CH	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N<img alt="piperazine ring" data-bbox="665 405 715 435"/>-N-CH<img alt="4-phenylbenzoic acid side chain" data-bbox="765 405 865 435"/>
16	H	0	CH=CH	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N<img alt="piperazine ring" data-bbox="665 445 715 475"/>-N-CH<img alt="1,2-diphenylethyl side chain" data-bbox="745 435 845 505"/>
17	H	0	CH=CH	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N<img alt="piperazine ring" data-bbox="665 515 715 545"/>-N-CH<img alt="1,2-diphenyl-2-carboxyethyl side chain" data-bbox="745 515 865 585"/>
18	H	0	CH=CH	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N<img alt="piperazine ring" data-bbox="665 585 715 615"/>-N-CH<img alt="1,2-diphenylethyl side chain" data-bbox="745 595 845 645"/>

00956001.061600

Table 1 (continuation)

Nr	R <sup>1</sup> - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
19	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	<chem>CCCCCN1CCN(CC1)Cc2c3ccccc3cc2</chem>
20	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	<chem>CCCCCN1CCN(CC1)Cc2ccc3ccccc3c2</chem>
21	H	0	OCH <sub>2</sub>	H	<chem>CCN1CCN(CC1)Cc2ccccn2</chem>
22	H	0	CH=CH	H	<chem>CCCCN1CCN(CC1)Cc2c3ccccc3n2</chem>
23	H	0	CH=CH·CH=CH	H	<chem>CCCCN1CCN(CC1)C2c3ccccc3C2</chem>
24	H	0	OCH <sub>2</sub>	H	<chem>CCN1CCN(CC1)Cc2ccccc2</chem>
25	H	0	CH <sub>2</sub> NHCH <sub>2</sub> CH <sub>2</sub>	H	<chem>CCN1CCN(CC1)Cc2ccccc2</chem>
26	H	0	C≡C	H	<chem>CCCCN1CCN(CC1)Cc2ccccc2</chem>
27	H	0	OCH <sub>2</sub>	H	<chem>CCCCN1CCN(CC1)Cc2ccccc2</chem>
28	H	0	CH=CH	H	<chem>CC#CCN1CCN(CC1)Cc2ccccc2</chem>

00956001-061600















Table 1 (continuation)

57

Nr	R <sup>1</sup> - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
29	H	0	CH=CH	H	$\text{CH}_2\text{CH}_2\text{CH}(\text{OH})\text{CH}_2-\text{N} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \text{N}-\text{CH}(\text{C}_6\text{H}_5)_2$
30	H	0	CH=CH	H	$\text{OCH}_2\text{CH}_2\text{CH}_2-\text{N} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \text{N}-\text{CH}(\text{C}_6\text{H}_5)_2$
31	H	0	CH=CH	H	$\text{CH}_2\text{CH}_2\text{CH}_2\text{C}(=\text{O})-\text{N} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \text{N}-\text{CH}(\text{C}_6\text{H}_5)_2$
32	H	0	CH=CH	H	$\text{CH}_2\text{CH}_2\text{CH}_2\text{SO}_2-\text{N} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \text{N}-\text{CH}(\text{C}_6\text{H}_5)_2$
33	H	0	CH=CH	H	$\text{CH}_2\text{CH}_2\text{NH}-\text{C}(=\text{O})-\text{N} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \text{N}-\text{CH}(\text{C}_6\text{H}_5)_2$
34	H	0	SCH <sub>2</sub>	H	$\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{N} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \text{N}-\text{CH}(\text{C}_6\text{H}_5)_2$
35	H	0	CH=CH	H	$\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_2-\text{N} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \text{N}-\text{CH}(\text{C}_6\text{H}_5)_2$
36	H	0	CH=CH	H	$\text{CH}_2\text{CH}_2\text{CH}_2\text{C}\equiv\text{CCH}_2-\text{N} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \text{N}-\text{CH}(\text{C}_6\text{H}_5)_2$
37	H	0	CH=CH	H	$\text{CH}_2\text{C}\equiv\text{C}-\text{CH}=\text{CHCH}_2-\text{N} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \text{N}-\text{CH}(\text{C}_6\text{H}_5)_2$
38	H	0	CH=CH	H	$\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{N} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \text{N}-\text{CH}(\text{C}_6\text{H}_5)_2$ <div style="text-align: center;"> </div>
39	H	0	CH=CH	H	$\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{N} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \text{N}-\text{CH}(\text{C}_6\text{H}_5)_2$ <div style="text-align: center;"> </div>
40	H	0	CH=CH	H	$\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{N} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \text{N}-\text{CH}(\text{C}_6\text{H}_5)_2$ <div style="text-align: center;"> </div>
41	H	0	CH=CH	H	$\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{N} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \text{N}-\text{CH}(\text{C}_6\text{H}_5)_2$ <div style="text-align: center;"> </div>

009596001, 061500

Table 1 (continuation)

Nr	R <sup>1</sup> -R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
42	H	0	CH=CH	H	 -CH(C <sub>6</sub> H <sub>5</sub> ) <sub>2</sub>
43	H	0	OCH <sub>2</sub>	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N  -CH <sub>2</sub> CH(C <sub>6</sub> H <sub>5</sub> ) <sub>2</sub>
44	H	0	CH=CH	H	CH <sub>2</sub> CH <sub>2</sub> -N  -CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH(C <sub>6</sub> H <sub>5</sub> ) <sub>2</sub>
45	H	0	CH=CH	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N  -CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH(C <sub>6</sub> H <sub>5</sub> ) <sub>2</sub>
46	S.F	0		H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N  -N 
47	H	0	CH=CH	H	CH <sub>2</sub> C≡CCH <sub>2</sub> -N  -N 
48	H	0	OCH <sub>2</sub>	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N  -N 
49	H	0		H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N  -N 

0956001.06160

Table 1 (continuation)

Nr	R <sup>1</sup> -R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
50	H	0	CH=CH	H	$\text{CH}_2\text{CH}_2\underset{\text{OH}}{\text{CH}}\text{CH}_2-\text{N} \begin{array}{c} \diagup \\ \diagdown \end{array} \begin{array}{c} \text{Cl} \\ \text{Cl} \end{array} \begin{array}{c} \text{C}_6\text{H}_4 \\ \text{C}_6\text{H}_4 \end{array}$
51	H	0	CH=CH	H	$\text{CH}_2\text{CH}_2\text{NH}-\text{C}(=\text{O})-\text{N} \begin{array}{c} \diagup \\ \diagdown \end{array} \begin{array}{c} \text{Cl} \\ \text{Cl} \end{array} \begin{array}{c} \text{C}_6\text{H}_4 \\ \text{C}_6\text{H}_4 \end{array}$
52	H	0	CH=CH	H	$\text{CH}_2\text{C}\equiv\text{CCH}_2-\text{N} \begin{array}{c} \diagup \\ \diagdown \end{array} \begin{array}{c} \text{Cl} \\ \text{Cl} \end{array} \begin{array}{c} \text{C}_6\text{H}_4 \\ \text{C}_6\text{H}_4 \end{array}$
53	H	0	CH=CH	H	$\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{N} \begin{array}{c} \diagup \\ \diagdown \end{array} \begin{array}{c} \text{OH} \\ \text{Cl} \end{array} \begin{array}{c} \text{C}_6\text{H}_4 \\ \text{C}_6\text{H}_4 \end{array}$
54	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	$\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{N} \begin{array}{c} \diagup \\ \diagdown \end{array} \begin{array}{c} \text{OH} \\ \text{Cl} \end{array} \begin{array}{c} \text{C}_6\text{H}_4 \\ \text{C}_6\text{H}_4 \end{array}$
55	H	0	CH=CH	H	$\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{N} \begin{array}{c} \diagup \\ \diagdown \end{array} \begin{array}{c} \text{COOH} \\ \text{Cl} \end{array} \begin{array}{c} \text{C}_6\text{H}_4 \\ \text{C}_6\text{H}_4 \end{array}$

009596001.051500

Table 1 (continuation)






60

Nr	R <sup>1</sup> -R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
56	H	0	CH=CH	H	
57	H	0	CH=CH	H	
58	6-C <sub>2</sub> H <sub>5</sub> S	0	CH=CH	H	
59	H	0		H	
60	H	0	OCH <sub>2</sub>	H	
61	6-CH <sub>3</sub>	0	CH=CH	H	
62	H	0	CH=CH	H	

0056001.061600

Table 1 (continuation)

61















Nr	R <sup>1</sup> - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
63	H	0	CH=CH	H	
64	H	0	CH2CH2	H	
65	H	0	CH2CH2	OH	"
66	H	0	CH2CH2	CH3	"
67	H	0	CH2CH2	$\begin{smallmatrix} C= \\   \\ CH=CH_2 \end{smallmatrix}$	"
68	H	0	CH=CH	H	"
69	H	1	CH=CH	H	"
70	5-F	0	CH=CH	H	"
71	6-CH3	0	CH=CH	H	"
72	6-CF3	0	CH=CH	H	"
73	H	0	OCH2	H	"
74	H	0	$\begin{smallmatrix} CH_2CH \\   \\ OH \end{smallmatrix}$	H	"
75	H	0	$\begin{smallmatrix} CH \equiv C \\   \\ CH_3 \end{smallmatrix}$	H	"
76	H	0	CH=CH-CH=CH	H	"
77	H	0	$\begin{smallmatrix} C=CH \\   \\ CH_3 \end{smallmatrix}$	H	
78	H	0	CH=CH-CH=CH	H	"
79	H	0	CH=CH	H	
80	H	0	CH2CH2	H	
81	H	0	CH=CH	H	"

00956001-051500



Table 1 (continuation)

62

Nr	R <sup>1</sup> - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
82	2-CH <sub>3</sub> O	0	CH=CH	H	"
83	H	0	CH=CH	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N  N 
84	6-CF <sub>3</sub>	0	CH=CH	H	"
85	H	0	CH=CH	H	CH <sub>2</sub> CH <sub>2</sub> -N  N 
86	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N  N 
87	H	0	CH=CH	H	"
88	H	1	CH=CH	H	"
89	6-CH <sub>3</sub> O	0	CH=CH	H	"
90	H	0	CH=CH	CH <sub>3</sub>	"
91	H	0	CH=CH-CH=CH	H	"
92	H	0	CH=CH	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N  N 
93	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N  N 
94	H	0	CH=CH	H	"
95	5-F	0	CH=CH	H	CH <sub>2</sub> CH=CHCH <sub>2</sub> -N  N 
96	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N  N 

09595001.081600

Table 1 (continuation)

Nr	R <sup>1</sup> - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
97	H	0	CH=CH	H	"
98	H	0	CH=CH	H	<chem>CH2CH2N1CCN(C1)c2c3ccccc3cc2</chem>
99	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	<chem>CH2CH2CH2CH2N1CCN(C1)c2c3ccccc3cc2</chem>
100	H	0	$\begin{array}{c} \text{CH}_2\text{CH} \\   \\ \text{OH} \end{array}$	H	"
101	H	0	CH=CH	H	"
102	6-C <sub>2</sub> H <sub>5</sub> S	0	CH=CH	H	"
103	5-F	0	CH=CH	H	"
104	H	1	CH=CH	H	"
105	H	0	CH <sub>2</sub> CH <sub>2</sub>	C <sub>2</sub> H <sub>5</sub>	"
106	H	0	OCH <sub>2</sub>	H	"
107	H	0	$\begin{array}{c} \text{CH}=\text{C} \\   \\ \text{CH}_3 \end{array}$	H	"
108	6-CH <sub>3</sub>	0	CH=CH	H	<chem>CH2CH=CHCH2N1CCN(C1)c2c3ccccc3cc2</chem>
109	H	0	CH=CH-CH=CH	H	"
110	H	0	CH=CH	H	<chem>CH2CH2CH2CH2CH2N1CCN(C1)c2c3ccccc3cc2</chem>

009190-10096660

Table 1 (continuation)





64

Nr	R <sup>1</sup> - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
111	H	0	CH=CH	H	<chem>CH2CH2OCH2CH2N1CCN(C1)c2c3ccccc3cc2</chem>
112	6-C <sub>6</sub> H <sub>5</sub> O	0	CH=CH	H	"
113	H	0	CH=CH	H	<chem>(CH2)6N1CCN(C1)c2c3ccccc3cc2</chem>
114	H	0	CH=CH	H	<chem>CH2CH2CH2CH2N1CCN(C1)c2c3ccccc3cc2C(=O)O</chem>
115	H	0	CH=CH	H	<chem>CH2CH2N1CCN(C1)c2c3ccccc3cc2</chem>
116	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	<chem>CH2CH2CH2CH2N1CCN(C1)c2c3ccccc3cc2</chem>
117	H	0	CH <sub>2</sub> CH <sub>2</sub>	OH	"
118	H	0	CH <sub>2</sub> CH   OH	H	"
119	H	0	CH=CH	H	"
120	H	1	CH=CH	H	"

00956001.061600

Table 1 (continuation)

65

Nr	R <sup>1</sup> - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
121	2,6-(CH <sub>3</sub> ) <sub>2</sub>	0	CH=CH	H	"
122	H	0	CH=C   C <sub>6</sub> H <sub>5</sub>	H	"
123	H	0	CH=CH-CH=CH	H	"
124	H	0	OCH <sub>2</sub>	H	
125	H	0	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub>	H	"
126	H	0	CH=CH	H	
127	H	0	CHCH <sub>2</sub>   CH <sub>3</sub>	H	
128	H	0	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub>	H	"
129	H	0	CH=CH	H	

00956001.061600






Table 1 (continuation)

66

Nr	R <sup>1</sup> -R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
130	H	0	CH=CH	H	
131	H	0	CH=CH	H	
132	H	0	CH=CH	H	
133	H	0	CH=CH	H	
134	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	
135	H	0	CH=CH	H	"
136	H	0	CH=CH	H	







00595001-061670

Table 1 (continuation)

Nr	R <sup>1</sup> - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
137	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	
138	H	0	CH=CH	H	
139	H	0	CH=CH	H	
140	H	0	OCH <sub>2</sub>	H	
141	H	0	CH=CH	H	"
142	H	0	CH=CH·CH=CH	H	"
143	H	0	CH=CH	H	

0956001.061600

Table 1 (continuation)

Nr	R <sup>1</sup> - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
144	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	
145	H	0	CH=CH	H	
146	H	0	CH=CH	H	
147	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	
148	H	0	CH=CH	H	
149	H	0	CH=CH	H	

09596001.061600

Table 1 (continuation)

69

Nr	R' - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
150	H	0	CH=CH	H	
151	H	0	CH=CH	H	
152	H	0	CH2CH2	H	
153	H	0	CH=CH	H	
154	H	0	CH=CH	H	
155	H	0	CH=CH	H	
156	H	0	OCH2	H	
157	H	0	CH=CH	H	
158	H	0	CH=CH	H	
159	H	0	CH=CH	H	

00596001-061600



Table 1 (continuation)

Nr	R <sup>1</sup> -R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
160	H	0	CH=CH	H	
161	H	0	CH=CH	H	
162	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	
163	H	0	CH=CH	H	
164	H	0	CH=CH	H	
165	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	
166	H	0	CH=CH	H	"
167	H	0	CH=CH	H	
168	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	
169	H	0	CH=CH	H	"
170	H	0	CH=CH	H	
171	H	0	CH=CH	H	
172	H	0	CH=CH	H	

00546001.061600

Table 1 (continuation)

71

Nr	R <sup>1</sup> -R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
173	H	0	CH=CH	H	
174	H	0	CH=CH	H	
175	H	0	CH2CH2	H	
176	H	0	CH=CH	H	
177	H	0	CH=CH	H	
178	H	0	CH2CH2	H	
179	H	0	CH=CH	H	
180	H	0	CH=CH	H	
181	H	0	CH=CH	H	
182	H	0	CH2CH2		
183	H	0	CH=CH	H	"

009180-009660-056600-001600

Table 1 (continuation)

Nr	R <sup>1</sup> -R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
184	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	
185	5-CF <sub>3</sub>	0	CH=CH	H	
186	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	
187	H	0	CH=CH:CH=CH	H	
188	H	0	CH=CH	H	
189	H	0	CH=CH	H	
190	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	
191	H	0	CH=CH	H	
192	H	0	CH=CH	H	
193	H	0	CH=CH	H	

Table 1 (continuation)

73

Nr	R <sup>1</sup> - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
194	H	0	CH=CH	H	<chem>CCCCN1CCN(CC1)C(=O)NC2CCCC2</chem>
195	H	0	CH=CH	H	<chem>CCCCN1CCN(CC1)C(=O)NC2c3ccccc3c4ccccc4c5ccccc25</chem>
196	H	0	CH=CH-CH=CH	H	<chem>CCCCN1CCN(CC1)C(=O)NC2=CC=CC=C2</chem>
197	H	0	CH=CH	H	<chem>CCCCN1CCN(CC1)C(=O)NC2CCCCC2</chem>
198	H	0	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub>	H	<chem>CCCCN1CCN(CC1)C(=O)NC2c3ccccc3c4ccccc24</chem>
199	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	<chem>CCCCN1CCN(CC1)C(=O)NC2c3ccccc3c4ccccc24</chem>
200	H	0	OCH <sub>2</sub>	H	<chem>CCCCN1CCN(CC1)C(=O)NC2c3ccccc3c4ccccc24</chem>
201	H	0	CH=CH	H	<chem>CCCCCN1CCN(CC1)C(=O)NC2c3ccccc3c4ccccc24</chem>
202	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	<chem>CCCCCN1CCN(CC1)C(=O)NC2c3ccccc3c4ccccc24</chem>

00573001-061600




Table 1 (continuation) 74

Nr	R <sup>1</sup> - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
203	H	0	CH=CH	H	
204	H	0	CH=CH	H	
205	6-CH <sub>3</sub>	0	CH=CH	H	
206	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	

05596001.061600

Table 1 (continuation)

75

Nr	R <sup>1</sup> - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
207	H	0	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub>	H	 <chem>c1ccccc1C(=O)N1CCN(C1)CCCC2=CN(C(=O)N2)C3=CC=CC=C3</chem>
205	H	0	CH=CH	H	 <chem>c1ccccc1C(=O)N1CCN(C1)CCCC2=CN(C(=O)N2)C3=CC=CC=C3</chem>
209	H	0	CH=CH	H	 <chem>c1ccccc1C(=O)N1CCN(C1)CCCC2=CN(C(=O)N2)C3=CC=CC=C3</chem>

00595001-051500

Table 1 (continuation)

Nr	R <sup>1</sup> - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
210	H	0	CH=CH	H	<chem>CCCCN1CCN(C1)C(=O)N2CCCC2</chem>
211	H	0	CH=CH	H	<chem>CCCCN1CCN(C1)C(=O)N2CCOCC2</chem>
212	H	0	CH=CH	H	<chem>CCCCN1CCN(C1)C(=O)N2c3ccccc3c2</chem>
213	H	0	CH=CH	H	<chem>CCN1CCN(C1)C(=O)N2c3ccccc3c2</chem>
214	H	0	CH=CH	H	<chem>CCCCN1CCN(C1)C(=O)N2c3ccccc3c2</chem>
215	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	<chem>CCN1CCN(C1)C(=O)N2c3ccccc3c2</chem>
216	H	c	CH=CH	H	<chem>(CCCCCCN1CCN(C1)C(=O)N2c3ccccc3c2</chem>
217	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	<chem>CCCCN1CCN(C1)C(=O)N2c3ccccc3c2</chem>

00595001.061600

Table 1 (continuation)

Nr	R <sup>1</sup> - R <sup>3</sup>	K	A	R <sup>4</sup>	D-E-G
218	H	0	CH=CH	H	
219	H	0	CH2CH2	H	
220	H	0	CH=CH	H	
221	H	0	CH=CH	H	
222	H	0	CH=CH:CH=CH	H	
223	H	0	CH2CH2	H	
224	H	0	CH=CH	H	
225	H	0	CH2CH2	H	"
226	H	0	CH2CH2	H	
227	H	0	CH=CH	H	"
228	H	0	CH=CH	H	



Table 1 (continuation)

Nr	R <sup>1</sup> - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
229	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N<img alt="piperidine ring" data-bbox="680 140 740 170"/>-N-SO <sub>2</sub> -<img alt="pyridine ring" data-bbox="800 140 860 170"/>
230	5-F	0	CH=CH	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N<img alt="piperidine ring" data-bbox="680 180 740 210"/>-N-SO <sub>2</sub> -<img alt="5-chloro-2-thienyl ring" data-bbox="760 180 870 210"/>
231	H	0	CH=CH	H	CH <sub>2</sub> CH <sub>2</sub> -N<img alt="piperidine ring" data-bbox="620 215 680 245"/>-N-SO <sub>2</sub> -<img alt="quinoline ring" data-bbox="750 215 810 275"/>
232	H	0	CH=CH	H	CH <sub>2</sub> CH <sub>2</sub> -N<img alt="piperidine ring" data-bbox="620 305 680 335"/>-N-P(=O)(<img alt="phenyl ring" data-bbox="715 280 775 315"/>)(<img alt="phenyl ring" data-bbox="715 325 775 360"/>)
233	H	0	OCH <sub>2</sub>	H	"
234	H	0	CH=CH	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N<img alt="piperidine ring" data-bbox="680 415 740 445"/>-N-P(=O)(<img alt="phenyl ring" data-bbox="770 385 830 420"/>)(<img alt="phenyl ring" data-bbox="770 425 830 460"/>)
235	H	0	CH=CH:CH=CH	H	"
236	6-CH <sub>3</sub>	0	CH=CH	H	CH <sub>2</sub> C≡CCH <sub>2</sub> -N<img alt="piperidine ring" data-bbox="680 510 740 540"/>-N-P(=O)(<img alt="phenyl ring" data-bbox="770 485 830 520"/>)(<img alt="phenyl ring" data-bbox="770 525 830 560"/>)
237	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N<img alt="piperidine ring" data-bbox="680 595 740 625"/>-N-P(=O)(<img alt="phenyl ring" data-bbox="795 565 855 600"/>)(<img alt="phenyl ring" data-bbox="795 605 855 640"/>)

00595001-061600

Table 1 (continuation)

Nr	R <sup>1</sup> - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
238	H	0	CH=CH	H	<chem>(CH2)6N1CCN(C1)P(=O)(c2ccccc2)c3ccccc3</chem>
239	H	0	CH=CH	H	<chem>Clc1ccc(cc1)P(=O)(c2ccccc2)N3CCN(CCC3)P(=O)(c4ccccc4)c5ccccc5</chem>
240	H	0	CH=CH	H	<chem>CC(F)(F)FOC(=O)N1CCN(CCC1)C2=CC=CC=C2</chem>
241	H	0	CH=CH-CH=CH	H	<chem>C=CCOC(=O)N1CCN(CCC1)C2=CC=CC=C2</chem>
242	H	0	CH2CH2	H	<chem>CC(C)(C)OC(=O)N1CCN(CCC1)C2=CC=CC=C2</chem>
243	H	0	CH=CH	H	"
244	H	0	CH=CH	H	<chem>CC(C)(C)OC(=O)N1CCN(CCC1)C2=CC=CC=C2</chem>
245	H	0		H	<chem>CC(C)(C)OC(=O)N1CCN(CCC1)C2=CC=CC=C2</chem>
246	H	0	C≡C	H	<chem>CC(C)(C)OC(=O)N1CCN(CCC1)C2=CC=CC=C2</chem>
247	H	0	CH=CH	H	<chem>CC(C)(C)OC(=O)N1CCN(CCC1)C2=CC=CC=C2</chem>
248	H	0	CH2CH2	H	<chem>c1ccccc1COC(=O)N1CCN(CCC1)C2=CC=CC=C2</chem>

00596001-061600

Table 1 (continuation)

80

Nr	R <sup>1</sup> - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
249	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	<chem>CCCCCN1CCN(C1)C(=O)OC2=CC=CC=C2</chem>
250	H	0	CH=CH	H	<chem>CCN1CCNCC1</chem>
251	H	0	CH=CH	H	<chem>CCCCN1CCNCC1</chem>
252	H	0	CH=CH	H	<chem>CCCN1CCN(C1)C2=CC=CC=C2</chem>
253	H	0	OCH <sub>2</sub>	H	<chem>CCCCN1CCN(C1)CC2=CC=CC=C2</chem>
254	H	0	CH=CH-CH=CH	H	<chem>CCN1CCN(C1)CC2=CC=CC=C2</chem>
255	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	<chem>CCCCN1CCN(C1)C(C)C2=CC=CC=C2</chem>
256	H	0	CH=CH	H	<chem>CCCCN1CCN(C1)CC2=CC=CC=C2</chem>
257	H	0	OCH <sub>2</sub>	H	<chem>CCCCN1CCN(C1)CC(C)(C)C2=CC=CC=C2</chem>
258	H	0	CH=CH	H	<chem>CC#CCN1CCN(C1)CC(C)(C)C2=CC=CC=C2</chem>
259	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	<chem>CCOCCN1CCN(C1)CC(C)(C)C2=CC=CC=C2</chem>

009596001.061600







Table 1 (continuation)

Nr	R <sup>1</sup> - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
250	H	0	CH <sub>2</sub> CHF	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N(CH(CH <sub>2</sub> C <sub>6</sub> H <sub>5</sub> ) <sub>2</sub> )
261	H	0	C≡C	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N(CH(CH <sub>2</sub> Cl)(CH <sub>2</sub> Cl))
262	H	0	CH=CH	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N(CH(CH <sub>2</sub> COOH)(CH <sub>2</sub> Cl))
263	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	CH <sub>2</sub> C≡CCH <sub>2</sub> -N(CH(CH <sub>2</sub> Cl)(CH <sub>2</sub> Cl))
264	H	0	CH=CH	H	CH <sub>2</sub> CH <sub>2</sub> -N(CH(CH <sub>2</sub> Cl)(CH <sub>2</sub> Cl))
265	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N(CH(CH <sub>2</sub> Cl)(CH <sub>2</sub> Cl))
266	H	0	CH=CH	H	"
267	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N(CH(CH <sub>2</sub> Cl)(CH <sub>2</sub> Cl))
268	H	0	CH=CH	H	"

00596001.061600

Nr	R <sup>1</sup> - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
269	H	0	$\text{CH}=\text{C}$   $\text{CH}_3$	H	"
270	H	0	$\text{CH}_2\text{CH}_2$	H	 $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{N}$
271	H	0	$\text{CH}=\text{CH}$	H	"
272	H	0	$\text{CH}=\text{CH}$	H	 $\text{CH}_2\text{CH}_2-\text{N}$
273	H	0	$\text{CH}_2\text{CH}_2$	H	 $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{N}$
274	H	0	$\text{OCH}_2$	H	"
275	H	0	$\text{CH}=\text{CH}$	H	"
276	$6\text{-CF}_3$	0	$\text{CH}=\text{CH}$	H	"
277	H	1	$\text{CH}=\text{CH}$	H	"
278	H	0	$\text{CH}=\text{CH}$	$\text{CH}_3$	"
279	H	0	$\text{CH}_2\text{CH}_2$	H	 $\text{CH}_2\text{CH}=\text{CHCH}_2-\text{N}$

[illegible]

Nr	R <sup>1</sup> - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
280	H	0	CH=CH	H	
281	H	0	CH=CH	H	
282	H	0	CH2CH2	H	
283	H	0	CH=CH	H	
284	H	0	CH2CH2	H	
285	H	0	CH=CH	H	

**06E01**

Table 1 (continuation)

Nr	R <sup>1</sup> - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
286	H	0	CH <sub>2</sub>	H	
287	H	0	CH=CH	H	
288	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	
289	H	0	CH=CH	H	
290	H	0	CH=CH	H	
291	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	
292	H	0	CH=CH	H	
293	H	0	CH=CH	H	
294	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	

Table 1 (continuation)

Nr	R' - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
255	H	0	CH=CH	H	<chem>CCCCN1CCN(C1)C(=O)Nc2ccc3ccccc3c2</chem>
256	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	<chem>CCCCN1CCN(C1)C(=O)Nc2cccnc2</chem>
297	H	0	CH=CH	H	<chem>CCCCN1CCN(C1)C(=O)N2c3ccccc3c4ccccc24</chem>
298	H	0	CH=CH	H	<chem>CCCCN1CCN(C1)C(=O)N2c3ccccc3c4cnc34</chem>
299	H	0	CH=CH	H	<chem>CCCCN1CCN(C1)C(=O)N2c3ccccc3c4c(=O)[nH]c4n2</chem>
300	H	0	OCH <sub>2</sub>	H	<chem>CCN1CCN(C1)S(=O)(=O)c2ccc(C)cc2</chem>
301	H	0	CH=CH	H	<chem>CCN1CCN(C1)S(=O)(=O)c2ccc3ccccc3c2</chem>
302	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	<chem>CCCCN1CCN(C1)S(=O)(=O)c2ccc3ccccc3c2</chem>

00596001.051600



Table 1 (continuation)

86







Nr	R <sup>1</sup> - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
303	H	0	CH=CH	H	
304	H	0	CH=CH	H	
305	H	0	CH=CH	H	
306	H	0	CH=CH	H	
307	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	
308	H	0	CH=CH	H	
309	H	0	CH=CH.CH=CH	H	"
310	H	0	CH=CH	H	

Table 1 (continuation)

Nr	R <sup>1</sup> - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
311	H	0	CH=CH	H	
312	H	0	CH=CH	H	
313	H	0	OCH <sub>2</sub>	H	
314	H	0	CH=CH	H	
315	H	0	CH=CH	H	
316	H	0	CH=CH	H	
317	H	0	CH=CH	H	
318	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	
319	H	0	CH=CH-CH=CH	H	"
320	H	0	CH=CH	CH <sub>3</sub>	

00596001.061600

Table 1 (continuation)

Nr	R <sup>1</sup> - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
321	H	0	CH=CH	H	$\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}-\overset{\text{O}}{\underset{\text{  }}{\text{C}}}-\text{N} \begin{array}{c} \diagup \diagdown \\ \text{---} \text{N} \text{---} \end{array} \text{CH}(\text{C}_6\text{H}_5)_2$
322	H	0	CH=CH	H	
323	H	0	CH2CH2	H	
324	H	0	CH=CH	H	"
325	H	0	CH=CH	H	
326	H	0	CH=CH	H	
327	H	0	CH2CH2	H	
328	H	0	CH=CH	H	

00505001-061600



Table 1 (continuation)

90









Nr	R <sup>1</sup> - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
338	H	0	SCH <sub>2</sub>	H	
339	H	0	SCH <sub>2</sub>	H	
340	H	0	CH=CH	H	
341	H	0	CH <sub>2</sub> CH <sub>2</sub>	CH <sub>3</sub>	
342	H	0	CH=CH	H	
343	H	0	SCH <sub>2</sub>	H	
344	H	0	CH=CH	CH <sub>3</sub>	
345	H	0	SCH <sub>2</sub>	H	

Table 1 (continuation)







91

Nr	R <sup>1</sup> - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
346	H	0	CH=CH	H	$\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}-\text{C}(=\text{O})-\text{N}(\text{piperazine})-\text{N}(\text{fluorenyl})$
347	H	0	CH=CH	H	$\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{N}(\text{piperazine})-\text{N}-\text{C}(=\text{O})-\text{C}_6\text{H}_4-\text{OH}$
348	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	$\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{N}(\text{piperazine})-\text{N}-\text{C}(=\text{O})-\text{C}_6\text{H}_3(\text{NH}_2)$
349	H	0	CH=CH	H	"
350	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	$\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{N}(\text{piperazine})-\text{N}-\text{C}(=\text{O})-\text{C}_6\text{H}_4-\text{COOH}$
351	H	0	CH=CH	H	"
352	H	0	CH=CH	H	$\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{N}(\text{piperazine})-\text{N}-\text{C}(=\text{O})-\text{C}_6\text{H}_4-\text{H}_2\text{NOC}$
353	H	0	CH=CH	H	$\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{N}(\text{piperazine})-\text{N}-\text{C}(=\text{O})-\text{C}_6\text{H}_4-\text{H}_3\text{CS}$
354	H	0	CH=CH	H	$\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{N}(\text{piperazine})-\text{N}-\text{C}(=\text{O})-\text{C}_6\text{H}_4-\text{CH}_2\text{CH}_2\text{COOH}$
355	H	0	CH=CH	H	$\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{N}(\text{piperazine})-\text{N}-\text{C}(=\text{O})-\text{C}_6\text{H}_3(\text{OH}, \text{NH}_2)$
356	H	0	CH=CH	H	$\text{CH}_2\text{CH}_2-\text{N}(\text{piperazine})-\text{N}-\text{C}(=\text{O})-\text{C}_6\text{H}_4-\text{N}(\text{fluorenyl})$

0355607-0516100

Table 1 (continuation)

92

Nr	R <sup>1</sup> - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
357	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	 <chem>CCCCN1CCN(C1=O)C2=CC=CC=C2</chem>
358	H	0	CH=CH	H	"
359	H	0	CH=CH	H	 <chem>CCCCN1CCN(C1=O)C2=CC=CC=C2</chem>
360	H	0	CH=CH	H	 <chem>CCCCN1CCN(C1=O)C2=CC=CC=C2</chem>
361	H	0	CH=CH-CH=CH	H	"
362	H	0	CH <sub>2</sub> CH <sub>2</sub>	H	 <chem>CCCCN1CCN(C1=O)C2=CC=CC=C2</chem>
363	H	0	CH=CH	H	"
364	H	0	CH=CH	H	 <chem>CCCCN1CCN(C1=O)C2=CC=CC=C2</chem>
365	H	0	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub>	H	 <chem>CCCCN1CCN(C1=O)C2=CC=CC=C2</chem>

059601.061600

Table 1 (continuation)

Nr	R <sup>1</sup> - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
366	H	0	CH=CH	H	<chem>CCCCN1CCN(C1)C(=O)CNc2c3ccccc3ccc2</chem>
367	H	0	CH=CH	H	<chem>CCOCCN1CCN(C1)C(=O)CNc2c3ccccc3ccc2</chem>
368	H	0	CH=CH	H	<chem>CCCCN1CCN(C1)C(=O)CNc2c3ccccc3ccc2</chem>
369	H	0	OCH <sub>2</sub>	H	"
370	H	0	CH=CH	H	<chem>CCCCN1CCN(C1)C(=O)CNc2c3ccccc3ccc2</chem>
371	H	0	SCH <sub>2</sub>	H	"
372	H	0	CH=CH	H	<chem>CCCCN1CCN(C1)C(=O)CNc2c3ccccc3ccc2</chem>
373	H	0	CH=CH	H	<chem>CCCCN1CCN(C1)C(=O)CNc2c3ccccc3ccc2</chem>




Table 1 (continuation)

Nr	R <sup>1</sup> - R <sup>3</sup>	k	A	R <sup>4</sup>	D-E-G
374	H	0	CH=CH	H	<chem>CCCC1CCN(C1)C(=O)OC(C)(C)C</chem>
375	H	0	SCH <sub>2</sub>	H	<chem>CCCCN1CCCN(C1)CCc2ccccc2c2ccccc2</chem>
376	H	0	CH=CH	CH <sub>3</sub>	<chem>CCN(C)C(=O)N1CCCN(C1)CCc2ccccc2c2ccccc2</chem>
377	H	0	CH=CH	H	<chem>CCCCN1CCCN(C1)C(=O)c2ccccc2N</chem>
378	H	0	CH=CH	H	<chem>CCCCN1CCCN(C1)C(=O)c2ccccc2C(=O)O</chem>
379	H	0	CH=CH	H	<chem>CCCCN1CCCN(C1)C(=O)c2c3ccccc3c4c2ccccc4N5C=CC=CC=C5</chem>
380	H	0	CH=CH	H	<chem>CCCCN1CCCN(C1)C(=O)N2C3C=CC=CC=C3C=Cc4ccccc42</chem>


Further subject-matter of the claims are known analogous methods for the production of the compounds of formula (I) according to the invention.

According to method variant (A), compounds of formula (I) are

- (a) obtained in the manner by reacting carboxylic acids of formula (II)



in which  $R^1$ ,  $R^2$ ,  $R^3$ , A and k have the meaning given above or their reactive derivatives with compounds of formula (III)



wherein D, E, G and  $R^4$  also have the above described meanings and the meanings given in the claims.

Reactive derivatives of compound (II) can be, for example, activated esters, anhydrides, acid halides (especially acid chlorides) or simple low alkyl esters. Suitable activated esters are, for example, p-nitrophenyl ester, 2,4,6-trichlorophenyl ester, pentachlorophenyl ester, cyanomethyl ester, esters of N-hydroxysuccinimide, N-hydroxyphthalimides, 1-hydroxybenzotriazole, N-hydroxypiperidine, 2-hydroxypyridine, 2-mercaptopyridine, etc. Anhydrides can be symmetric anhydrides or mixed, as they are obtained, for example, with

pivaloyl chloride or with chloroformates. Aromatic (for example chloroformic phenyl ester), araliphatic (for example chloroformic benzyl ester) or aliphatic chloroformates (for example chloroformic methyl ester, -ethyl ester or -isobutyl ester) can be used for this.

Reaction of compounds (II) with compounds (III) can also be carried out in the presence of condensation agents such as dicyclohexylcarbodiimide, 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide hydrochloride, N,N'-carbonyldiimidazole, 1-ethoxycarbonyl-2-ethoxy-1,2-dihydroquinoline, etc. If carbodiimides are used as the condensation agent, reagents such as N-hydroxysuccinimide, N-hydroxyphthalimide, 1-hydroxybenzotriazole, N-hydroxypiperidine, etc. can be advantageously added.

Compounds of formula (III) can be used for reaction as free bases as well as in the form of their acid addition salts. For this, the salts of inorganic acids are to be preferred, i.e. hydrochlorides, hydrobromides or sulfates for example.

Reaction of compounds (II) or their reactive derivatives with compounds (III) are normally carried out in a suitable, preferably inert solvent. As examples, aromatic hydrocarbons such as benzene, toluene, xylene, halogenated hydrocarbons (for example dichloromethane, chloroform, 1,2-dichloroethane, trichloroethylene), ethers (for example diethyl ether, tetrahydrofuran, dioxane, glycol dimethyl ether), ethyl acetate, acetonitrile or polar aprotic solvents such as, for example, dimethylsulfoxide, dimethylformamide or N-methylpyrrolidone are to be named. Pure solvents, as well as mixtures of two or more of them, can be used.

The reaction is optionally carried out in the presence of an auxiliary base. Suitable examples for this are alkali metal carbonates (sodium carbonate, potassium carbonate), alkali metal hydrogen carbonates (sodium hydrogen carbonate, potassium hydrogen carbonate), or organic bases such as, for

example, triethylamine, ethyl diisopropylamine, tributylamine, N-methylmorpholine or pyridine. A suitable excess of compound (III) can also be used as a base. If compounds (III) are used in form of their acid addition salts, then it is appropriate to consider the amount of auxiliary base used as equivalent.

The reaction temperatures can - depending on reactivity of the starting materials - vary in a wide range. Generally, the reaction is carried out at temperatures between -40°C and 180°C, preferably between -10°C and 130°C, especially at the boiling point of the solvent used.

The starting compounds and/or intermediates (II) and (III) are known and/or can be produced according to known methods in an analogous manner. Moreover, their production is further described below by means of representative examples.

Compounds of formula (I) can also be produced according to the variant pursuant to **Method B** by reaction of compounds of formula (I) wherein G is hydrogen, whereby the latter which themselves are anti-proliferative active ingredients according to the invention (as follows from the definitions for the general formula). Thereby, the reaction of compounds according to formula (I) occurs with a compound of formula (IV),



(IV)

in which G has the meaning given above, with the exception of hydrogen, and L represents a suitable nucleofuge or reactive group. The type of nucleofuge or reactive group L and the conditions of the reaction are dependent of the nature of group G. According to a further variant pursuant to **method (B1)** compounds of formula (I), in which G, with the exception of hydrogen, has the meaning of G<sup>1</sup> according to the above definition can also be synthesized by reacting compounds of formula (I), in which G is hydrogen, with a suitable

alkylation agent and/or arylation agent of formula (IV), wherein G is an alkyl-, alkenyl-, alkynyl-, cycloalkyl-, aryl-, aralkyl-, heteroaryl- or heteroaralkyl residue according to definition and the leaving group L can be a reactive derivative of an alcohol, for example, a halogen atom such as chlorine, bromine or iodine or a sulfonic acid ester, i.e. for example a methanesulfonyloxy group, trifluoromethanesulfonyloxy-, ethanesulfonyloxy-, benzene-sulfonyloxy-, p-toluenesulfonyloxy-, p-bromobenzene-sulfonyloxy-, m-nitrobenzenesulfonyloxy group, etc. A reactive group L can be a terminal epoxide group.


The reaction of compounds (I), in which G is a hydrogen, and (IV) is usually conducted in a suitably inert solvent. Such solvents can be for example, aromatic hydrocarbons (benzene, toluene, xylene), ethers (for example tetrahydrofuran, dioxane, glycol dimethyl ether), ethyl acetate, acetonitrile, ketones (acetone, ethyl methyl ketone), polar protic solvents such as alcohols (ethanol, isopropanol, butanol, glycol monomethyl ether) or polar aprotic solvents such as, for example, dimethylsulfoxide, dimethylformamide or N-methylpyrrolidone. Pure solvents as well as mixtures of two or more can also be used. Preferably, the reactions are carried out in the presence of bases, whereby the same bases as named in method (a) above can be used. If chlorides or bromides are used as compound (IV), the reaction can be accelerated by the addition of alkali metal iodides, for example sodium iodide, potassium iodide. The reaction temperatures can vary between 0°C and 180°C depending on the reactivity of the educts, but preferably lie between 20°C and 130°C. Finally, according to the variant pursuant to method (B2) compounds of formula (I), in which G represents an acyl residue, a carbamoyl residue, a sulfonyl residue or a phosphinoyl residue according to the above definition, can also be produced in a manner by reacting compounds of formula (I), wherein G is hydrogen, with a carboxylic acid, carbamic acid, sulfonic acid and/or phosphinic acid of formula (V),



(V)

wherein G is an acyl residue, carbamoyl residue, sulfonyl residue or phosphinoyl residue according to definition, or their derivatives capable of reaction. Preferred derivatives of carboxylic acids and/or sulfonic acids (V) which are capable of reaction are symmetric or unsymmetric carboxylic acid anhydrides and/or sulfonic acid anhydrides or acyl- and/or sulfonyl halides, especially acyl- and/or sulfonyl chlorides. Preferred derivatives of carbamates and/or phosphinic acids which are capable of reaction are the carbamoyl halides and/or phosphinyl halides, especially carbamyl- and/or phosphinyl chlorides. The reaction of the acids (V) and/or their reactive derivatives with compounds (I), in which G is hydrogen, preferably occurs in the presence of auxiliary bases in solvents and under conditions as they are described in method (A).

Compounds of formula (I), wherein G represents a carbamoyl residue according to the definition (G2b) with  $r = 0$ , i.e. is a group



can also be produced pursuant to the variant according to method (B3) by reacting compounds of formula (I), in which G is hydrogen, with a carbonyl group transmitter to an intermediate product and subsequently reacting this directly with a primary or secondary amine with the formula (VI)




(VI)


wherein  $R^{13}$  and  $R^{15}$  and/or the grouping  $—NR^{13}R^{15}$  have the above meanings without purifying or isolating the intermediate product.

Bis-trichloromethyl carbonate (triphosgene) and carbonyldiimidazole have been proven as particularly reactive carbonyl group transmitters. The reaction of compounds of formula (I), wherein G is hydrogen, with triphosgene and/or carbonyldiimidazole are typically conducted in an absolute, inert solvent in the presence of a tertiary organic amine as an auxiliary base in such a manner that the solution of compounds (I) and the auxiliary base are slowly poured into a solution of an equivalent amount of carbonyl group transmitter. Thereby, the reaction requires molar ratios of 1 : 1 for the reaction of compound (I) and carbonyldiimidazole, and, in contrast, a ratio of 1 : 0.35 for the use of triphosgene. After complete reaction of the components to the intermediate product, compound (VI) is added in stoichiometric amounts or in excess as a solution or a solid, whereby the reaction is typically completed at elevated temperature. Suitable inert solvents are, for example hydrocarbons such as hexane, heptane, benzene, toluene, xylene, chlorinated hydrocarbons (for example dichloromethane, chloroform, 1,2-dichloroethane, trichloroethylene), ethers (for example diethyl ether, tetrahydrofuran, dioxane), esters such as ethyl acetate, butyl acetate, acetonitrile or polar aprotic solvents such as formamide or dimethylformamide. Pure solvents as well as mixtures of various solvents can be used. Sometimes it is of advantage to carry out the first partial reaction at low temperature in a low-viscosity, highly-volatile solvent and to remove the solvent after formation of the intermediate and replace it by a higher boiling solvent. Amines such as for example triethylamine, ethyl diisopropylamine, tributylamine, N-methylmorpholine or pyridine are suitable as auxiliary bases.

If compounds (I) or (VI) are used as salts, the amount of the auxiliary base is increased accordingly. The reaction temperatures can lie between  $-40^{\circ}\text{C}$  and  $50^{\circ}\text{C}$  for the first partial reaction, preferably  $0^{\circ}\text{C}$  to  $30^{\circ}\text{C}$ , and between  $0^{\circ}\text{C}$  and  $150^{\circ}\text{C}$  for the second partial reaction, preferably  $20^{\circ}\text{C}$  to  $120^{\circ}\text{C}$ . Finally, according to variant pursuant to method (B4) compounds of formula (I), wherein G represents a carbamoyl residue according to the definition (G2b) with  $r = 0$  and  $R^{15} = \text{hydrogen}$ , i.e. the group represents



can also be produced by reacting the compounds of formula (I) in which G is hydrogen, with an isocyanate of formula (VII) in which  $R^{13}$  has the meaning according to definition



Reaction of the compounds of formula (I), in which G is hydrogen, with the isocyanates of formula (VII) are conducted thereby in an absolute, inert solvent which can be a hydrocarbon such as pentane, hexane, heptane, benzene, toluene, or xylene, chlorinated hydrocarbons (such as dichloromethane, chloroform, 1,2-dichloroethane, trichloroethylene), an ether (for example, diethyl ether, tetrahydrofuran, dioxane), esters such as ethyl acetate, butyl acetate, or polar aprotic solvents such as formamide or dimethylformamide. Mixtures of various solvents can also be used. Thereby, the reaction temperatures can vary in the region from  $-20^{\circ}\text{C}$  to  $150^{\circ}\text{C}$ , but preferably lie at  $20^{\circ}\text{C}$  to  $100^{\circ}\text{C}$ .

As already mentioned, the compounds of formula (I), wherein G is hydrogen, are themselves active ingredients according to



the invention with tumor growth inhibiting activity. However, independent of their therapeutic applicability, they also represent useful intermediate compounds for the production of a multitude of other compounds according to the invention corresponding to the method variants (B1) to (B4).

In principle, the compounds of the formula (I), in which G represents hydrogen, can be produced according to method A by reacting a carboxylic acid of formula (II) with amines of formula (III) in which G is hydrogen as described above. However, since the compounds of formula (III) with hydrogen as G represent  $\alpha,\omega$ -diamines, the formation of product mixtures is always to be expected in their reaction with carboxylic acids (II) or their reactive derivatives making a subsequent separation necessary.

In contrast, compounds of formula (I), in which G is hydrogen, are essentially more advantageously produced from other compounds of formula (I), in which G is a selectively cleavable group under mild conditions, i.e. corresponds to a nitrogen protective group.

In this connection, among the compounds according to formula (I) with tumor growth inhibiting properties, compounds in which G represents a benzyl group, a 4-methoxybenzyl group, a diphenylmethyl group, a triphenylmethyl group, a benzyloxy-carbonyl group, a methoxy- and/or ethoxycarbonyl group, a tert-butoxycarbonyl group, an allyloxycarbonyl group or a trifluoroacetyl group are particularly suitable. Thus, compounds of formula (I) with benzyl, diphenylmethyl, triphenylmethyl or benzyloxy-carbonyl groups as G can already be catalytically transformed into compounds of formula (I) with hydrogen as G at room temperature under mild conditions with elementary hydrogen or by transfer hydration. Compounds of formula (I) with a 4-methoxybenzyl group are converted to compounds of formula (I) with hydrogen as G by selective oxidation with ammonium-cer(IV)-nitrate. The cleavage of

simple alkoxy carbonyl groups such as the methoxy- or ethoxy carbonyl group as well as the trifluoroacetyl group as G in compounds of formula (I) succeed by alkali hydrolysis under mild conditions without cleaving the A and D linked amide function. This is suitably valid for the cleavage of the triphenylmethyl group and the tert-butoxycarbonyl group in the form of G in compounds of formula (I) which occurs in acidic medium under mild conditions. Finally, compounds of formula (I) with an allyloxycarbonyl group as the meaning for G can be converted into such with hydrogen as the meaning for G in neutral medium with palladium catalyst. All these methods are fully familiar to the person skilled in the art, and are furthermore also documented in various monographs, see for example Greene, Wuts: Protective Groups in Organic Synthesis, New York, 1991.

The compounds of formula (I) produced according to the methods (A) to (E) can be isolated and purified in a known manner, for example by subjecting the residue after distillation of the solvent to partition, extraction, re-precipitation or re-crystallization or another purification method. For this, column chromatography on a suitable support or preparative middle or high pressure liquid chromatography (HPLC) are preferred for this.

The compounds (I) are first normally obtained in form of their free bases or their hydrates or solvates, depending on the type of isolation and purification. Their addition salts with pharmaceutically suitable acids are obtained in a typical manner by converting the base with the desired acid in a suitable solvent. Depending on the number of basic centers of compounds (I), one or more equivalent acids per mole of base can be bound.

Suitable solvents are, for example, chlorinated hydrocarbons such as dichloromethane or chloroform; ethers such as diethyl ether, dioxane or tetrahydrofuran; acetonitrile; ketones such as acetone or ethyl methyl ketone; esters such as methyl

The bases can be recovered from the salts by alkalization, for example with aqueous ammonia solution, alkali carbonate or diluted sodium hydroxide solution.

SYNTHETIC EXAMPLES  
for the

In the production examples for the end products, the abbreviations stand for the following terms:

MP = melting point,  
RT = room temperature,  
MPLC = intermediate pressure liquid chromatography  
THF = tetrahydrofuran,  
DMF = dimethylformamide,  
abs. = absolute,  
CDI = carbonyldiimidazole,  
EDC = N-(3-dimethylaminopropyl)-N'-ethyl-carbodiimide  
hydrochloride,  
HOBT = 1-hydroxybenzotriazole,  
TEA = triethylamine.

$^1\text{H}$ -NMR-Spectrum = proton resonance spectrum, taken at 100 MHz. The chemical shifts are given in ppm against TMS as a standard ( $\delta = 0.0$ ), whereby

s = singlet,  
 d = doublet,  
 t = triplet,  
 dt = doublet-triplet,  
 m = multiplet,  
 ar = aromatic,  
 py = pyridine.

#### Example 1

N-[3-(4-diphenylmethylpiperazin-1-yl)-propoxyl]-3-pyridin-3-yl-acrylamide (substance 30)

3.8 g (22,9 mmol) of 3-(3-pyridyl)-acrylic acid are suspended in 40 ml absolute dichloromethane and after addition of two drops pyridine, are cooled to ca.  $0^\circ\text{C}$  in an ice bath under moisture exclusion. 5 ml (58,5 mmol) of oxalyl chloride are slowly added and the mixture is first stirred for 30 min under ice cooling and then stirred overnight at RT. Subsequently, the solvent and excess oxalyl chloride are distilled off on a rotary evaporator. In order to completely remove the oxalyl chloride, the colorless residue is dried further for two hours under high-vacuum. The acid chloride obtained in this manner is suspended without further purification in 30 ml absolute dichloromethane and cooled to ca.  $0^\circ\text{C}$  in an ice bath under moisture exclusion. 7.45 g (22,9 mmol) 3-(4-diphenylmethyl-piperazinyl)-propyl-hydroxylamine are dissolved in 40 ml absolute dichloromethane and are added dropwise to this suspension. After complete addition, the ice bath is removed and the reaction is stirred for a further two hours at RT. The mixture is subsequently concentrated, taken up in 10% sodium hydroxide solution and extracted three times with acetic acid ethyl ester. The combined organic phases are washed with saturated NaCl solution, dried over sodium sulfate and the solvent is

removed under vacuum. The residue is chromatographically purified over silica gel with  $\text{CHCl}_3/\text{CH}_3\text{OH}$  (99/1 to 85/15) and crystallized twice from acetic ethyl acid ester after evaporation of the solvent: Colorless crystals with a MP. of 115 - 117°C; yield 3.66 g (35%).

$\text{C}_{28}\text{H}_{32}\text{N}_4\text{O}_2$  (456.6)

IR-Spectrum (KBr):

n(NH)	3200 $\text{cm}^{-1}$
n(C=O)	1660 $\text{cm}^{-1}$
n(C=C)	1630 $\text{cm}^{-1}$

$^1\text{H}$ -NMR-Spectrum (DMSO- $\text{D}_6$ ):

1.50 - 1.90	(2H, m, C- $\text{CH}_2$ -C)
2.05 - 2.65	(10H, m, piperazine, N- $\text{CH}_2$ )
3.86	(2H, t, $\text{OCH}_2$ , J=6.1 Hz)
4.24	(1H, s, $\text{Ar}_2\text{CH}$ )
6.54	(1H, d, $\text{CH}=\text{CHCO}$ , J=16.0 Hz)
7.05 - 7.70	(13H, m, Ar, Py, NH, CH=CHCO)
7.90 - 8.15	(1H, m, Py)
8.50 - 8.70	(1H, m, Py)
8.70 - 8.90	(1H, m, Py)

## Example 2

N-[4-(4-diphenylphosphinoyl-piperazin-1-yl)-butyl]-3-pyridin-3-yl-acrylamide (substance 234)

2.7 g (18 mmol) 3-(3-pyridyl)-acrylic acid and 3.1 g (19 mmol) CDI are heated under reflux in 100 ml absolute THF under moisture exclusion. After an hour, this is cooled to RT and 7.0 g (19 mmol) 4-(4-diphenyl-phosphinoyl-piperazin-1-yl)-butylamine, dissolved in 30 ml absolute THF, is added dropwise. Following addition, this is further stirred for three hours at RT and left standing overnight. The mixture is poured into 150 ml water and extracted three times with acetic acid ethyl ester by shaking. The combined organic

phases are washed with saturated NaCl solution, dried over sodium sulphate and the solution is removed under vacuum. The residue is chromatographically pre-purified over silica gel with  $\text{CHCl}_3/\text{CH}_3\text{OH}$  (85/15). After removal of the solvent, the colorless oily residue is further purified by preparative middle pressure chromatography with  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  (93/7): yield 3.5 g (40%) as amorphous solid material.

$\text{C}_{28}\text{H}_{33}\text{N}_4\text{O}_2\text{P}$  (488.5)

$^1\text{H}$ -NMR-Spectrum ( $\text{CDCl}_3$ ):

1.40 - 1.80 (4H, m, C- $\text{CH}_2$ - $\text{CH}_2$ -C)  
 2.20 - 2.60 (6H, m, piperazine, N- $\text{CH}_2$ )  
 2.90 - 3.25 (4H, m, piperazine)  
 3.25 - 3.55 (2H, m,  $\text{CONHCH}_2$ )  
 6.53 (1H, d,  $\text{CH}=\text{CHCO}$ ,  $J=15.7$  Hz)  
 6.70 - 6.95 (1H, m, NH)  
 7.00 - 8.00 (13H, m, Ar, Py,  $\text{CH}=\text{CHCO}$ )  
 8.54 (1H, dd, Py,  $J=1.4$  Hz,  $J=4.8$  Hz)  
 8.70 (1H, d, Py,  $J=1.8$  Hz)

### Example 3

N-[4-(4-diphenylmethyl-piperazin-1-yl)-3-hydroxy-butyl]-3-pyridin-3-yl-acrylamide (substance 29)

Production analogues to Example 2.

Batch size: 2.0 g (13.3 mmol) 3-(3-pyridyl)-acrylic acid, 2.4 g (14.6 mmol) CDI and 4.5 g (13.3 mmol) 4-(4-diphenyl-methyl-piperazin-1-yl)-3-hydroxy-butylamine. The addition of the amine occurs at  $-10^\circ\text{C}$ . Subsequently, this is stirred further for an hour at  $0^\circ\text{C}$ .

In the purification, chromatography first occurs with  $\text{CHCl}_3/\text{CH}_3\text{OH}$  (95/5); subsequently, crystallization occurs once from 20 ml ethanol and twice each from 50 ml acetic acid

ethyl ester: Colorless crystals of MP. 168 - 171°C; yield 0.4 g (6%).

$C_{29}H_{34}N_4O_2$  (470.6)

IR-Spectrum (KBr):

n(NH)	3270 $cm^{-1}$
n(C=O)	1660, 1565 $cm^{-1}$
n(C=C)	1615 $cm^{-1}$

$^1H$ -NMR-Spectrum ( $CDCl_3$ ):

1.25 - 2.00	(2H, m, C-CH <sub>2</sub> -C)
2.05 - 2.85	(10H, m, piperazine, N-CH <sub>2</sub> )
3.00 - 3.60	(2H, m, CH-OH)
3.60 - 4.00	(2H, m, CONHCH <sub>2</sub> )
4.22	(1H, s, Ar <sub>2</sub> CH)
6.44	(1H, d, CH=CHCO, J=15.7 Hz)
6.60 - 8.85	(1H, m, NH)
6.95 - 7.60	(11H, m, Ar, Py)
7.58	(1H, d, CH=CHCO, J=15.7 Hz)
7.65 - 7.90	(1H, m, Py)
8.45 - 8.65	(1H, m, Py)
8.65 - 8.85	(1H, m, Py)

#### Example 4

N-[4-(4-diphenylmethyl-piperazin-1-yl)-4-oxo-butyl]-3-pyridin-3-yl-acrylamide (substance 31)

Production analogous to Example 1.

Batch size: 2.3 g (15.4 mmol) 3-(3-pyridyl)-acrylic acid, 4 ml (46.8 mmol) oxalyl chloride and 5.2 g (15.4 mmol) 4-(4-diphenyl-methyl-piperazin-1-yl)-4-oxo-butylamine.

In the purification, chromatography first occurs with  $CHCl_3/CH_3OH$  (90/10); subsequently, crystallization occurs twice from 400 ml acetic acid ethyl ester and 300 ml ethyl methyl ketone: colorless crystals of MP. 179 - 180°C; yield 3.2 g (45%).

$C_{29}H_{32}N_4O_2$  (468.6)

IR-Spectrum (KBr):

n(NH) 3240  $cm^{-1}$   
 n(C=O) 1665, 1550  $cm^{-1}$   
 n(C=C) 1630  $cm^{-1}$

$^1H$ -NMR-Spectrum ( $CDCl_3$ ):

1.70 - 2.10 (2H, m, C-CH<sub>2</sub>-C)  
 2.15 - 2.65 (6H, m, piperazine,  
 CO-CH<sub>2</sub>)  
 3.20 - 3.80 (6H, m, CONHCH<sub>2</sub>,  
 piperazine)  
 4.20 (1H, s, Ar<sub>2</sub>CH)  
 6.45 (1H, d, CH=CHCO, J=15.7 Hz)  
 6.75 - 7.00 (1H, m; NH)  
 7.05 - 7.55 (11H, m, Ar, Py)  
 7.58 (1H, d, CH=CHCO, J=15.7 Hz)  
 7.70 - 7.90 (1H, m, Py)  
 8.45 - 8.65 (1H, m, Py)  
 8.65 - 8.85 (1H, m, Py)

#### Example 5

N-[3-(4-diphenylmethyl-piperazin-1-yl-sulfonyl)-propyl]-3-pyridin-3-yl-acrylamide (substance 32)

Production analogous to Example 1.

Batch size: 0.5 g (3.3 mmol) 3-(3-pyridyl)-acrylic acid, 2 ml (23.4 mmol) oxalyl chloride and 1.24 g (3.3 mmol) 3-(4-diphenyl-methyl-piperazin-1-yl-sulfonyl)-propylamine.

In the purification, chromatography first occurs  $CHCl_3/CH_3OH$  (95/5); subsequently, crystallization occurs from 75 ml acetic acid ethyl ester: colorless crystals of MP. 167 - 168°C; yield 0.7 g (84%).

$C_{28}H_{32}N_4O_3S$  (504.6)

IR-Spectrum (KBr):



n(NH)	3360 cm <sup>-1</sup>
n(C=O)	1660, 1540 cm <sup>-1</sup>
n(C=C)	1630 cm <sup>-1</sup>

<sup>1</sup>H-NMR-Spectrum (CDCl<sub>3</sub>):

2.00 - 2.35	(2H, m, C-CH <sub>2</sub> -C)
2.35 - 2.65	(4H, m, piperazine)
3.00	(2H, t, SO <sub>2</sub> -CH <sub>2</sub> , J=7.1 Hz)
3.20 - 3.40	(4H, m, piperazine)
3.57	(2H, dt, CONHCH <sub>2</sub> , J=6.4 Hz, J=12.8 Hz)
4.27	(1H, s, Ar <sub>2</sub> CH)
6.10 - 6.35	(1H, m, NH)
6.47	(1H, d, CH=CHCO, J=15.7 Hz)
7.05 - 7.55	(11H, m, Ar, Py)
7.63	(1H, d, CH=CHCO, J=15.7 Hz)
7.70 - 7.95	(1H, m, Py)
8.50 - 8.65	(1H, m, Py)
8.70 - 8.85	(1H, m, Py)

## Example 6

N-{2-[2-(4-diphenylmethyl-piperazin-1-yl)-ethoxy]-ethyl}-3-pyridin-3-yl-acrylamide trihydrochloride (substance 35 as trihydrochloride)

Production analogous to Example 1.

Batch size: 6.0 g (40 mmol) 3-(3-pyridyl)-acrylic acid, 9.4 ml (110 mmol) oxalyl chloride and 12.4 g (36.5 mmol) 2-[2-(4-diphenylmethyl-piperazin-1-yl)-ethoxy]-ethylamine. The crude product is chromatographically pre-purified over silica gel with CHCl<sub>3</sub>/CH<sub>3</sub>OH (98/2 to 95/5). After removal of the solvent, the residue is dissolved in isopropanol and mixed with isopropanolic HCl solution. The mixture is rotated in and the HCl salt is crystallized from 50 ml methanol/6 drops diisopropyl ether: Colorless crystals of MP. 157 - 159°C; yield 0.6 g (3%).

C<sub>29</sub>H<sub>34</sub>N<sub>4</sub>O<sub>2</sub> • 3HCl (580.0)

## IR-Spectrum (KBr):

n(NH)	3240 $\text{cm}^{-1}$
n(C=O)	1670, 1550 $\text{cm}^{-1}$
n(C=C)	1630 $\text{cm}^{-1}$

 $^1\text{H}$ -NMR-Spectrum ( $\text{CD}_3\text{OD}$ ):

3.40 - 4.00	(16H, m, piperazine, CH <sub>2</sub> -CH <sub>2</sub> -O-CH <sub>2</sub> -CH <sub>2</sub> )
5.58	(1H, s, Ar <sub>2</sub> CH)
7.14	(1H, d, CH=CHCO, J=15.8 Hz)
7.30 - 7.95	(10H, m, Ar)
7.67	(1H, d, CH=CHCO, J=15.8 Hz)
8.10 - 8.25	(1H, m, Py)
8.80 - 9.00	(2H, m, Py)
9.15 - 9.20	(1H, m, Py)

## Example 7

N-{4-[4-(bis-(4-fluorophenyl)-methyl)-piperazin-1-yl]-but-2-in-yl}-3-pyridin-3-yl-acrylamide•trihydrochloride (substance 47 as trihydrochloride)

Production analogous to Example 1.

Batch size: 2.5 g (16.9 mmol) 3-(3-pyridyl)-acrylic acid, 2 ml (23 mmol) oxalyl chloride and 6.0 g (16.9 mmol) 4-{4-[bis-(4-fluorophenyl)-methyl]-piperazin-1-yl}-but-2-inylamine. The crude product is chromatographically pre-purified over silica gel with  $\text{CHCl}_3/\text{CH}_3\text{OH}$  (95/5). After removal of the solvent, the residue is dissolved in methanol and mixed with methanolic HCl solution. The drawn off HCl salt is first crystallized from isopropanol and subsequently from ethanol/diisopropylether. Colorless crystals of MP. 160 - 163°C; yield 3.5 g (35%).

$\text{C}_{29}\text{H}_{28}\text{F}_2\text{N}_4\text{O} \cdot 3\text{HCl}$  (595.9)

## IR-Spectrum (KBr):

n(NH)	3240 $\text{cm}^{-1}$
n(C=O)	1670, 1550 $\text{cm}^{-1}$
n(C=C)	1630 $\text{cm}^{-1}$

<sup>1</sup>H-NMR-Spectrum (D<sub>2</sub>O):

2.95 - 3.55	(8H, m, piperazine)
3.80 - 4.10	(4H, m, CH <sub>2</sub> -CC-CH <sub>2</sub> )
5.04	(1H, s, Ar <sub>2</sub> CH)
6.72	(1H, d, CH=CHCO, J=15.9 Hz)
6.85 - 7.60	(9H, m, Ar, CH=CHCO)
7.80 - 8.00	(1H, m, Py)
8.50 - 8.70	(2H, m, Py)
8.70 - 8.85	(1H, m, Py)

## Example 8

3-pyridin-3-yl-N-{4-[4-(1,2,3,4-tetrahydronaphthalin-1-yl)-piperazin-1-yl]-butyl}-acrylamide (substance 87)

Production analogous to Example 2.

Batch size: 2.7 g (18 mmol) 3-(3-pyridyl)-acrylic acid, 3.15 g (19 mmol) CDI and 5.0 g (17.4 mmol) 4-[4-(1,2,3,4-tetrahydro-naphthalin-1-yl)-piperazin-1-yl]-butylamine.

In the purification, chromatography first occurs with CHCl<sub>3</sub>/CH<sub>3</sub>OH (95/5 to 90/10); subsequently, crystallization occurs twice each from 40 ml 1-chlorobutane: Colorless crystals of MP. 110 - 114°C; yield 3.7 g (50%).

C<sub>26</sub>H<sub>34</sub>N<sub>4</sub>O (418,6)

## IR-Spectrum (KBr):

n(NH)	3300 cm <sup>-1</sup>
n(C=O)	1650, 1530 cm <sup>-1</sup>
n(C=C)	1620 cm <sup>-1</sup>

<sup>1</sup>H-NMR-Spectrum (CDCl<sub>3</sub>):

1.40 - 2.10	(8H, m, C-CH <sub>2</sub> -CH <sub>2</sub> -C, cyclohexyl)
2.25 - 3.10	(12H, m, piperazine, N-CH <sub>2</sub> , cyclohexyl)
3.30 - 3.60	(2H, m, CONHCH <sub>2</sub> )
3.70 - 4.00	(1H, m, cyclohexyl)
6.50	(1H, d, CH=CHCO, J=15.7 Hz)

6.90 - 7.45 (5H, m, Ar, Py, NH)  
 7.62 (1H, d,  $\text{CH}=\text{CHCO}$ ,  $J=15.7$  Hz)  
 7.50 - 7.90 (2H, m, Py, Ar)  
 8.50 - 8.65 (1H, m, Py)  
 8.70 - 8.80 (1H, m, Py)

### Example 9

3-pyridin-3-yl-N-[4-(4-(5,6,7,8-tetrahydro-naphthalin-1-yl)-piperazin-1-yl)-butyl]-acrylamide (substance 94)

1.6 g (11.1 mmol) 3-(3-pyridyl)-acrylic acid and 6.2 ml (44.3 mmol) TEA are suspended in 80 ml absolute dichloromethane and cooled to ca. 0°C under moisture exclusion. 2.0 g (12.1 mmol) 81% HOBT and 2.3 g (12.1 mmol) EDC are added and the mixture is stirred for 30 min under ice cooling. 4.0 g (10.1 mmol) 4-[4-(5,6,7,8-tetrahydro-naphthalin-1-yl)-piperazin-1-yl]-butylamine are added and the mixture is stirred without cooling overnight at RT. Subsequently, the batch is washed twice with 25 ml 2M sodium hydroxide solution and 25 ml water. The organic phase is dried over sodium sulfate and the solvent is removed under vacuum. The residue is chromatographically pre-purified over silica gel with  $\text{CHCl}_3/\text{CH}_3\text{OH}$  (95/5) and crystallized twice from acetonitrile (25 ml and 15 ml): Colorless crystals of MP. 108-109°C; yield 2.7 g (64%).

$\text{C}_{26}\text{H}_{34}\text{N}_4\text{O}$  (418.6)

IR-Spectrum (KBr):

n(NH)	3260 $\text{cm}^{-1}$
n(C=O)	1650, 1555 $\text{cm}^{-1}$
n(C=C)	1620 $\text{cm}^{-1}$

$^1\text{H}$ -NMR-Spectrum ( $\text{CDCl}_3$ ):

1.40 - 2.10 (8H, m, $\text{C}-\text{CH}_2-\text{CH}_2-\text{C}$ , cyclohexyl)
2.25 - 3.30 (14H, m, piperazine, $\text{N}-\text{CH}_2$ , cyclohexyl)

3.30 - 3.70 (2H, m, CONHCH<sub>2</sub>)  
 6.51 (1H, d, CH=CHCO, J=15.6 Hz)  
 6.70 - 7.45 (5H, m, Ar, Py, NH)  
 7.63 (1H, d, CH=CHCO, J=15.6 Hz)  
 7.70 - 7.95 (1H, m, Py)  
 8.45 - 8.65 (1H, m, Py)  
 8.65 - 8.85 (1H, m, Py)

#### Example 10

N-{4-[(naphthalin-1-yl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-acrylamide (substance 97)

Production analogous to Example 1.

Batch size: 3.38 g (22.7 mmol) 3-(3-pyridyl)-acrylic acid, 7.85 g (61.8 mmol) oxalyl chloride and 8.55 g (20.6 mmol) 4-[(naphthalin-1-yl)-piperazin-1-yl]-butylamine. In the recovery, 40 ml 10% sodium hydroxide solution is added to the reaction solution. The aqueous phase is extracted with dichloromethane. The combined organic phases are washed with 15 ml water, dried over sodium sulfate and the solvent is removed under vacuum. The residue is chromatographically purified over silica gel with CHCl<sub>3</sub>/CH<sub>3</sub>OH (98/2 to 96/4). After removal of the solvent, this is crystallized three times each with 40 ml acetic acid ethyl ester under addition of 5 drops diisopropylether respectively: solid with MP. 124-125°C; yield 1.5 g (18%).

C<sub>26</sub>H<sub>30</sub>N<sub>4</sub>O (414.6)

IR-Spectrum (KBr):

n(NH)	3280 cm <sup>-1</sup>
n(C=O)	1650, 1545 cm <sup>-1</sup>
n(C=C)	1620 cm <sup>-1</sup>

<sup>1</sup>H-NMR-Spectrum (CDCl<sub>3</sub>):

1.45 - 1.95	(4H, m, C-CH <sub>2</sub> -CH <sub>2</sub> -C)
2.30 - 3.40	(10H, m, piperazine, N-CH <sub>2</sub> )
3.30 - 3.60	(2H, m, CONHCH <sub>2</sub> )

6.50 (1H, d, CH=CHCO, J=15.7 Hz)  
 6.55 - 6.85 (1H, m, NH)  
 6.95 - 7.95 (8H, m, Ar, Py)  
 7.63 (1H, d, CH=CHCO, J=15.7 Hz)  
 8.05 - 8.35 (1H, m, Py)  
 8.50 - 8.70 (1H, m, Py)  
 8.70 - 8.85 (1H, m, Py)

# Example 11

N-{2-[4-(6,11-dihydrodibenzo[b,e]thiepin-11-yl)-piperazin-1-yl]-ethyl}-3-pyridin-3-yl-acrylamide (substance 138)

5.0 g (14.1 mmol) N-[2-(piperazin-1-yl)-ethyl]-3-pyridin-3-yl-acrylamidetrihydrochloride (substance 317 as trihydrochloride) and 5.8 ml (42.3 mmol) TEA are suspended in 65 ml absolute dichloromethane. 4.7 g (15.5 mmol) 11-methanesulfonyloxy-6,11-dihydrodibenzo[b,e]thiepine dissolved in 60 ml absolute dichloromethane is added dropwise under moisture exclusion. The mixture is stirred overnight at RT. Subsequently, the batch is washed three times each with 50 ml water. The organic phase is dried over sodium sulfate and the solvent is removed under vacuum. The residue is chromatographically purified over silica gel with CHCl<sub>3</sub>/CH<sub>3</sub>OH (95/5). Subsequently, a further purification occurs by means of MPLC with CHCl<sub>3</sub>/CH<sub>3</sub>OH (95/5): yield 4.3 g (44%) as an amorphic solid.

C<sub>28</sub>H<sub>30</sub>N<sub>4</sub>OS (470,6)

IR-Spectrum (KBr):

n(NH)	3270 cm <sup>-1</sup>
n(C=O)	1655, 1535 cm <sup>-1</sup>
n(C=C)	1620 cm <sup>-1</sup>

<sup>1</sup>H-NMR-Spectrum (CDCl<sub>3</sub>):

2.00 - 2.90 (10H, m, Piperazine,  
 Piperazin-CH<sub>2</sub>)  
 3.20 - 3.70 (3H, m, CONHCH<sub>2</sub>, SCH<sub>2</sub>)  
 4.10 (1H, s, Ar<sub>2</sub>CH)

5.90 - 6.50 (2H, m, NH, SCH<sub>2</sub>)  
 6.49 (1H, d, CH=CHCO, J=15.7 Hz)  
 6.90 - 7.50 (9H, m, Ar, Py)  
 7.62 (1H, d, CH=CHCO, J=15.7 Hz)  
 7.75 - 7.95 (1H, m, Py)  
 8.50 - 8.70 (1H, m, Py)  
 8.70 - 8.90 (1H, m, Py)

#### Example 12

N-[4-(4-biphenyl-2-yl-piperazin-1-yl)-butyl]-3-pyridin-3-yl-acrylamide (substance 68)

Production analogous to Example 1.

Batch size: 5.8 g (39.1 mmol) 3-(3-pyridyl)-acrylic acid, 9.1 ml (106 mmol) oxalyl chloride and 5.0 g (16.1 mmol) 4-(4-biphenyl-2-yl-piperazin-1-yl)-butylamine.

In the recovery, 60 ml 10% sodium hydroxide solution is added to the reaction solution. The aqueous phase is extracted twice each with 15 ml dichloromethane. The combined organic phases are washed twice each with 15 ml water, dried over sodium sulfate and the solvent is removed under vacuum. The residue is chromatographically purified over silica gel with CHCl<sub>3</sub>/CH<sub>3</sub>OH/NH<sub>4</sub>OH (85/15/2). After removal of the solvent, this is crystallized twice with 1-chlorobutane: solid with MP. 115°C; yield 3.3 g (46%).

C<sub>28</sub>H<sub>32</sub>N<sub>4</sub>O (440.6)

IR-Spectrum (KBr):

n(NH)	3280 cm <sup>-1</sup>
n(C=O)	1650, 1545 cm <sup>-1</sup>
n(C=C)	1620 cm <sup>-1</sup>

<sup>1</sup>H-NMR-Spectrum (CDCl<sub>3</sub>):

1.35 - 1.85 (4H, m, C-CH<sub>2</sub>-CH<sub>2</sub>-C)  
 2.10 - 2.55 (6H, m, piperazine,  
 N-CH<sub>2</sub>)  
 2.75 - 3.00 (4H, m, piperazine)  
 3.20 - 3.50 (2H, m, CONHCH<sub>2</sub>)  
 6.44 (1H, d, CH=CHCO, J=15.6 Hz)

6.45 - 6.70 (1H, m, NH)  
 6.95 - 7.85 (12H, m, Ar,  $\text{CH}=\text{CHCO}$ , Py)  
 8.45 - 8.65 (1H, m, Py)  
 8.65 - 8.80 (1H, m, Py)

### Example 13

N-{4-[4-(9H-fluorenyl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-acrylamide (substance 101)

Production analogous to Example 1.

Batch size: 19.2 g (129 mmol) 3-(3-pyridyl)-acrylic acid, 15.1 ml (176 mmol) oxalyl chloride and 37.6 g (117 mmol) 4-(9-fluorenyl)-piperazin-1-yl-butylamine.

In the recovery, 150 ml 10% sodium hydroxide solution is added to the reaction solution. The organic phase is washed three times each with 60 ml water, dried over sodium sulfate and the solvent is removed under vacuum. The residue is chromatographically purified over silica gel with  $\text{CHCl}_3/\text{CH}_3\text{OH}$  (90/10). After removal of the solvent, this is crystallized with 1700 ml acetic acid ethyl ester: solid with MP. 145-147°C; yield 39.0 g (73%).

$\text{C}_{29}\text{H}_{32}\text{N}_4\text{O}$  (452.6)

IR-Spectrum (KBr):

n(NH)	3300 $\text{cm}^{-1}$
n(C=O)	1655, 1530 $\text{cm}^{-1}$
n(C=C)	1620 $\text{cm}^{-1}$

$^1\text{H}$ -NMR-Spectrum ( $\text{CDCl}_3$ ):

1.35 - 1.80	(4H, m, C-CH <sub>2</sub> -CH <sub>2</sub> -C)
2.10 - 2.80	(10H, m, piperazine, N-CH <sub>2</sub> )
3.20 - 3.50	(2H, m, CONHCH <sub>2</sub> )
4.81	(1H, s, Ar <sub>2</sub> CH)
6.34	(1H, d, CH=CHCO, J=15.7 Hz)
6.75 - 7.05	(1H, m, NH)
7.00 - 7.80	(11H, m, Ar, CH=CHCO, Py)
8.40 - 8.60	(1H, m, Py)



8.60 - 8.70 (1H, m, Py)

## Example 14

N-{4-[4-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-yl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-acrylamide (substance 119)

Production analogous to Example 1.

Batch size: 4.16 g (27.9 mmol) 3-(3-pyridyl)-acrylic acid, 6.5 ml (76.2 mmol) oxalyl chloride and 12.2 g (25.4 mmol) 4-[4-(10, 11-dihydro-5H-dibenzo[a,d]cyclohepten-5-yl)-piperazin-1-yl]-butylamine.

In the recovery, 50 ml 10% sodium hydroxide solution is added to the reaction solution. The aqueous phase is extracted with dichloromethane. The combined organic phases are washed with 50 ml water, dried over sodium sulfate and the solvent is removed under vacuum. The residue is chromatographically pre-purified over silica gel with  $\text{CHCl}_3/\text{CH}_3\text{OH}$  (98/2 to 90/10). After removal of the solvent, this is crystallized four-fold each with 50 ml acetic acid ethyl ester: Colorless solid with MP. 119-120°C; yield 4.9 g (40%).

$\text{C}_{31}\text{H}_{36}\text{N}_4\text{O}$  (480.7)

IR-Spectrum (KBr):

n(NH)	3280 $\text{cm}^{-1}$
n(C=O)	1670, 1540 $\text{cm}^{-1}$
n(C=C)	1625 $\text{cm}^{-1}$

$^1\text{H}$ -NMR-Spectrum ( $\text{CDCl}_3$ ):

1.40 - 1.80	(4H, m, $\text{C}-(\text{CH}_2)_2\text{-C}$ )
2.10 - 2.55	(10H, m, piperazine, $\text{N-CH}_2$ )
2.55 - 3.00	(2H, m, $\text{Ar-CH}_2\text{-CH}_2\text{-Ar}$ )
3.20 - 3.50	(2H, m, $\text{CONHCH}_2$ )
4.75 - 4.20	(3H, m, $\text{Ar}_2\text{CH}$ , $\text{Ar-CH}_2\text{-CH}_2\text{-Ar}$ )
6.50	(1H, d, $\text{CH=CHCO}$ , $J=15.7$ Hz)
6.90 - 7.40	(10H, m, Ar, Py, NH)

7.61 (1H, d,  $\text{CH}=\text{CHCO}$ ,  $J=15.7$  Hz)  
 7.70 - 7.85 (1H, m, Py)  
 8.50 - 8.65 (1H, m, Py)  
 8.70 - 8.80 (1H, m, Py)

### Example 15

N-[2-(4-diphenylacetyl-piperazin-1-yl)-ethyl]-3-pyridin-3-yl-acrylamide (substance 158)

8.0 g (22.6 mmol) N-[2-(piperazin-1-yl)-ethyl]-3-pyridin-3-yl-acrylic amide trihydrochloride (substance 317 as a trihydrochloride) and 13 ml (92.7 mmol) TEA are present in 100 ml absolute dichloromethane and cooled to ca. 0°C under moisture exclusion. 6.3 g (24.9 mmol) diphenylacetyl chloride (90%) are dissolved in 70 ml absolute dichloromethane and added dropwise. The mixture is stirred overnight at RT without further cooling. Subsequently, 200 ml dichloromethane are added and the batch is washed three times each with 100 ml water. The organic phase is dried over sodium sulfate and the solvent is removed under vacuum. The residue is crystallized twice from 40 ml and 30 ml acetonitrile. Beige coloured crystals of MP. 174°C. yield 4.6 g (44%).

$\text{C}_{26}\text{H}_{30}\text{N}_4\text{O}_2$  (454.6)

IR-Spectrum (KBr):

n(NH)	3320 $\text{cm}^{-1}$
n(C=O)	1675, 1550 $\text{cm}^{-1}$
n(C=C)	1610 $\text{cm}^{-1}$

$^1\text{H}$ -NMR-Spectrum ( $\text{CDCl}_3$ ):

$\text{CH}_2$ )	2.10 - 2.60 (6H, m, piperazine, N-
	3.30 - 3.85 (6H, m, piperazine,
	CONHCH <sub>2</sub> )
	5.20 (1H, s, $\text{Ar}_2\text{CH}$ )
	6.20 - 6.40 (1H, m, NH)

6.46 (1H, d, CH=CHCO, J=15.7 Hz)  
 7.10 - 7.40 (11H, m, Ar, Py)  
 7.61 (1H, d, CH=CHCO, J=15.7 Hz)  
 7.70 - 7.90 (1H, m, Py)  
 8.50 - 8.65 (1H, m, Py)  
 8.70 - 8.80 (1H, m, Py)

#### Example 16

N-{2-[4-(10,11-dihydro-dibenzo[b,f]azepin-5-yl-carbonyl)-piperazin-1-yl]-ethyl}-3-pyridin-3-yl-propionamide (substance 215)

Production analogous to Example 15.

Batch size: 8.0 g (21.5 mmol) N-[2-(piperazin-1-yl)-ethyl]-3-pyridin-3-yl-propionamide-trihydrochloride, 12.3 ml (88.1 mmol) TEA and 6.1 g (23.6 mmol) 10,11-dihydro-dibenzo[b,f]azepin-5-carbonyl chloride in 170 ml absolute dichloromethane.

In the purification, this is first chromatographically pre-purified over silica gel with CHCl<sub>3</sub>/CH<sub>3</sub>OH (100/0 to 90/10) and, after removal of the solvent, crystallized from 10 ml acetonitrile. Colorless crystals of MP. 146 - 147°C. yield 0.7 g (6%).

C<sub>29</sub>H<sub>33</sub>N<sub>5</sub>O<sub>2</sub> (483.6)

IR-Spectrum (KBr):

n(NH)	3330 cm <sup>-1</sup>
n(C=O)	1660, 1535 cm <sup>-1</sup>
n(C=C)	1630 cm <sup>-1</sup>

<sup>1</sup>H-NMR-Spectrum (CDCl<sub>3</sub>):

2.10 - 2.60	(8H, m, piperazine, CO-CH <sub>2</sub> , N-CH <sub>2</sub> )
2.96	(2H, t, Py-CH <sub>2</sub> , J=7.4 Hz)
3.10 - 3.45	(10H, m, CONHCH <sub>2</sub> , piperazine, Ar-CH <sub>2</sub> -CH <sub>2</sub> -Ar)

5.80 - 6.00 (1H, m, NH)  
 7.00 - 7.60 (10H, m, Ar, Py)  
 8.35 - 8.55 (2H, m, Py)

#### Example 17

N-{2-[4-(naphthalin-2-yl-sulfonyl)-piperazin-1-yl]-ethyl}-3-pyridin-3-yl-acrylamide (substance 227)

Production analogous to Example 15.

Batch size: 8.0 g (22.6 mmol) N-[2-(piperazin-1-yl)-ethyl]-3-pyridin-3-yl-acrylamide trihydrochloride (substance 317 as a trihydrochloride), 13 ml (92.7 mmol) TEA and 5.6 g (24.9 mmol) naphthalin-2-sulfonic acid chloride in 180 ml absolute dichloromethane.

For purification, this is crystallized twice from 150 ml and 100 ml acetonitrile. Beige coloured crystals of MP. 183 - 184°C. yield 4.0 g (39%).

$C_{24}H_{26}N_4O_3S$  (450.6)

IR-Spectrum (Kbr):

n(NH)	3250 $cm^{-1}$
n(C=O)	1665, 1555 $cm^{-1}$
n(C=C)	1625 $cm^{-1}$

$^1H$ -NMR-Spectrum ( $CDCl_3$ ):

2.35 - 2.80 (6H, m, piperazine, N-CH <sub>2</sub> )
3.00 - 3.35 (4H, m, piperazine)
3.44 (2H, dd, CONHCH <sub>2</sub> , J=5.5 Hz, J=11.2 Hz)
5.90 - 6.15 (1H, m, NH)
6.35 (1H, d, CH=CHCO, J=15.6 Hz)
7.15 - 8.15 (9H, m, Ar, CH=CHCO, Py)
8.35 (1H, bs, Ar)
8.45 - 8.60 (1H, m, Py)
8.60 - 8.75 (1H, m, Py)

## Example 18

N-{2-[4-(tert-butoxycarbonyl)-piperazin-1-yl]-ethyl}-3-pyridin-3-yl-acrylamide (substance 374)

Production analogous to Example 1.

Batch size: 36.1 g (242 mmol) 3-(3-pyridyl)-acrylic acid, 23.1 ml (264 mmol) oxalyl chloride, 50 g (15.4 mmol) 2-[4-(tert-but-oxycarbonyl)-piperazin-1-yl]-ethylamine and 30.4 ml (220 mmol) TEA in 400 ml absolute dichloromethane.

In the purification, this is crystallized twice from 100 ml acetic acid ethyl ester and 150 ml 1-chlorbutane: Colorless crystals of MP. 92 - 93°C; yield 38.4 g (48%).

$C_{19}H_{28}N_4O_3$  (360.5)

IR-Spectrum (Kbr):

n(NH)	3320 $cm^{-1}$
n(C=O)	1670, 1530 $cm^{-1}$
n(C=C)	1620 $cm^{-1}$

$^1H$ -NMR-Spectrum ( $CDCl_3$ ):

1.47 (9H, s, tert. Butyl )
2.30 - 2.80 (6H, m, piperazine, N-CH <sub>2</sub> )
3.30 - 3.70 (6H, m, piperazine CONHCH <sub>2</sub> )
6.30 - 6.55 (1H, m, NH)
6.52 (1H, d, CH=CHCO, J=15.7 Hz)
7.20 - 7.40 (1H, m, Py)
7.63 (1H, d, CH=CHCO, J=15.7 Hz)
7.70 - 7.90 (1H, m, Py)
8.45 - 8.65 (1H, m, Py)
8.65 - 8.85 (1H, m, Py)

## Example 19

N-[2-(piperazin-1-yl)-ethyl]-3-pyridin-3-yl-acrylamide•tri-hydrochloride (substance 317 as a trihydrochloride)

38 g (105 mmol) N-{2-[4-(tert-butoxycarbonyl)-piperazin-1-yl]-ethyl}-3-pyridin-3-yl-acrylamide (substance 374) are dissolved in 380 ml methanol and 42 ml concentrated hydrochloric acid are added. The mixture is stirred for four hours under reflux. After cooling of the solution, the solvent is removed under vacuum. The residue is crystallized from 185 ml methanol. Colorless crystals of MP. 216 - 226°C (under decomposition). yield 34.8 g (93%).

$C_{14}H_{20}N_4O \cdot 3HCl$  (369,7)

IR-Spectrum (KBr):

n(NH)	3150 $cm^{-1}$
n(C=O)	1670, 1540 $cm^{-1}$
n(C=C)	1610 $cm^{-1}$


$^1H$ -NMR-Spectrum ( $D_2O$ ):

3.20 - 3.75	(12H, m, piperazine, N-CH <sub>2</sub> -CH <sub>2</sub> )
6.74	(1H, d, CH=CHCO, J=15.9 Hz)
7.44	(1H, d, CH=CHCO, J=15.9 Hz)
7.80 - 8.00	(1H, m, Py)
8.50 - 8.70	(2H, m, Py)
8.80 - 8.90	(1H, m, Py)









In the following Table 2, further synthesized end products according to formula (I) are listed:

Table 2:















synthesized compounds  
of formula (I)



Nr	$R^1 - R^3$	A	D-E-G	mp. [°C] (solvent) <sup>1</sup>
1	H	$CH_2CH_2$	$CH_2CH_2CH_2CH_2-N$ (piperidine ring)	oil <sup>1</sup>
2	H	$CH=CH$	$CH_2CH_2CH_2CH_2-N$ (piperidine ring)	240-242 <sup>m</sup> (EtOH)
25	H	$CH_2NHCH_2CH_2$	$CH_2CH_2-N$ (piperidine ring) $-CH(C_6H_5)_2$	amorph <sup>2,v</sup> ( $CHCl_3/MeOH/$ $NH_3$ 90/9/1)
27	H	$OCH_2$	$CH_2CH_2CH_2CH_2-N$ (piperidine ring) $-CH(C_6H_5)_2$	91-93 (PE)
29	H	$CH=CH$	$CH_2CH_2CH(OH)CH_2-N$ (piperidine ring) $-CH(C_6H_5)_2$	168-171 (EE)
30	H	$CH=CH$	$OCH_2CH_2CH_2-N$ (piperidine ring) $-CH(C_6H_5)_2$	105-107 (EE)
31	H	$CH=CH$	$CH_2CH_2CH_2C(=O)-N$ (piperidine ring) $-CH(C_6H_5)_2$	179-180 (MEK)
32	H	$CH=CH$	$CH_2CH_2CH_2SO_2-N$ (piperidine ring) $-CH(C_6H_5)_2$	167-168 (EE)
33	H	$CH=CH$	$CH_2CH_2NH-C(=O)-N$ (piperidine ring) $-CH(C_6H_5)_2$	187-188 (iPrOH)
35	H	$CH=CH$	$CH_2CH_2OCH_2CH_2-N$ (piperidine ring) $-CH(C_6H_5)_2$	157-159 <sup>3</sup> (MeOH/iPr <sub>2</sub> O)

Nr	R <sup>1</sup> - R <sup>3</sup>	A	D-E-G	mp. [°C] (solvent) <sup>i</sup>
36	H	CH=CH	$\text{CH}_2\text{CH}_2\text{CH}_2\text{C}\equiv\text{CCH}_2-\text{N} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \text{N}-\text{CH}(\text{C}_6\text{H}_5)_2$	162-164° (90% iPrOH)
37	H	CH=CH	$\text{CH}_2\text{C}\equiv\text{C}-\text{CH}=\text{CHCH}_2-\text{N} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \text{N}-\text{CH}(\text{C}_6\text{H}_5)_2$	amorph? (CHCl <sub>3</sub> /MeOH)
38	H	CH=CH	$\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{N} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \text{N}-\text{CH}(\text{C}_6\text{H}_5)_2$ 	141-145° (EtOH)
42	H	CH=CH	$\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{N} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \text{N}-\text{CH}(\text{C}_6\text{H}_5)_2$ 	125-127 (MeCN)
47	H	CH=CH	$\text{CH}_2\text{C}\equiv\text{CCH}_2-\text{N} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \text{N}-\text{CH}(\text{C}_6\text{H}_4\text{F})_2$ 	160-163° (EtOH/iPr <sub>2</sub> O)
60	H	OCH <sub>2</sub>	$\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{N} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \text{N}-\text{CH}(\text{C}_6\text{H}_5)_2$ 	resin? (CHCl <sub>3</sub> /MeOH/ NH <sub>3</sub> 90/9/1)
68	H	CH=CH	$\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{N} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \text{N}-\text{CH}(\text{C}_6\text{H}_5)_2$ 	115 (BuCl)
75	H	$\text{CH}=\text{C} \begin{array}{c} \diagup \\ \text{CH}_3 \end{array}$	$\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{N} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \text{N}-\text{CH}(\text{C}_6\text{H}_5)_2$ 	186-188° (EtOH)
83	H	CH=CH	$\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{N} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \text{N}-\text{CH}(\text{C}_6\text{H}_5)_2$ 	178-180 (iPrOH)
85	H	CH <sub>2</sub> CH <sub>2</sub>	$\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{N} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \text{N}-\text{CH}(\text{C}_6\text{H}_5)_2$ 	93 (BuCl)



Nr	R <sup>1</sup> - R <sup>3</sup>	A	D-E-G	mp.[°C] (solvent) <sup>i</sup>
87	H	CH=CH	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N  N 	110-114 (BuCl)
93	H	CH <sub>2</sub> CH <sub>2</sub>	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N  N 	81-82 (MeCN)
94	H	CH=CH	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N  N 	108-109 (MeCN)
96	H	CH <sub>2</sub> CH <sub>2</sub>	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N  N 	104-105 (EE)
97	H	CH=CH	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N  N 	124-125 (EE/Pr <sub>2</sub> O)
101	H	CH=CH	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N  N 	145-147 (EE)
119	H	CH=CH	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -N  N 	119-120 (EE)

Nr	R <sup>1</sup> - R <sup>3</sup>	A	D-E-G	mp.[°C] (solvent) <sup>1</sup>
138	H	CH=CH		amorph <sup>2</sup> (CHCl <sub>3</sub> /MeOH)
158	H	CH=CH		174 (MeCN)
208	H	CH=CH		150 (acetone)
215	H	CH <sub>2</sub> CH <sub>2</sub>		146-147 (MeCN)
227	H	CH=CH		183-184 (MeCN)
231	H	CH=CH		amorph <sup>2</sup> (CHCl <sub>3</sub> /MeOH)
234	H	CH=CH		amorph <sup>2</sup>
251	H	CH=CH		Oil <sup>2</sup>

Nr	R <sup>1</sup> - R <sup>3</sup>	A	D-E-G	mp.[°C] (solvent) <sup>i</sup>
305	H	CH=CH		Oil <sup>2</sup>
317	H	CH=CH		216-226/Zers. <sup>3</sup> (MeOH)
349	H	CH=CH		amorph (MeCN)
374	H	CH=CH		92-93 (EE/BuCl)

- 1
- PE = Petroleum ether  
 EE = Ethyl acetate  
 MEK = Methyl ethyl ketone  
 iPrOH = Isopropanol  
 iPr<sub>2</sub>O = Diisopropyl ether  
 MeOH = Methanol  
 EtOH = Ethanol

MeCN = Acetonitrile  
 BuCl = 1-Chlorobutane

- 2 Purified by column chromatography  
 3 as a Trihydrochloride  
 4 as a Tetrahydrochloride  
 5 as a Trioxalate  
 6 as a Sulfate

## Production of the Starting Substances

## Example A

## 4-(4-diphenylphosphinoyl-piperazin-1-yl)-butylamine

## a) N-diphenylphosphinoyl-piperazine:

20 g (84.5 mmol) diphenylphosphinate chloride dissolved in 30 ml DMF are added dropwise to a solution of 21.8 g (253 mmol) piperazine in DMF. After four hours, the solution is concentrated under vacuum, taken up in chloroform and extracted by shaking with 10% hydroxide solution. The organic phase is dried over sodium sulfate and solvent is removed under vacuum. The residue is chromatographically purified over silica gel with  $\text{CHCl}_3/\text{CH}_3\text{OH}/\text{TEA}$  (90/10/0 to 90/10/6): yield 12 g (49%).

b) 2-[4-(4-diphenylphosphinoyl)-butyl]-isoindolin-1,3-dione: 12 g (41.9 mmol) N-diphenylphosphinoyl-piperazine, 11.8 g (42 mmol) N-(4-bromobutyl)-phthalimide, 5.8 g (42 mmol) potassium carbonate and 1.4 g (8 mmol) potassium iodide are heated in ethyl methyl ketone for 6 hours under reflux. After cooling, the reaction mixture is concentrated under vacuum. The residue is taken up in acetic acid ethyl ester and extracted by shaking with water. The organic phase is dried over sodium sulfate and the solvent is removed under vacuum: yield 20 g (100%).

c) 4-(4-diphenylphosphinoyl-piperazin-1-yl)-butylamine: 20 g (40 mmol) 2-[4-(4-diphenylphosphinoyl)-butyl]-isoindolin-1,3-dione and 4 ml (80 mmol) hydrazine hydrate are heated in 400 ml ethanol for three hours under reflux. The cooling solution is concentrated under vacuum and the residue is taken up in acetic acid ethyl ester. The suspension is filtered and the residue is washed with toluene. The filtrate and washing fluid are concentrated under vacuum until dry. Subsequently, the residue is taken up in chloroform and shaken with 10% sodium sulfate. The organic phase is dried over sodium sulfate and the solvent is removed under vacuum. The accumulated crude product is further processed without further purification: yield 13.3 g (95%).

## Example B

## 4-(4-diphenylmethyl-piperazin-1-yl)-3-hydroxy-butylamine

## a) 2-(but-3-enyl)-isoindolin-1,3-dione:

50 g (370 mmol) 4-bromo-1-butene and 68.5 g (370 mmol) phthalimide potassium salt are suspended in 800 ml ethyl methyl ketone and heated under reflux for 14 hours. After cooling, the mixture is filtrated and the filtrate is concentrated under vacuum. The residue is taken up in acetic acid ethyl ester and washed with water. The organic phase is dried over sodium sulfate and the solvent is removed under vacuum: yield 50 g (67%).

## b) 2-(3,4-epoxybutyl)-isoindolin-1,3-dione:

70 g (350 mmol) 2-(but-3-enyl)-isoindolin-1,3-dione are dissolved in dichloromethane. The solution is cooled to ca. 0°C and a suspension of 120.8 g (350 mmol) 50% 3-chloroperoxy-benzoic acid in dichloromethane is added under cooling. The mixture is left standing without further cooling at room temperature for two days. After addition of 250 ml saturated  $\text{NaHCO}_3$  solution the organic phase is separated and washed three times each with 200 ml saturated  $\text{NaHCO}_3$  solution and once with water. The organic phase is dried over sodium sulfate and the solvent is removed under vacuum: yield 80 g.

## c) 2-[4-(4-diphenylmethyl-piperazin-1-yl)-3-hydroxy-butyl]-isoindolin-1,3-dione:

5 g (~25 mmol) 2-(3,4-epoxybutyl)-isoindolin-1,3-dione, 7.5 g (30 mmol) benzhydrylpiperazine and 3.5 g (25 mmol) potassium carbonate are stirred in DMF for 6 hours 80°C. After cooling, the reaction mixture is filtered and concentrated under vacuum. The residue is taken up in 300 ml acetic acid ethyl ester and washed three times each with 20 ml water. The organic phase is dried over sodium sulfate and the solvent is removed under vacuum. The residue is chromatographically purified over silica gel with  $\text{CHCl}_3/\text{CH}_3\text{OH}$  (100/10).

### Example C

a) 1-(4-chlorobutyl)-4-diphenylmethyl-piperazine:  
25 g (99 mmol) benzhydrylpiperazine and 15.2 ml (109 mmol) TEA are present in 200 ml absolute THF and cooled to ca. 0°C under moisture exclusion. 14 g (99 mmol) 4-chlorobutyric chloride are dissolved in 40 ml absolute THF and added dropwise. The mixture is stirred an additional three hours at RT and subsequently filtered. The filtrate is concentrated under vacuum, the residue is taken up in acetic acid ethyl ester and washed with saturated NaCl solution. The organic phase is dried over sodium sulfate and concentrated under vacuum until dry. The resin is further processed without further purification: yield 35.1 g (99%).

b) 1-(4-aminobutyl)-4-diphenylmethyl-piperazine:  
8.9 g (24.9 mmol) 1-(4-chlorobutyl)-4-diphenylmethylpiperazine, 4.8 g (73.8 mmol) sodium azide, 1 g potassium iodide and 1 g molecular sieve 4A are stirred in 70 ml DMF for five hours at 70°C. After cooling, the reaction mixture is filtered and the filtrate is concentrated under vacuum. The accumulated crude product is dissolved in methanol and mixed with a spatula tip of palladium-carbon (10%). The mixture is stirred for two days at RT under hydrogen atmosphere. The

mixture is filtered from the catalyst and the solvent is removed under vacuum. The residue is chromatographically purified over silica gel with  $\text{CHCl}_3/\text{CH}_3\text{OH}/\text{TEA}$  (90/10/5): yield 5.2 g (62%) as a colorless amorphous solid.

#### Example D

##### 3-(4-diphenylmethyl-piperazin-1-yl-sulfonyl)-propylamine

a) 1-diphenylmethyl-4-(3-chloropropanesulfonyl)-piperazine: 39.2 g (155 mmol) benzhydrylpiperazine and 19.7 ml (141 mmol) TEA are present in 100 ml absolute dichloromethane and cooled to ca. 0°C under moisture exclusion. 25 g (141 mmol) 3-chloropropanesulfonyl chloride are dissolved in 70 ml absolute dichloromethane and added dropwise. The mixture is stirred for two hours under cooling and subsequently mixed with chloroform and washed with saturated NaCl solution. The organic phase is dried over sodium sulfate and concentrated under vacuum until dry. The solid is further processed without further purification: yield 59.2 g.

b) 2-[3-(4-diphenylmethyl-piperazin-1-yl-sulfonyl)-propyl]-isoindolin-1,3-dione:

The reaction of the halogenide to phthalimide occurs analogously to Example B)a).

Batch size: 15 g (38 mmol) 1-diphenylmethyl-4-(3-chloropropane sulfonyl)-piperazine and 7.2 g (39 mmol) phthalimide-potassium salt in DMF at 80°C.

The purification occurs by chromatography on silica gel with petroleum ether/acetic acid ethyl ester (4/1). Yield 14 g (71%).

c) 3-(4-diphenylmethyl-piperazin-1-yl-sulfonyl)-propylamine:

The reaction of the phthalimide to the amine occurs analogously to Example B)d).

Batch size: 14g (27.8 mmol) 2-[3-(4-diphenylmethyl-piperazin-1-yl-sulfonyl)-propyl]-isoindolin-1,3-dione and 2.8 ml (55.6 mmol) hydrazine-hydrate.

The resulting resin is further processed without further purification: Yield 1.5 g (15%).

## Example E

## 2-[2-(4-diphenylmethyl-piperazin-1-yl)-ethoxy]-ethylamine

a) 2-[2-(4-diphenylmethyl-piperazin-1-yl)-ethoxy]-ethanol: 71.7 g (284 mmol) benzhydrylpiperazine, 45 g (361 mmol) 2-(2-chloroethoxy)-ethanol, 43.2 g (312 mmol) potassium carbonate and 9.4 g (57 mmol) potassium iodide are stirred in 400 ml absolute DMF for 8 hours at 75°C. After cooling, the solution is concentrated under vacuum. The residue is distributed between acetic acid ethyl ester and water. The aqueous phase is extracted twice with acetic acid ethyl ester and the combined organic phases are washed three times with saturated sodium chloride solution. The organic phase is dried over sodium sulfate and the solvent is removed under vacuum. The residue is chromatographically purified over silica gel with  $\text{CHCl}_3/\text{CH}_3\text{OH}$  (90/10): Yield 104g.

b) 2-{2-[2-(4-diphenylmethyl-piperazin-1-yl)-ethoxy]-ethyl}-isoindolin-1,3-dione:  
40 g (~118 mmol) 2-[2-(4-diphenylmethyl-piperazin-1-yl)-ethoxy]-ethanol, 31.1 g (119 mmol) triphenylphosphine and 17.3 g (118 mmol) phthalimide are suspended in 200 ml THF and 24.2 ml (119 mmol) azodicarbonic acid diethyl ester are added dropwise under protective atmosphere and light cooling (to ca. 15°C). The mixture is stirred without further cooling for three hours and subsequently, the solvent is removed under vacuum. The residue is taken up in 1N HCl and washed twice each with 50 ml acetic acid ethyl ester respectively. The aqueous phase is neutralized with ca. 50 g sodium hydrogen carbonate and extracted four times each with 125 ml chloroform. The combined chloroform phases are dried over sodium sulfate and the solvent is removed under vacuum. The residue is chromatographically purified over silica gel with  $\text{CHCl}_3/\text{CH}_3\text{OH}$  (99/1 to 90/10): Yield 11 g (19%).

c) 2-[2-(4-diphenylmethyl-piperazin-1-yl)-ethoxy]-ethylamine:  
The reaction of the phthalimide to the amine occurs analogously to Example B)d).



Batch size: 27g (55.6 mmol) 2-{2-[2-(4-diphenylmethyl-piperazinyl)-ethoxy]-ethyl}-isoindol-1,3-dione and 5.4 ml (110 mmol) hydrazine hydrate.

The purification occurs by chromatography on silica gel with  $\text{CHCl}_3/\text{CH}_3\text{OH}/\text{TEA}$  (9/1/0 to 9/1/1): Yield 12.4 g (66%).

#### Example F

4-{4-[bis-(4-fluorophenyl)-methyl]-piperazin-1-yl}-but-2-in-ylamine

a) 2-propinyl-isoindolin-1,3-dione:

32.3 g (271 mmol) 3-bromopropine are dissolved in 150 ml DMF and 50.3 g (271 mmol) phthalimide potassium salt are added under ice cooling. The suspension is warmed at 70°C for eight hours. The mixture is concentrated under a vacuum and the residue is distributed between acetic acid ethyl ester and water. The organic phase is dried over sodium sulfate and the solvent is removed under vacuum. The residue is crystallized from acetic acid ethyl ester: Yield 36.4 g (72%) colorless crystals.

b) 2-(4-{4-[bis-(4-fluorophenyl)-methyl]-piperazin-1-yl}-but-2-inyl)-isoindolin-1,3-dione:

15 g (81 mmol) 2-propinyl-isoindolin-1,3-dione, 15 g (52 mmol) [bis-(4-fluorophenyl)-methyl]-piperazine, 2.5 g (81 mmol) paraformaldehyde and 0.2 g copper sulfate are stirred for three hours in 200 ml dioxane at 100°C. The cool solution is concentrated under vacuum and the residue is distributed between acetic acid ethyl ester and saturated NaCl solution. The organic phase is dried over sodium sulfate and concentrated under vacuum until dry. The residue is chromatographically purified over silica gel with acetic acid ethyl ester/petroleum ether (1/1): Yield 23.4 g (93%) yellow amorphous solid.

c) 4-{4-[bis-(4-fluorophenyl)-methyl]-piperazin-1-yl}-but-2-inylamine:

00595001.051500

The reaction of the phthalimide to the amine occurs analogously to Example 21)d): 23.3 g (48 mmol) 2-(4-{4-[bis-(4-fluorophenyl)-methyl]-piperazin-1-yl}-but-2-ynyl)-isoindolin-1,3-dione and 4.7 ml (96 mmol) hydrazine hydrate. The purification occurs by chromatography on silica gel with  $\text{CHCl}_3/\text{CH}_3\text{OH}$  (90/10 to 85/15): Yield 9.8 g (58%).

#### Example G

4-[4-(1,2,3,4-tetrahydronaphthalin-1-yl)-piperazin-1-yl]-butylamine

a) 2-(4-(4-(1,2,3,4-tetrahydronaphthalin-1-yl)-piperazin-1-yl)-butyl)-isoindolin-1,3-dione dihydrochloride: 30 g (138,6 mmol) 1-(1,2,3,4-tetrahydronaphthyl)-1-piperazine, 40.3 g (140 mmol) N-(4-bromobutyl)-phthalimide and 27.6 g (200 mmol) potassium carbonate are stirred in 150 ml DMF for three hours at RT. The mixture is concentrated under a vacuum and the residue is distributed between 200 ml acetic acid ethyl ester and 150 ml water. The aqueous phase is extracted 50 ml acetic acid ethyl ester and the combined organic phases are washed four times with water. The organic phase is dried over sodium sulfate and concentrated under vacuum until dry. The residue is dissolved in 400 ml methanol and mixed with 70 ml 6.6M isopropanolic hydrochloric acid. The salt precipitated in the cold is drawn off and dried. Colorless crystals of MP.175-180°C: Yield 53.2 g (78%).

b) 4-[4-(1,2,3,4-tetrahydronaphthalin-1-yl)-piperazin-1-yl]-butylamine:  
The reaction of the phthalimide to the amine occurs analogously to Example B)d).  
Batch size: 52 g (106 mmol) 2-(4-[4-(1,2,3,4-tetrahydronaphthalin-1-yl)-piperazin-1-yl]-butyl)-isoindolin-1,3-dione dihydrochloride and 14.6 ml (300 mmol) hydrazine hydrate in 500 ml ethanol.

The accumulated crude product is further processed without further purification: Yield 27.4 g (89%).

## Example H

4-[4-(5,6,7,8-tetrahydro-naphthalin-1-yl)-piperazin-1-yl]-butylamine•trihydrochloride

a) 2-{4-[4-(5,6,7,8-tetrahydro-naphthalin-1-yl)-piperazin-1-yl]-butyl}-isoindolin-1,3-dione:

The reaction of the piperazine with the phthalimide occurs analogously to Example 26a.

Batch size: 24 g (110.9 mmol) 1-(5,6,7,8,-tetrahydronaphthalin-1-yl)-piperazine, 32.6 g (115.4 mmol) N-(4-bromobutyl)-phthalimide and 30.6 g (221.8 mmol) potassium carbonate in 240 ml DMF.

The purification occurs by chromatography over silica gel with  $\text{CHCl}_3/\text{CH}_3\text{OH}$  (100/0 to 98/2): Yield 41.6 g (89%).

b) 4-[4-(5,6,7,8-tetrahydronaphthalin-1-yl)-piperazin-1-yl]-butylamine.Trihydrochloride:

The reaction of the phthalimide to the amine occurs analogously to Example 21d.

Batch size: 41.5 g (99.4 mmol) 2-{4-[4-(5,6,7,8-tetrahydronaphthalin-1-yl)-piperazin-1-yl]-butyl}-isoindolin-1,3-dione and 9.5 ml (198.8 mmol) hydrazine hydrate in 400 ml ethanol.

The purification occurs by a chromatography over silica gel with  $\text{CHCl}_3/\text{CH}_3\text{OH}/\text{NH}_4\text{OH}$  (90/10/0 to 90/9/1). After removing the solvent the residue is dissolved in 300 ml isopropanol and mixed with 47 ml 6M isopropanolic hydrochloric acid. The salts precipitating in the cold is filtered off and dried: Yield 23.5 g (59%).

## Example I

4-[4-(naphthalin-1-yl)-piperazin-1-yl]-butylamine

a) 2-[4-(naphthalin-1-yl)-piperazin-1-yl]-butyl]-isoindolin-1,3-dione:

The reaction of the piperazine with the phthalimide occurs analogously to Example G)a).

Batch size: 21 g (100 mmol) 1-(1-naphthyl)-piperazine  
 [Production according to Glennon et al., J. Med. Chem. 29, 2375 (1986)], 28.3 g (100 mmol) N-(4-bromobutyl)-phthalimide and 20.8 g (150 mmol) potassium carbonate in 250 ml DMF.  
 The accumulated crude product is further processed without further purification: Yield 30 g (68%).

b) 4-[4-(naphthalin-1-yl)-piperazin-1-yl]-butylamine:  
 The reaction of phthalimide to the amine occurs analogously to Example B)d).  
 Batch size: 30 g (68 mmol) 2-[4-(naphthalin-1-yl)-piperazin-1-yl]-butyl]-isoindolin-1,3-dione and 6.9g (137 mmol) hydrazine hydrate in ethanol.

The accumulated crude product is further processed without further purification: Yield 14.6 g (75%).

#### Example J

##### 11-methanesulfonyloxy-6,11-dihydrodibenzo[b,e]-thiepine

a) 6.11-dihydro-dibenzo[b,e]thiepin-11-ol:  
 48 g (212 mmol) 6.11-dihydro-dibenzo[b,e]thiepin-11-one are dissolved in 480 ml absolute methanol and cooled to ca. -10°C. 19.2 g (507 mmol) sodium borohydride is added portion wise to this solution. The mixture is stirred for three hours at RT without further cooling. After the careful addition of 30 ml water, the suspension is concentrated under vacuum. The residue is taken up in 500 ml dichloromethane and washed twice each with 150 ml water. The organic phase is dried over sodium sulfate and the solvent is removed under vacuum. The residue is crystallized from 180 ml toluene. Colorless crystals of MP. 108°C: Yield 41.2 g (85%).

b) 11-methanesulfonyloxy-6,11-dihydrodibenzo[b,e]-thiepine:  
 3.5 g (15.5 mmol) 6,11-dihydro-dibenzo[b,e]thiepin-11-ol and 2.4 ml (17 mmol) TEA are dissolved in 50 ml absolute dichloromethane. The mixture is cooled to ca. 5°C and a solution of 1.3 ml (16.3 mmol) methanesulfonyl chloride in 10 ml absolute dichloromethane is added dropwise. The mixture

is additionally stirred for two at RT and directly employed in Example 11.

#### Example K

##### 2-(4-tert-butoxycarbonyl-piperazin-1-yl)-ethylamine

a) 2-{2-[4-(tert-butoxycarbonyl)-piperazin-1-yl]-ethyl}-isoindolin-1,3-dione:  
44.7 g (240 mmol) N-(tert-butoxycarbonyl)-piperazine, 60.9 g (240 mmol) N-(2-bromoethyl)-phthalimide, 49.8 g (360 mmol) potassium carbonate and 49.5 g (330 mmol) sodium iodide are heated in 1000 ml ethyl methyl ketone for five hours under reflux. After cooling the reaction mixture and concentrated under vacuum. The residue is taken up in 700 ml chloroform and extracted twice by shaking each with 50 ml water. The organic phase is dried over sodium sulfate and the solvent is distilled off. The residue is crystallized from 110 ml methanol. Colorless crystals of MP. 136 - 138°C: Yield 47.6 g (55%).

b) 2-[4-(tert-butoxycarbonyl)-piperazin-1-yl]-ethylamine:  
The reaction of the phthalimide to the amine occurs analogously to Example B)d).  
Batch size: 42.2 g (120 mmol) 2-{2-[4-(tert-butoxycarbonyl)-piperazin-1-yl]-ethyl}-isoindolin-1,3-dione and 11.6 ml (240 mmol) hydrazine hydrate in 450 ml ethanol.

The accumulated crude product is further processed without further purification: Yield 24.8 g (90%).

#### Example L

##### 4-(4-biphenyl-2-yl-piperazin-1-yl)-butylamine

a) 2-[4-(4-biphenyl-2-yl-piperazin-1-yl)-butyl]-isoindolin-1,3-dione:  
The reaction of piperazine with the phthalimide occurs analogously to Example G)a).

Batch size: 15.3 g (64.2 mmol) 1-(o-biphenyl)-piperazine, 18.5 g (64.2 mmol) N-(4-bromobutyl)-phthalimide and 13.3 g (96 mmol) potassium carbonate in 270 ml ethyl methyl ketone. The purification occurs by chromatography over silica gel with  $\text{CHCl}_3/\text{CH}_3\text{OH}$  (98/2): Yield 29 g (99%).

b) 4-(4-biphenyl-2-yl-piperazin-1-yl)-butylamine:

The reaction of the phthalimide to the amine occurs analogously to Example B)d).

Batch size: 20.8 g (47.3 mmol) 2-[4-(4-biphenyl-2-yl-piperazin-1-yl)-butyl]-isoindolin-1,3-dione and 4.6 ml (94.6 mmol) hydrazine-hydrate in 185 ml ethanol.

The accumulated crude product is further processed without further purification: Yield 11.6 g (79%).

#### Example M

4-[4-(9H-fluoreen-9-yl)-piperazin-1-yl]-butylamine

a) 2-[4-[4-(9H-fluoreen-9-yl)-piperazin-1-yl]-butyl]-isoindol-in-1,3-dione:

The reaction of the piperazine with the phthalimide occurs analogously to Example 26a.

Batch size: 25 g (77.3 mmol) 1-(9-fluorenyl)-piperazine-dihydrochloride, 22.9 g (81 mmol) N-(4-bromobutyl)-phthalimide and 34 g (246 mmol) potassium carbonate in 80 ml DMF.

The purification occurs by chromatography over silica gel with  $\text{CHCl}_3/\text{CH}_3\text{OH}$  (99/1 to 90/10): Yield 30 g (86%).

b) 4-[4-(9H-fluoreen-9-yl)-piperazin-1-yl]-butylamine:

The reaction to the phthalimide to the amine occurs analogously to Example 21d.

Batch size: 33 g (76.4 mmol) 2-[4-[4-(9H-fluoreen-9-yl)-piperazin-1-yl]-butyl]-isoindolin-1,3-dione and 7.4 ml (153 mmol) hydrazine-hydrate in ethanol.

The accumulated crude product is further processed without further purification: Yield 11.5 g (46%).

4- [4- (10,11-dihydro-5H-dibenzo[a,d] cyclohepten-5-yl) -1-piperazin-1-yl] -butylamine

48 g (210 mmol) 5-chloro-10,11-dihydro-5H-dibenzo[a,d]dicyclo-heptene are dissolved in 500 ml dioxane and after addition of 45 g (522 mmol) piperazine, the mixture is heated under reflux for 7 hours with stirring. After cooling, this is concentrated under vacuum and the residue is distributed between 500 ml chloroform and 300 ml water. The aqueous phase is additionally washed three times each with 200 ml chloroform. The combined organic phases are dried over sodium sulphate and the solvent is removed under vacuum. The residue is chromatographically purified over silica gel with  $\text{CHCl}_3/\text{CH}_3\text{OH}$  (98/2 to 90/10): Yield 34.5 g (58%).

The reaction of the piperazine with the phthalimide occurs analogously to Example G)a).

Batch size: 34.5 g (124 mmol) 1-(10,11-dihydro-5H-dibenzo-[a,d]cyclohepten-5-yl)-piperazine, 35 g (124 mmol) N-(4-bromobutyl)-phthalimide and 3.7 g (24.8 mmol) sodium iodide in 80 ml DMF.

The purification occurs by chromatography over silica gel with  $\text{CHCl}_3/\text{CH}_3\text{OH}$  (98/2): Yield 55.3 g (93%).

c) 4-[4-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-yl)-1-piperazin-1-yl]-butylamine;

The reaction of the phthalimide to the amine occurs analogously to Example B)d).

Batch size: 30 g (62.5 mmol) 2-{4-[4-(10,11-dihydro-5H-di-benzo[a,d]cyclohepten-5-yl)-piperazin-1-yl]-butyl}-isoindolin-1,3-dione and 6.2 ml (125 mmol) hydrazine•hydrate in 250 ml ethanol.

100

The active ingredients according to the invention can be processed to the desired medicaments in the form of their acid addition salts, hydrates or solvates individually or in combination with each other, optionally under addition of other active ingredients. In the case of the combination of active ingredients according to the invention with other medicinals, these can also optionally be separately present next to each other in the medicine packaging, for example as tablets next to vials, depending on the requirements.

Further subject-matter of the invention is a method for the treatment of the human or animal body in which a compound or compound mixture according to formula (I), wherein the substituents have the above described meanings, is administered for treatment of tumors and/or as a cytostatic agent, cancerostatic agent or as an immunosuppressing agent, optionally in combination with further cytostatic or immunosuppressive active ingredients or other active ingredients suitable for the named indications.

Furthermore, the invention relates to a compound or compound mixture according to formula (I) for use in a therapeutic method in which the therapeutic use is carried out in connection with one or more medical indications with tumors or for immunosuppression, optimally in combination with further pharmaceuticals suitable for the named indications.

The use of one or more compounds according to formula (I), for the production of medicaments for the treatment of the human or animal body, especially in connection with one or more medical indications in the treatment of tumors or for immunosuppression, optimally in combination with further pharmaceuticals suitable in these indications or the use of compounds according to formula (I) in a corresponding diagnosis method also represent an embodiment according to



the invention, whereby the compounds for the designated medical indications are included that are excluded from claims 1 and 2 in view of the definition of group G1. The medical indications according to the invention of the compounds excluded from the protective scope of the compound claims are new.

The respective suitable tumor indications are illustrated in the last section of the description in the discussion of the pharmacological test results.

A method for the production of medicaments with an amount of one or more compounds according to formula (I) which are suitable for the processing of these active ingredients together with respective suitable pharmaceutically acceptable carriers and adjuvants for finished medicinal forms equally belongs to the scope of protection according to the invention.

Depending on the medical indication being considered, the respective suitable medicinal form is selected for the suitable therapeutic application, whereby especially 0.001 or 0.01 to 2 mg and/or 0.1, 1, 2, 5, 10, 20, 25, 30, 50, 100, 200, 300, 500, 600, 800, 1000, 2000, 3000, 4000 or 5000 mg of active ingredient according to the claims 1 to 7, 9 and 10 are applied as a dosage and/or dose unit for single administration.

The invention also relates to the use of the compounds according to formula (I) for treatment in the above indications, as well as a diagnostic agent.

The production methods of the respective suitable medicaments as well as a series of examples of medicinal forms are described in the following for better understanding of the invention.

## Therapeutic Administration Forms

The production of medicaments with an amount of one or more compounds according to the invention and/or their use in the application according to the invention occurs in the customary manner by means of common pharmaceutical technology methods. For this, the active ingredients as such or in the form of their salts are processed together with suitable, pharmaceutically acceptable adjuvants and carriers to medicinal forms suitable for the various indications and types of application. Thereby, the medicaments can be produced in such a manner that the respective desired release rate is obtained, for example a quick flooding and/or a sustained or depot effect.

Preparations for parenteral use, to which injections and infusions belong, are among the most important systemically employed medicaments for tumor treatment as well as for other indications.

Preferably, injections are administered for the treatment of tumors. These are prepared either in the form of vials or also as so-called ready-to-use injection preparations, for example as ready-to-use syringes or single use syringes in addition to perforation bottles for multiple withdrawals. Administration of the injection preparations can occur in the form of subcutaneous (s.c.), intramuscular (i.m.), intravenous (i.v.) or intracutaneous (i.c.) application. The respective suitable injection forms can especially be produced as solutions, crystal suspensions, nanoparticulate or colloid-disperse systems, such as for example, hydrosols.

The injectable formulations can also be produced as concentrates which can be adjusted with aqueous isotonic dilution agents to the desired active ingredient dosage. Furthermore, they can also be produced as powders, such as for example lyophilisates, which are then preferably dissolved or dispersed immediately before application with

03596001-051600

suitable diluents. The infusions can also be formulated in the form of isotonic solutions, fat emulsions, liposome formulations, microemulsions and liquids based on mixed micells, for example, based on phospholipids. As with injection preparations, infusion formulations can also be prepared in the form of concentrates to dilute. The injectable formulations can also be applied in the form of continuous infusions as in stationary as well as in out-patient therapy, for example in the form of mini-pumps.

Albumin, plasma expanders, surface active compounds, organic solvents, pH influencing compounds, complex forming compounds or polymeric compounds can be added to the parenteral medicinal forms, especially as substances for influencing the adsorption of the active ingredients to protein or polymers or also with the aim of decreasing the adsorption of the active ingredient to materials such as injection instruments or packaging materials, for example plastic or glass.

The active ingredients can be bound to nanoparticles in the preparations for parenteral use, for example on finely dispersed particles based on poly(meth)acrylates, polyacetates, polyglycolates, polyamino acids or polyether urethanes. The parenteral formulations can also be constructively modified as depot preparations, for example on the multiple unit principle, where the active ingredients are incorporated in a most finely distributed and/or dispersed, suspended form or as crystal suspensions, or on the single unit principle, where the active ingredient is enclosed in a medicinal form, for example, a tablet or a seed which is subsequently implanted. Often, these implantations or depot medicaments in single unit and multiple unit medicinal forms consist of so-called biodegradable polymers, such as for example, polyether urethanes of lactic and glycolic acid, polyether urethanes, polyamino acids, poly(meth)acrylates or polysaccharides.

009190.1009650

00596001-051800

Sterilized water, pH value influencing substances, such as for example organic and inorganic acids or bases as well as their salts, buffer substances for setting the pH value, agents for isotonicity, such as for example sodium chloride, monosodium carbonate, glucose and fructose, tensides and/or surface active substances and emulsifiers, such as for example, partial fatty acid esters of polyoxyethylene sorbitan (Tween®) or for example fatty acid esters of polyoxethylene (Cremophor®), fatty oils such as for example peanut oil, soybean oil and castor oil, synthetic fatty acid esters, such as for example ethyl oleate, isopropyl myristate and neutral oil (Miglyol®) as well as polymer adjuvants such as for example gelatine, dextran, polyvinylpyrrolidone, organic solvent additives which increase solubility, such as for example propylene glycol, ethanol, N,N-dimethylacetamide, propylene glycol or complex forming compounds such as for example citrates and urea, preservatives, such as for example hydroxypropyl benzoate and hydroxymethyl benzoate, benzyl alcohol, anti-oxidants, such as for example sodium sulphite and stabilizers, such as for example EDTA, are suitable as adjuvants and carriers in the production of preparations for parenteral use.

In suspensions, addition of thickening agents to prevent the settling of the active ingredients from tensides and peptizers, to secure the ability of the sediment to be shaken, or complex formers, such as EDTA, ensues. This can also be achieved with the various polymeric agent complexes, for example with polyethylene glycols, polystyrol, carboxymethylcellulose, Pluronic® or polyethylene glycol sorbitan fatty acid esters. The active ingredient can also be incorporated in liquid formulations in the form of inclusion compounds, for example with cyclodextrins. As further adjuvants, dispersion agents are also suitable. For production of lyophilisates, builders are also used, such as for example mannite, dextran, saccharose, human albumin, lactose, PVP or gelatine varieties.

As long as the active ingredients are not incorporated in the liquid medicinal formulations in the form of a base, they are used in the form of their acid addition salts, hydrates or solvates in the preparations for parenteral use.

A further systemic application form of importance is peroral administration as tablets, hard or soft gelatine capsules, coated tablets, powders, pellets, microcapsules, oblong compressives, granules, chewable tablets, lozenges, gums or sachets. These solid peroral administration forms can also be prepared as sustained action and/or depot systems. Among these are medicaments with an amount of one or more micronized active ingredients, diffusions and erosion forms based on matrices, for example by using fats, wax-like and/or polymeric compounds, or so-called reservoir systems. As a retarding agent and/or agent for controlled release, film or matrix forming substances, such as for example ethylcellulose, hydroxypropylmethylcellulose, poly(meth)acrylate derivatives (for example Eudragit®), hydroxypropylmethylcellulose phthalate are suitable in organic solutions as well as in the form of aqueous dispersions. In this connection, so-called bio-adhesive preparations are also to be named in which the increased retention time in the body is achieved by intensive contact with the mucus membranes of the body. An example of a bio-adhesive polymer is the group of Carbomers®.

For sublingual application, compressives, such as for example non-disintegrating tablets in oblong form of a suitable size with a slow release of active ingredient, are especially suitable. For purposes of a targeted release of active ingredients in the various sections of the gastrointestinal tract, mixtures of pellets which release at the various places are employable, for example mixtures of gastric fluid soluble and small intestine soluble and/or gastric fluid resistant and large intestine soluble pellets. The same goal

0956001.051500

of releasing at various sections of the gastrointestinal tract can also be conceived by suitably produced laminated tablets with a core, whereby the coating of the agent is quickly released in gastric fluid and the core of the agent is slowly released in the small intestine milieu. The goal of controlled release at various sections of the gastrointestinal tract can also be attained by multilayer tablets. The pellet mixtures with differentially released agent can be filled into hard gelatine capsules.

Anti-stick and lubricant and separating agents, dispersion agents such as flame dispersed silicone dioxide, disintegrates, such as various starch types, PVC, cellulose esters as granulating or retarding agents, such as for example wax-like and/or polymeric compounds on the basis of Eudragit®, cellulose or Cremophor® are used as a further adjuvants for the production of compressives, such as for example tablets or hard and soft gelatine capsules as well as coated tablets and granulates.

Anti-oxidants, sweetening agents, such as for example saccharose, xylite or mannite, masking flavors, aromatics, preservatives, colorants, buffer substances, direct tableting agents, such as for example microcrystalline cellulose, starch and starch hydrolysates (for example Celutab®), lactose, polyethylene glycols, polyvinylpyrrolidone and dicalcium phosphate, lubricants, fillers, such as lactose or starch, binding agents in the form of lactose, starch varieties, such as for example wheat or corn and/or rice starch, cellulose derivatives, for example methylcellulose, hydroxypropylcellulose or silica, talcum powder, stearates, such as for example magnesium stearate, aluminium stearate, calcium stearate, talc, siliconized talc, stearic acid, acetyl alcohol or hydrated fats, etc. are also used.

In this connection, oral therapeutic systems constructed especially on osmotic principles, such as for example GIT

009190.10096580

(gastrointestinal therapeutic system) or OROS (oral osmotic system), are also to be mentioned.

Effervescent tablets or tabsolute both of which represent immediately drinkable instant medicinal forms which are quickly dissolved or suspended in water are among the perorally administrable compressives.

Among the perorally administrable forms are also solutions, for example drops, juices and suspensions, which can be produced according to the above given method, and can still contain preservatives for increasing stability and optionally aromatics for reasons of easier intake, and colorants for better differentiation as well as antioxidants and/or vitamins and sweeteners such as sugar or artificial sweetening agents. This is also true for inspissated juices which are formulated with water before ingestion. Ion exchange resins in combination with one or more active ingredients are also to be mentioned for the production of liquid injestable forms.

A special release form consists in the preparation of so-called floating medicinal forms, for example based on tablets or pellets which develop gas after contact with body fluids and therefore float on the surface of the gastric fluid. Furthermore, so-called electronically controlled release systems can also be formulated by which active ingredient release can be selectively adjusted to individual needs.

A further group of systemic administration and also optionally topically effective medicinal forms are represented by rectally applicable medicaments. Among these are suppositories and enema formulations. The enema formulations can be prepared based on tablets with aqueous solvents for producing this administration form. Rectal capsules can also be made available based on gelatine or other carriers.

0055001.051600

Hardened fat, such as for example Witepsol®, Massa Estarinum®, Novata®, coconut fat, glycerol-gelatine masses, glycerol-soap-gels and polyethylene glycols are suitable as suppository bases.

For long-term application with a systematic active ingredient release up to several weeks, pressed implants are suitable which are preferably formulated on the basis of so-called biodegradable polymers.

As a further important group of systemically active medicaments, transdermal systems are also to be emphasized which distinguish themselves, as with the above-mentioned rectal forms, by circumventing the liver circulation system and/or liver metabolism. These plasters can be especially prepared as transdermal systems which are capable of releasing the active ingredient in a controlled manner over longer or shorter time periods based on different layers and/or mixtures of suitable adjuvants and carriers. Aside from suitable adjuvants and carriers such as solvents and polymeric components, for example based on Eudragit®, membrane infiltration increasing substances and/or permeation promoters, such as for example oleic acid, Azone®, adipinic acid derivatives, ethanol, urea, propylglycol are suitable in the production of transdermal systems of this type for the purpose of improved and/or accelerated penetration.

As topically, locally or regionally administration medicaments, the following are suitable as special formulations: vaginally or genitally applicable emulsions, creams, foam tablets, depot implants, ovular or transurethral administration instillation solutions. For ophthalmological application, highly sterile eye ointments, solutions and/or drops or creams and emulsions are suitable.

In the same manner, corresponding otological drops, ointments or creams can be designated for application to the ear. For

00956001-051500



both of the above-mentioned applications, the administration of semi-solid formulations, such as for example gels based on Carbopols® or other polymer compounds such as for example polyvinylpyrrolidone and cellulose derivatives is also possible.

For customary application to the skin or also to the mucus membrane, normal emulsions, gels, ointments, creams or mixed phase and/or amphiphilic emulsion systems (oil/water-water/oil mixed phase) as well as liposomes and transfersomes can be named. Sodium alginate as a gel builder for production of a suitable foundation or cellulose derivatives, such as for example guar or xanthene gum, inorganic gel builders, such as for example aluminium hydroxides or bentonites (so-called thixotropic gel builder), polyacrylic acid derivatives, such as for example Carbopol®, polyvinylpyrrolidone, microcrystalline cellulose or carboxymethylcellulose are suitable as adjuvants and/or carriers. Furthermore, amphiphilic low and high molecular weight compounds as well as phospholipids are suitable. The gels can be present either as hydrogels based on water or as hydrophobic organogels, for example based on mixtures of low and high molecular paraffin hydrocarbons and Vaseline.

Anionic, cationic or neutral tensides can be employed as emulsifiers, for example alkalized soaps, methyl soaps, amine soaps, sulfonated compounds, cationic soaps, high fatty alcohols, partial fatty acid esters of sorbitan and polyoxyethylene sorbitan, for example lanette types, wool wax, lanolin, or other synthetic products for the production of oil/water and/or water/oil emulsions.

Hydrophilic organogels can be formulated, for example, on the basis of high molecular polyethylene glycols. These gel-like forms are washable. Vaseline, natural or synthetic waxes, fatty acids, fatty alcohols, fatty acid esters, for example as mono-, di-, or triglycerides, paraffin oil or vegetable

09596001.061600

oils, hardened castor oil or coconut oil, pig fat, synthetic fats, for example based on acrylic, caprinic, lauric and stearic acid, such as for example Softisan® or triglyceride mixtures such as Miglyol® are employed as lipids in the form of fat and/or oil and/or wax-like components for the production of ointments, creams or emulsions.

Osmotically effective acids and bases, such as for example hydrochloric acid, citric acid, sodium hydroxide solution, potassium hydroxide solution, monosodium carbonate, further buffer systems, such as for example citrate, phosphate, tries-buffer or triethanolamine are used for adjusting the pH value.

Preservatives, for example such as methyl- or propyl benzoate (parabenes) or sorbic acid can be added for increasing stability.

Pastes, powders or solutions are to be mentioned as further topically applicable forms. Pastes often contain lipophilic and hydrophilic auxiliary agents with very high amounts of fatty matter as a consistency-giving base.

Powders or topically applicable powders can contain for example starch varieties such as wheat or rice starch, flame dispersed silicon dioxide or silica, which also serve as diluents, for increasing flowability as well as lubricity as well as for preventing agglomerates.

Nose drops or nose sprays serve as nasal application forms. In this connection, nebulizers or nose creams or ointments can come to use.

Furthermore, nose spray or dry powder formulations as well as controlled dosage aerosols are also suitable for systemic administration of the active ingredients.

09596001-051600

These pressure and/or controlled dosage aerosols and dry powder formulations can be inhaled and/or insufflated. Administration forms of this type also certainly have importance for direct, regional application in the lung or bronchi and larynx. Thereby, the dry powder compositions can be formulated for example as active ingredient-soft pellets, as an active ingredient-pellet mixture with suitable carriers, such as for example lactose and/or glucose. For inhalation or insufflation, common applicators are suitable which are suitable for the treatment of the nose, mouth and/or pharynx. The active ingredients can also be applied by means of an ultrasonic nebulizing device. As a propellant gas for aerosol spray formulations and/or controlled dosage aerosols, tetrafluoroethane or HFC 134a and /or heptafluoropropane or HFC 227 are suitable, wherein non-fluorinated hydrocarbons or other propellants which are gaseous at normal pressure and room temperature, such as for example propane, butane or dimethyl ether can be preferred. Instead of controlled dosage aerosols, propellant-free, manual pump systems can also be used.

The propellant gas aerosols can also suitably contain surface active adjuvants, such as for example isopropyl myristate, polyoxyethylene sorbitan fatty acid ester, sorbitan trioleate, lecithins or soya lecithin.

For regional application in situ, solutions for instillation, for example for transurethral administration in bladder tumors or genital tumors, or for profusion in liver tumors or other organ carcinomas are suitable.

The respective suitable medicinal forms can be produced in accordance with the prescription and procedures based on pharmaceutical-physical fundamentals as they are described for example in the following handbooks and are included in the present inventive subject-matter with respect to the production of the respective suitable medicaments:

09596001.061500

Physical Pharmacy (A.N. Martin, J. Swarbrick, A. Cammarata), 2nd Ed., Philadelphia Pennsylvania, (1970), German version: Physikalische Pharmazie, (1987), 3rd edition, Stuttgart;

R. Voigt, M. Bornschein, Lehrbuch der pharmazeutischen Technologie, Verlag Chemie, Weinheim, (1984), 5th edition;

P.H. List, Arzneimittelformenlehre, Wissenschaftliche Verlagsgesellschaft mbH, Stuttgart, (1985), 4th edition;

H. Sucker, P. Fuchs, P. Speiser, Pharmazeutische Technologie, Georg Thieme Verlag, Stuttgart - New York, (1991), 2nd edition;

A.T. Florence, D. Attwood, Physicochemical Principles of Pharmacy, The Maximillan Press Ltd., Hong Kong, (1981);

L.A. Trissel, Handbook on Injectable Drugs, American Society of Hospital Pharmacists, (1994), 8th edition;

Y.W. Chien, Transdermal Controlled Systemic Medications, Marcel Dekker Inc., New York - Basel, (1987);

K.E. Avis, L. Lachmann, H.A. Liebermann, Pharmaceutical Dosage Forms: Parenteral Medications, volume 2, Marcel Dekker Inc., New York - Basel, (1986);

B.W. Müller, Controlled Drug Delivery, Paperback APV, volume 17, Wissenschaftliche Verlagsgesellschaft mbH, Stuttgart, (1987);

H. Asch, D. acetic, P.C. Schmidt, Technologie von Salben, Suspensionen and Emulsionen, Wissenschaftliche Verlagsgesellschaft mbH, Stuttgart, (1984);

H.A. Liebermann, L. Lachman, J.B. Schwartz, Pharmaceutical Desage forms: Tablets, Volume 1, Marcel Dekker Inc., New York, 2nd Edition (1989);

D. Chulin, M. Deleuil, Y. Pourcelot, Powder Technology and Pharmaceutical Processes, in J.C. Williams, T. Allen, Handbook of Powder Technology, Elsevier Amsterdam - London - New York - Tokyo, (1994);

J.T. Carstensen, Pharmaceutical Principles of Solid Dosage Forms, Technomic Publishing Co., Inc., Lancaster - Basel, (1993).

009790-10092920

## PRODUCTION EXAMPLES

## 1. Injection therapeutics

## a) Parenteral Solution

active ingredient used according	
to the invention	5.000 g
acid sodium phosphate	5.000 g
sodium tartrate	12.000 g
benzyl alcohol	7.500 g
water for injection purposes	to 1000.000 ml

The solution is produced according to the customary method, sterilized and filled into 10 ml vials. One vial contains 50 mg of the compound according to the invention.

## b) Parenteral Solution

active ingredient used according to the	
invention	1.000 g
hydrochloric acid, dilute	5.000 g
sodium chloride	6.000 g
water for injection purposes	to 1000.000 ml

The solution is produced according to a customary method by stirring; the medicinal form is adjusted to a suitable pH value by acid addition and subsequently filled into 100 ml vials and sterilized. A vial contains 100 mg of the compound according to the invention.

## c) Parenteral Dispersion

active ingredient used according to	
the invention	10.000 g
soya lecithin	20.000 g
saturated triglycerides	100.000 g
sodium hydroxide	7.650 g
water for injection purposes	to 1000.000 ml

0956001.06300

The active ingredient(s) used according to the invention is dispersed in the saturated triglycerides. Then the soya lecithin is added under stirring, and subsequent to this, the aqueous solution of sodium hydroxide is added with subsequent homogenization. The dispersion is sterilized and filled into 10 ml vials. A vial contains 50 mg of the compound according to the invention.

d) Biodegradable Parenteral Depot Medicinal Form

active ingredient used according to the invention	10.000 g
polylactic acid /polyglycolic acid polymer	70.000 g
polyvinylpyrrolidone	0.200 g
gelatine	2.000 g
soya lecithin	2.000 g
isotonic sodium chloride solution	to 1000.000 ml

First, the active ingredient is incorporated into the biodegradable polymer comprising polylactic acid and polyglycolic acid by a suitable method (spray drying, solvent-evaporation or phase separation) and subsequently subjected to a sterilization process. The particles are introduced into a 2-chamber ready-made syringe in which the adjuvant solution, which is also produced in a sterile manner, is filled. The biodegradable microparticles are mixed with the dispersion agent shortly before application and dispersed. A ready-made syringe contains 200 mg of the active compound according to the invention.

e) Parenteral Dispersion for Subcutaneous Instillation

active ingredient used according to the invention	25,000 g
soya lecithin	25,000 g
arachis oil	400,000 g
benzyl alcohol	50,000 g

Miglyole®

to 1000,000 g

The active ingredient is dispersed together with soya lecithin and arachis oil. The benzyl alcohol is dissolved in Miglyole® and added to the dispersion. The entire dispersion is sterilized and subsequently filled into vials with 2 ml content. A vial contains 50 mg active ingredient.

#### f) Parenteral Perfusions Solution

The solution named under example b) can also be used for perfusion of liver for example.

According to need, instead of ampules with injection solution, so-called perforation bottles (vials), which can also be optionally preserved, and infusion solutions with an amount of one or more active ingredients according to the invention can also be made available in the customary manner under addition of buffer substances for adjustment of physiological pH value and/or the isotonicity and/or a best possible suitable pH value for the medicinal form (euhydria) and optional further required nutrients, vitamins, amino acids, stabilizers and other necessary adjuvants, possibly in combination with further medicinal agents suitable for the mentioned indications.

## 2. Solid, Peroral Administrable Medicaments

### a) Tablets

active ingredient used according to	
the invention	10.000 g
lactose	5.200 g
starch, soluble	1.800 g
hydroxypropylmethylcellulose	900 g
magnesium stearate	100 g



The above components are mixed with each other and compacted in a conventional manner, wherein a tablet weight of 180 mg is set. Each tablet contains 100 mg active ingredient. If desired, the tablets obtained in this manner are coated, provided with a film coat and/or enterically coated.

b) Coated Tablet Core

active ingredient used according to the invention	10.000 g
flame dispersed silicon dioxide	500 g
corn starch	2.250 g
stearic acid	350 g
ethanol	3.0 l
gelatine	900 g
purified water	10.0 l
talcum	300 g
magnesium stearate	180 g

From these components, a granulate is produced which is pressed to the desired coated tablet cores. Each core contains 50 mg of active ingredient. The core can be further processed in a customary manner to coated tablets. If desired, a gastric fluid resistant or retarding film coat can be applied in a known manner.

c) Vials for Drinking

active ingredient used according to the invention	0.050 g
glycerine	0.500 g
sorbite, 70% solution	0.500 g
sodium saccharinate	0.010 g
methyl-p-hydroxybenzoate	0.040 g
aromatic agent	q.s.
sterile water	q.s. to 5 ml

The above-mentioned components are mixed in a customary manner to a suspension and filled in a suitable drink vial having 5 ml content.

d) Poorly Soluble Sublingual Tablets

active ingredient used according to		
the invention		0.030 g
lactose		0.100 g
stearic acid		0.004 g
talcum purum		0.015 g
sweetener	q.s.	
aromatic agent	q.s.	
rice starch	q.s.	to 0.500 g

The active ingredient is compacted together with the adjuvants under high pressure to sublingual tablets, favourably in oblong form.

e) Soft Gel Capsule

active ingredient used according to		
the invention		0.050 g
fatty acid glyceride mixture (Miglyole®)	q.s. to	0.500 g

The active ingredient is impasted together with the fluid carrier mixture and mixed together with further adjuvants suitable for the encapsulation and filled into elastic soft gelatine capsules which are sealed.

f) Hard Gelatine Capsules

active ingredient used according to		
the invention		0.150 g
microcrystalline cellulose		0.100 g
hydroxypropylmethylcellulose		0.030 g
mannite		0.100 g
ethylcellulose		0.050 g

triethyl citrate

0.010 g

The active ingredient is mixed together with the adjuvants, microcrystalline cellulose, hydroxypropylmethylcellulose and mannite, wet with granulation liquid and formed into pellets. These are subsequently coated with a solution of ethylcellulose and triethyl citrate in organic solvents in a fluidized-bed apparatus. A hard gelatine capsule contains 150 mg of active ingredient.

### 3. Topically Administrable Medicinal Forms

#### a) Hydrophilic Ointment

active ingredient used according to

the invention	0.500 g
Eucerinum <sup>®</sup> anhydricum	60.000 g
microcrystalline wax	15.000 g
Vaseline oil	q.s. to 100.000 g

The above-mentioned adjuvants are melted and further processed together with the active ingredient to an ointment in a customary manner.

#### b) Lipophilic Ointment

active ingredient used according to

the invention	10.000 g
propylene glycol	50.000 g
paraffin, liquid	100.000 g
paraffin wax	100.000 g
Vaseline	to 1000.000 ml

The active ingredient(s) used according to the invention is dissolved in propylene glycol at ca. 60°C. At the same time, the lipophilic components are melted at 60-70°C and

subsequently combined with the active ingredient solution. The ointment is emulsified at first at 60-70°C and subsequently cooled to 35-40°C under constant emulsification and then filled in 10 g tubes. A tube contains 100 mg of the compound according to the invention.

#### 4. Inhalation Therapeutic Agent

Further subject-matter is a pharmaceutical formulation which is characterized in that it contains an active ingredient(s) used according to the invention as a base or a physiologically acceptable salt thereof together with carriers and/or diluents customary for this and suitable for administration by means of inhalation.

In connection with the production of the medicaments, particularly suitable physiologically acceptable salts of the active ingredients are, as already illustrated in the synthesis section, acid addition salts derived from inorganic or organic acids such as for example especially hydrochloride, hydrobromide, sulfate, phosphate, maleate, tartrate, citrate, benzoate, 4-methoxybenzoate, 2- or 4-hydroxybenzoate, 4-chlorobenzoate, p-tosylate, methane sulfonate, ascorbate, salicylate, acetate, formate, succinate, lactate, glutarate, gluconate or tricarballlylate.

The administration of the active ingredient(s) used of the invention by means of inhalation occurs according to the invention in conventional ways customary for administrations of this form, for example in the form of a commercial controlled dosage aerosol or in combination with a spacer. In controlled dosage aerosols, a metering valve is delivered with whose help, a dosed amount of the composition is administered. For spraying, the present compositions can be formulated for example as aqueous solutions or suspensions and be administered by means of an atomizer. Aerosol spray formulations in which the active ingredient is either suspended with one or two stabilizers in a propellant as a

carrier and/or diluent, for example tetrafluoroethane or HFC 134a and/or heptafluoropropane or HFC 227 can equally be used, whereby however, non-fluorinated hydrocarbons or other propellants which are gaseous at normal pressure and room temperature, such as propane, butane or dimethyl ether, can be preferred. Thereby, propellant-free manual pump systems or dry powder systems as described below can also be used.

Suitably, the propellant aerosols can also contain surface active adjuvants, such as for example isopropyl myristate, polyoxyethylene sorbitan fatty acid ester, sorbitan trioleate, lecithins, oleic acid.

For administration by means of inhalation and/or insufflation, the medicaments with an amount of compounds according to the invention can also be formulated in the form of dry powder compositions, for example as active ingredient-soft pellets or as an active ingredient-powder mixture with a suitable carrier, such as for example lactose and/or glucose. The powder compositions can be formulated and administered as single doses or as multiple doses.

The compounds according to the invention are preferably administered by means of a controlled dosage aerosol or in the form of a dry powder dosage formulation, wherein the latter preferably contains glucose and/or lactose as a carrier substance.

As applicators for inhalation of the pharmaceutical formulations containing one or more of the active ingredient(s) used according to the invention, all applicators are generally suitable which are suitable for controlled dosage aerosols and/or a dry powder dosage formulation, such as for example usual applicators for the nose, mouth and or pharynx, or also devices standing under propellant gas for the delivery of a spray (as controlled dosage aerosol or dry powder dosage formulation) as they are also used for inhalations in the nose, mouth and/or pharynx.

A further embodiment can also consist of an aqueous solution of the active ingredient(s) used according to the invention, which also optionally contains further active ingredients and/or additives, which are applied by means of an ultrasound atomizer.

a) Controlled Dosage Aerosol

	Intended dose per stroke	per aerosol % by weight
active ingredient used according to the		
invention	0.500 mg	0.66
stabilizer	0.075 mg	0.10
HFC 134a	75.500 mg	99.24

b) Controlled Dosage Aerosol

	Intended dose per stroke	per aerosol % by weight
active ingredient used according to the		
invention	0.250 mg	0.32
Stabilizer	0.038 mg	0.05
HFC 227	79.180 mg	99.63

In the examples a) and b) the micronized active ingredient is, after previous dispersion in a small amount of the stabilizer, placed in a suspension vessel in which the bulk amount of propellant gas solution is found. The corresponding suspension is dispersed by means of a suitable stirring system (for example high performance mixer or ultrasound mixer) until an ultra-fine dispersion results. The suspension is then continuously held in flux in a filling apparatus suitable for cold propellants or pressure fillings.

Alternatively, the suspension can also be produced in a suitable cooled stabilizer solution in HFC 134a/227.

The examples c) to d) describe the composition and production of dosage dry powder formulations.

c) Dosage-Dry Powder Formulation

mg/dose

active ingredient used according  
to the invention

0.500 mg

d) Dosage-Dry Powder Formulation

mg/dose

active ingredient used according  
to the invention  
lactose Ph.Eur.

0.500 mg

to 2.5 mg or

to 5.0 mg

e) Dosage-Dry Powder Formulation

mg/dose

active ingredient used according  
to the invention  
lactose Ph.Eur.

0.250 mg

to 2.5 mg or

to 5.0 mg

In example c) the active ingredient is formulated after micronization under addition of steam as pellets with an MMAD between 0,1 and 0,3 mm diameter and brought to use in a multi-dose powder applicator.

In the examples d) and e) the active ingredient is micronized, thereafter, bulk material is mixed with the

009330-1009560

lactose in the given amounts, and subsequently, filled in a multi-dose powder inhalator.

In all of the examples set forth above, the active ingredient or the medicinal agent in the form of the respective suitable pharmaceutical acceptable salt and/or acid addition salts can be present, insofar as the base is not preferred in each case.

## PHARMACEUTICAL EXPERIMENTAL SECTION

### 1. Growth Inhibition of Human Tumor Cells

The tumor growth inhibiting activity of the substances was determined on human tumor cells in standardized in vitro test systems. In the screening tests, the substances gave  $IC_{50}$ -values in a concentration range of 0.1 nM to 10  $\mu$ M.

#### Example 1

HepG2 cells derived from a human liver carcinoma plated at a density of 20,000 cells/ml in 12-well plastic dishes. Cultivation occurred in Richters IMEM-ZO nutrient medium with 5% foetal calf serum (FCS) in a tissue culture incubator with a gas mixture of 5%  $CO_2$  and 95% air at a temperature of 37°C. One day after plating, the culture medium was aspirated from the cells and replaced by fresh medium which contained the respective concentrations of the test substances. For the individual concentrations and the controls without test substances, three-fold batches were done for each.

Three days after the beginning of treatment, the medium was again renewed with the test compounds. After six days of substance incubation, the test was ended and the protein



amount in the individual wells was determined with the sulforhodamin-B-method (according to P. Skehan et al.: New Colorimetric Cytotoxicity Assay for Anticancer-Drug Screening. J. Natl. Cancer Inst. 82: 1107-1112, 1990). The IC<sub>50</sub>-values (defined as that concentration in which the cell growth was inhibited by 50%) was taken from the dose-response curves and given as a comparative measurement for the activity of the test compounds.

The results obtained are depicted in the following Table:

Test substance No.	IC <sub>50</sub> -value [ $\mu$ M]
29	0.6
31	0.8
32	0.5
37	0.5
47	1
158	0.2
208	0.3
227	0.6

#### Example 2

A549 cells derived from a human lung carcinoma plated at a density of 20,000 cells/ml in 12-well plastic dishes. Cultivation occurred in Richters IMEM-ZO nutrient medium with 5% foetal calf serum (FCS) in a tissue culture incubator with a gas mixture of 5% CO<sub>2</sub> and 95% air at a temperature of 37°C. One day after plating, the culture medium was aspirated from the cells and replaced by fresh medium which contained the respective concentrations of the test substances. For the individual concentrations and the controls without test substances, three-fold batches were done for each. Three days after the beginning of treatment, the medium was again renewed with the test compounds.

After four days of substance incubation, the test was ended and the protein amount in the individual wells was determined with the sulforhodamin-B-method (according to P. Skehan et al.: New Colorimetric Cytotoxicity Assay for Anticancer-Drug Screening. J. Natl. Cancer Inst. 82: 1107-1112, 1990). The  $IC_{50}$ -values (defined as that concentration in which the cell growth was inhibited by 50%) was taken from the dose-response curves and given as a comparative measurement for the activity of the test compounds.

The following results were obtained:

Test substance No.	$IC_{50}$ -value [ $\mu M$ ]
32	0.5
60	2
87	0.5
94	2
101	0.4
138	4
215	5
234	0.1

#### Example 3.

HT-29 cells derived from a human colon carcinoma plated at a density of 20,000 cells/ml in 12-well plastic dishes. Cultivation occurred in Richters IMEM-ZO nutrient medium with 5% foetal calf serum (FCS) in a tissue culture incubator with a gas mixture of 5%  $CO_2$  and 95% air at a temperature of 37°C. One day after plating, the culture medium was aspirated from the cells and replaced by fresh medium which contained the respective concentrations of the test substances. For the individual concentrations and the controls without test substances, three-fold batches were done for each. Three days after the beginning of treatment, the medium was again renewed with the test compounds. After four days of substance incubation, the test was ended and the protein amount in the

individual wells was determined with the sulforhodamin-B-method (according to P. Skehan et al.: New Colorimetric Cytotoxicity Assay for Anticancer-Drug Screening. J. Natl. Cancer Inst. 82: 1107-1112, 1990). The  $IC_{50}$ -values (defined as that concentration in which the cell growth was inhibited by 50%) was taken from the dose-response curves and given as a comparative measurement for the activity of the test compounds.

The following results were obtained:

Test substance No.	$IC_{50}$ -value [ $\mu M$ ]
30	0.7
35	3
38	0.4
97	0.1
119	0.2

#### Example 4

THP-1 cells derived from a human monocytic leukemia plated at a density of 200,000 cells/ml in 96-well plastic dishes. Cultivation occurred in RPMI 1640 nutrient medium with 10% foetal calf serum (FCS) in a tissue culture incubator with a gas mixture of 5%  $CO_2$  and 95% air at a temperature of 37°C. For the individual concentrations and the controls without test substances as well as for the background with nutrient medium but without cells, three-fold batches were done for each. After four days of substance incubation 20  $\mu l$  WST-1 reagent (Boehringer Mannheim) was respectfully pipetted in each individual well. After 30 to 60 minute incubation in the tissue culture incubator at 37°C and 5%  $CO_2$ , the light extinction was measured in an ELISA reader at 450 nm wave length. The backgrounds were each subtracted from the typical measured values. (The  $IC_{50}$ -values (defined as that concentration in which the cell growth was inhibited by 50%) was taken from the dose-response curves and given as a

comparative measurement for the activity of the test compounds.

The following results were obtained:

Test substance No.	IC <sub>50</sub> -value [ $\mu$ M]
68	0.01
87	0.02
101	0.3
119	0.02

## 2. Indications

The compounds of formula (I) and their salts permit a therapeutic use in malignant illnesses of humans and animals through their excellent inhibition of the growth of tumor cells. The anti-neoplastic activity of the described substances can be used for prophylactic, adjunct, palliative, and curative treatment of solid tumors, leukemic illnesses and lymphomas as well as for decreasing or preventing metastasis formation in humans and animals. The therapeutic use is possible in the following illnesses for example: gynaecological tumors, ovarian carcinomas, testicle tumors, prostate carcinomas, skin cancer, kidney cancer, bladder tumors, oesophagus carcinomas, stomach cancer, rectal carcinomas, pancreas carcinomas, thyroid cancer, adrenal tumors, leukemia and lymphomas, Hodgkin's disease, tumor illnesses of the CNS, soft-tissue sarcomas, bone sarcomas, benign and malignant mesotheliomas, but especially intestine cancer, liver cancer, breast cancer, bronchial and lung carcinomas, melanomas, acute and chronic leukemias. Benign papillomatosis tumors can also be limited in their growth with the named substances. The broad effectiveness of the new compounds were tested for example in very different human

tumor cells in vitro according to the methods described in point 1. Thereby, the following IC<sub>50</sub> values were obtained for the compound Nr. 119:

Cell line	Source	IC <sub>50</sub> -values
		[mM]
HT-29	colon carcinoma	0.2
A549	lung carcinoma	0.2
HepG2	hepatocellular carcinoma	0.08
THP-1	monocytic leukemia	0.02

The novelty of the compounds can be expected to have an independent activity profile in the effectiveness against the various tumor types. Thus, tumors which are resistant to customary cytostatic agents, for example, can respond entirely to these substances. In addition, based on the independent characteristics, combinations of the new compounds with known pharmaceuticals used in chemotherapy are promising as long as their properties are complimented in a suitable manner. The integration of the new structures in a therapy scheme could be successful with one or more substances from the following classes for example: anti-metabolites (for example cytarabine, 5-fluorouracil, 6-mercaptopurine, methotrexate), alkylating agents (for example busulfane, carmustine, cisplatin, carboplatin, cyclophosphamide, dacarbazine, melphalan, thiotepa), DNA-intercalating substances and topoisomerase inhibitors (for example actinomycin D, daunorubicin, doxorubicin, mitomycin C, mitoxantrone, etoposide, teniposide, topotecan, irinotecan), spindle poisons (for example vincristine, navelbine, taxol, taxotere), hormonally active agents (for example tamoxifen, flutamide, formestane, goserelin) or other cytostatic agents with complex modes of action (for example L-asparaginase, bleomycin, hydroxyurea). Resistant tumor cells can be made sensitive again for example by interaction of the new compounds with a mechanism of

resistance for common cytostatic agents (for example P-glycoprotein, MRP, glutathione-S-transferase, metallothionein).

### 3. Immuno Suppressing Activity

Many anti-tumor agents have not only a cytotoxic effect on tumor cells, but also on the blood cell system. This leads to a weakening of the immune defence, which can, in turn, be specifically employed to suppress the rejection reaction after an organ transplantation for example. Also use of the main compounds, optionally in combination with other immunological diseases (for example, psoriasis or autoimmune diseases) seems likely. In order to test the possibility for a therapeutic use in illnesses of this type, the substance activity was tested on freshly isolated lymphocytes as follows:

The spleen of a Swiss mouse served as a lymphocyte source. The lymphocyte population was isolated from the spleen cell suspension over a ficoll gradient and taken up in IMEM-ZO culture medium with 0,1% dextran 70,000 and 2% foetal calf serum. The cells were plated at a density of ca. 500,000 cells/well/ml in a 12-well plate, 1 ml doubly concentrated test substance solution was pipetted per well and this was subsequently incubated in a tissue culture incubator at 37°C and 5% CO<sub>2</sub>. After 2 days, a 1 ml-aliquot with 5 µl of the fluorescent dye solutions propidium iodide (8 mg/ml) and 3,3'-dihexyloxacarbocyanin iodide (40 µg/ml) each was added per well, and incubated for 3 minutes at room temperature. Subsequently, 10,000 cells per each sample were measured on a flow-through cytometer and the percentage amount of vital cells in the population was determined. By means of the dose-response curves, IC<sub>50</sub>-values were calculated which were also employed in the following Tables for the characterization of the individual substances:


Test Substance No.	IC <sub>50</sub> value [ $\mu$ M]
87	0.09
101	0.07
119	0.03

The independent structural class of the compounds can also be expected to be successful for an efficient combination with known immunosuppressive agents such as for example cyclosporin A, tacrolimus, rapamycin, azathioprine and glucocorticoids.

00595001.061600

## CLAIMS

1. Compounds of the general formula (I)



(I)

wherein

- $R^1$  is selected from  
hydrogen, hydroxy, halogen, cyano, aminocarbonyl,  
carboxy,

saturated, single or several-fold unsaturated, branched  
or straight chained or cyclic hydrocarbon residues such  
as alkyl, alkenyl, alkynyl or cycloalkyl,

aryl such as phenyl or heteroaryl such as pyridyl,

alkoxy, cycloalkyloxy, alkenyloxy or alkynyloxy or  
aralkyloxy such as the benzyloxy group, alkoxycarbonyl,  
alkylaminocarbonyl, dialkylaminocarbonyl, alkanoyloxy,  
alkoxycarbonyloxy, alkylthio, cycloalkylthio,  
alkenylthio, alkynylthio, aryloxy such as phenoxy,  
heteroaryloxy such as pyridyloxy, arylthio such as  
phenylthio, heteroarylthio such as pyridylthio,



trifluoromethyl,

hydroxyalkyl,

$\text{NR}^5\text{R}^6$ , wherein

$\text{R}^5$  and  $\text{R}^6$  are selected independent of each other from hydrogen, saturated or unsaturated hydrocarbon residues such as alkyl, alkenyl, alkynyl, or aryl such as phenyl and aralkyl such as benzyl;

$\text{R}^2$  is selected from hydrogen, halogen, cyano, saturated hydrocarbon residues such as alkyl, or halogenated hydrocarbon residues such as trifluoromethyl, hydroxy, alkoxy, aralkyloxy residues such as benzyloxy, as well as alkanoyloxy,

whereby  $\text{R}^1$  and  $\text{R}^2$ , in the case that they are immediately adjacent to each other, optionally form a bridge which is selected from

$-(\text{CH}_2)_4-$  and  $-(\text{CH}=\text{CH})_2-$  and  $-\text{CH}_2\text{O}-\text{CR}^7\text{R}^8-\text{O}-$ , wherein

$\text{R}^7$  and  $\text{R}^8$  are selected independently of each other from hydrogen and alkyl residues;

$\text{R}^3$  is selected from Hydrogen, halogen, saturated hydrocarbon residues such as alkyl, or halogenated hydrocarbon residues such as trifluoromethyl, or hydroxyalkyl;

$\text{R}^4$  is selected from hydrogen, hydroxy, or single or several-fold unsaturated, branched or straight chained or cyclic hydrocarbon residues such as alkyl, alkenyl, alkynyl or cycloalkyl, alkoxy and aralkyloxy such as benzyloxy;

$k$  is 0 or 1;

[illegible]

**6090**

**6090**

**6090**

**6090**

[illegible]

**6090**


**6090**

alkynylene, which is optionally substituted once or twice by alkyl, hydroxy, or alkoxy, as well as

alkylene, alkenylene or alkynylene, wherein one to three methylene units is each isosterically replaced by O, S, NR<sup>10</sup>, CO, SO or SO<sub>2</sub>, wherein

R<sup>10</sup> has the same meaning as R<sup>9</sup> but is selected independently thereof;

E



whereby

q is 1, 2 or 3;


R<sup>11</sup> is selected from hydrogen, alkyl, hydroxy, hydroxymethyl, carboxy, or alkoxy carbonyl and

R<sup>12</sup> is selected from hydrogen, alkyl or an oxo group immediately adjacent to a nitrogen atom or

R<sup>11</sup> and R<sup>12</sup> optionally together, form an alkylene bridge under formation of a bicyclic ring systems;

G is selected from G<sup>1</sup>, G<sup>2</sup>, G<sup>3</sup>, G<sup>4</sup> or G<sup>5</sup>, wherein

G<sup>1</sup> is



(G1)

whereby

$r$  has the meaning 0 to 3,

$s$  is 0 or 1;

$\text{R}^{13}$  is selected from

hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl;  
saturated or unsaturated, four to eight-membered  
heterocycles which can contain one or two hetero-atoms  
that are selected from N and/or S and /or O;  
benzyl, phenyl;  
monocyclic aromatic five- or six-membered heterocycles  
which can contain 1 to 3 hetero-atoms that are selected  
from N and/or S and/or O and are either directly bound  
or bound over a methylene group;

anellated bi- and tricyclic aromatic or partially  
hydrated carbocyclic ring systems with 8 to 16 ring  
atoms and at least one aromatic ring, whereby the bond  
can occur either over an aromatic or a hydrated ring and  
either directly or over a methylene group;

anellated bi- and tricyclic aromatic or partially  
hydrated heterocyclic ring systems with 8 to 16 ring  
atoms and at least one aromatic ring, whereby one to  
three ring atoms can be selected from N and/or S and/or  
O and the bond can occur either over an aromatic or a  
hydrated ring and either directly or over a methylene  
group;

$\text{R}^{14}$  has the same meaning as  $\text{R}^{13}$  but is selected  
independently thereof;

$\text{R}^{15}$  is selected from


hydrogen, hydroxy,  $C_1$ - $C_3$ -alkyl, aralkyl such as benzyl or aryl such as phenyl,

monocyclic aromatic five or six-membered heterocycles, which can contain one to three hetero-atoms selected from the group N and/or S and/or O and are either bound directly or over a methylene group,

annelated bi- and tricyclic aromatic or partially hydrated carbocyclic ring systems with 8 to 16 ring atoms and at least one aromatic ring, wherein the linkage can occur either over an aromatic or a hydrated ring and either directly or over a methylene group,

annelated bi- and tricyclic aromatic or partially hydrated heterocyclic ring systems with 8 to 16 ring atoms and at least one aromatic ring, wherein one to three ring atoms can be selected from N and/or S and/or O and the linkage can occur either over an aromatic ring or a hydrated ring and either directly or over a methylene group,

with the exception of compounds in which G has the meaning



in the case that the following substituents are simultaneously signify

$R^{13}$


pyridyl or (optionally halogen-, alkyl-, alkoxy- or Trifluoromethyl-substituted) phenyl,

$R^{14}$


hydrogen or (phenyl optionally substituted with halogen-, alkyl-, alkoxy- or Trifluoromethyl,

$R^{15}$  is hydrogen, and  
 A represents alkylene, optionally substituted  
 ethenylene or butadienylene,  
 D alkylene or alkenylene as well as  
 E piperazine or homopiperazine and  
 $s = 1$ ;

$G^2$  is selected from



or



whereby  $r$  and  $s$  as well as the substituents  $R^{13}$  to  $R^{15}$  can have the above meaning, or the grouping



can also be a nitrogen heterocycle bound over the nitrogen atom selected from

saturated or unsaturated monocyclic, four to eight-membered heterocycles, which, aside from the essential nitrogen atom, can still optionally contain one or two further hetero-atoms selected from N and/or S and/or O, or

saturated or unsaturated bi- or tricyclic, anellated or bridged heterocycles with 8 to 16 ring atoms, that aside from the essential nitrogen atom, can optionally still contain one or two further hetero-atoms that are selected from N and/or S and/or O;

G<sup>3</sup> has the meaning  $\text{—SO}_2\text{—(CH}_2\text{)}_r\text{—R}^{13}$  (G3).

wherein  $r$  and  $R^{13}$  have the above definition,

G<sup>4</sup> has the meaning



whereby

Ar<sup>1</sup> and Ar<sup>2</sup> can be selected independently from each other from phenyl, pyridyl or naphthyl;

G<sup>5</sup> has the meaning



whereby

R<sup>16</sup> is selected from trifluoromethyl, alkoxy, alkenyloxy, and aralkyloxy such as benzyloxy,

whereby aromatic ring systems in the substituents R<sup>1</sup>, R<sup>2</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>13</sup>, R<sup>14</sup>, R<sup>15</sup>, R<sup>16</sup>, Ar<sup>1</sup> and Ar<sup>2</sup> and/or in the ring system  $\text{—NR}^{13}\text{R}^{15}$  can be substituted independently from each other by one to three of the same or different groups selected from

halogen, cyano, alkyl, halogen alkyl such as trifluoromethyl, cycloalkyl, aryl such as phenyl, arylalkyl such as benzyl; hydroxy, hydroxy alkyl, alkoxy, alkoxy entirely or partially substituted by fluorine, aralkyloxy such as benzyloxy, aryloxy such as phenoxy; mercapto, alkylthio, carboxy, carboxyalkyl, carboxyalkenyl, alkoxycarbonyl, aralkyloxycarbonyl such as benzyloxycarbonyl, nitro, amino, monoalkylamino, dialkylamino and in the case of two adjacent residues on the aromatic ring, also methylenedioxy, and


whereby alkyl-, alkenyl- and cycloalkyl residues in the groups  $G^1$ ,  $G^2$  and  $G^3$  can be substituted by one or two of the same or different groups which are selected from hydroxy, carboxy, alkoxy, carbonyl, aralkyloxy, carbonyl such as benzyloxy, carbonyl, amino, monoalkylamino and dialkylamino;

their cis- and trans-isomers, E- and Z-isomers, especially in case that A is a cyclopropane ring or D contains one or more double bonds, including the enantiomers, diastereomers and other isomers as well as their racemic or non-racemic mixtures and the corresponding endo- and exo-isomers for the case that the ring system E is bicyclic;

their tautomeres;

their acid addition salts including their hydrates and solvates.

2. Pyridylalkane, pyridylalkene and pyridylalkine carboxamides of the formula (I)



(I)

wherein:



- R<sup>1</sup>** is selected from  
hydrogen, halogen, cyano, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-alkenyl,  
C<sub>2</sub>-C<sub>6</sub>-alkinyl, trifluoromethyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl, C<sub>1</sub>-C<sub>6</sub>-  
hydroxyalkyl, hydroxy, C<sub>1</sub>-C<sub>6</sub>-alkoxy, C<sub>3</sub>-C<sub>6</sub>-alkenyloxy,  
C<sub>3</sub>-C<sub>6</sub>-alkinyloxy, benzyloxy, C<sub>1</sub>-C<sub>7</sub>-alkanoyloxy, C<sub>2</sub>-C<sub>7</sub>-  
alkoxycarbonyloxy, C<sub>1</sub>-C<sub>6</sub>-alkylthio, C<sub>3</sub>-C<sub>6</sub>-alkenylthio,  
C<sub>3</sub>-C<sub>6</sub>-alkinylthio, C<sub>3</sub>-C<sub>8</sub>-cycloalkyloxy, C<sub>3</sub>-C<sub>8</sub>-  
cycloalkylthio, C<sub>2</sub>-C<sub>7</sub>-alkoxycarbonyl, aminocarbonyl, C<sub>2</sub>-  
C<sub>7</sub>-alkylaminocarbonyl, C<sub>3</sub>-C<sub>13</sub>-dialkylaminocarbonyl,  
carboxy, phenyl, phenoxy, phenylthio, pyridyloxy,  
pyridylthio, and NR<sup>5</sup>R<sup>6</sup>, wherein
- R<sup>5</sup>** and **R<sup>6</sup>** are selected independently of each other from  
hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-alkenyl, C<sub>3</sub>-C<sub>6</sub>-alkinyl,  
benzyl and phenyl;
- R<sup>2</sup>** is selected from  
hydrogen, halogen, cyano, C<sub>1</sub>-C<sub>6</sub>-alkyl, trifluoromethyl,  
hydroxy, C<sub>1</sub>-C<sub>6</sub>-alkoxy, benzyloxy and C<sub>1</sub>-C<sub>7</sub>-alkanoyloxy;  
**R<sup>1</sup>** and **R<sup>2</sup>**, if adjacent, optionally form a bridge  
selected from  
- (CH<sub>2</sub>)<sub>4</sub> - and - (CH=CH)<sub>2</sub> - or -CH<sub>2</sub>O-CR<sup>7</sup>R<sup>8</sup>-O-, wherein
- R<sup>7</sup>** and **R<sup>8</sup>** are selected independently from each other from  
hydrogen and C<sub>1</sub>-C<sub>6</sub>-alkyl;
- R<sup>3</sup>** is selected from  
hydrogen, halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, trifluoromethyl and  
C<sub>1</sub>-C<sub>6</sub>-hydroxyalkyl;
- R<sup>4</sup>** is selected from  
hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-alkenyl, C<sub>3</sub>-C<sub>6</sub>-alkinyl, C<sub>3</sub>-  
C<sub>6</sub>-cycloalkyl, hydroxy, C<sub>1</sub>-C<sub>6</sub>-alkoxy and benzyloxy;

k is 0 or 1,

A is selected from  
 $C_1-C_6$ -alkylene, optionally substituted one to three-fold  
 by  $C_1-C_3$ -alkyl, hydroxy,  $C_1-C_3$ -alkoxy, fluorine, or  
 phenyl,

$C_2-C_6$ -alkylene, in which a methylene unit is  
 isosterically replaced by O, S,  $NR^9$ , CO, SO or  $SO_2$ ,  
 whereby, with the exception of CO, the isosteric  
 substitution cannot be adjacent to the amide group and,  
 in  $NR^9$ , the residue  $R^9$  is selected from hydrogen,  $C_1-C_6$ -  
 alkyl,  $C_3-C_6$ -alkenyl,  $C_3-C_6$ -alkinyl,  $C_1-C_6$ -acyl or  $C_1$ -  
 $C_6$ -alkanesulfonyl,

1,2-cyclopropylene,

$C_2-C_6$ -Alkenylene, optionally substituted once to three-  
 fold by  $C_1-C_3$ -alkyl, hydroxy,  $C_1-C_3$ -alkoxy, fluorine,  
 cyano or phenyl,

$C_4-C_6$ -alkadienylene, optionally substituted once to two-  
 fold by  $C_1-C_3$ -alkyl, fluorine, cyano or phenyl;

1,3,5-hexatrienylene, optionally substituted by  $C_1-C_3$ -  
 alkyl, fluorine, cyano or phenyl, and

ethynylene


D is selected from  
 $C_2-C_{10}$ -alkylene, optionally substituted once or twice by  
 $C_1-C_6$ -alkyl, hydroxy, or  $C_1-C_6$ -alkoxy;

C<sub>4</sub>-C<sub>10</sub>-alkenylene, optionally substituted once or twice by C<sub>1</sub>-C<sub>6</sub>-alkyl, hydroxy, or C<sub>1</sub>-C<sub>6</sub>-alkoxy;

C<sub>4</sub>-C<sub>10</sub>-alkynylene, optionally substituted once or twice by C<sub>1</sub>-C<sub>6</sub>-alkyl, hydroxy, or C<sub>1</sub>-C<sub>6</sub>-alkoxy; as well as

C<sub>2</sub>-C<sub>10</sub>-alkylene, C<sub>4</sub>-C<sub>10</sub>-alkenylene or C<sub>4</sub>-C<sub>10</sub>-alkynylene, in which one to three methylene units are isosterically replaced by O, S, NR<sup>10</sup>, CO, SO or SO<sub>2</sub>, whereby R<sup>10</sup> has the same meaning as R<sup>9</sup>, but is selected independently thereof;

E signifies



whereby

q has the meaning 1, 2 or 3;

R<sup>11</sup> is selected from hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, hydroxy, hydroxymethyl, carboxy, or C<sub>2</sub>-C<sub>7</sub>-alkoxycarbonyl and

R<sup>12</sup> is selected from hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl or an oxo group adjacent to a nitrogen atom, or R<sup>11</sup> and R<sup>12</sup> optionally together, form a 1-C<sub>3</sub>-alkylene bridge under formation of a bicyclic ring system;

G is selected from G1, G2, G3, G4 or G5, whereby

G<sup>1</sup> represents  $\text{---}(\text{CH}_2)_r\text{---}(\text{CR}^{14}\text{R}^{15})_s\text{---R}^{13}$  (G<sup>1</sup>)

r has the meaning 0 to 3,

s is 0 or 1 and

R<sup>13</sup> is selected from

hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-alkenyl, C<sub>3</sub>-C<sub>6</sub>-alkinyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl;  
saturated or unsaturated, four to eight-membered heterocycles, which can contain one or two hetero-atoms that are selected from N and/or S and/or O;  
benzyl, phenyl;

monocyclic aromatic five or six-membered heterocycles, which can contain one to three hetero-atoms selected from the group N and/or S and/or O and are either bound directly or over a methylene group,

anellated bi- and tricyclic aromatic or partially hydrated carbocyclic ring systems with 8 to 16 ring atoms and at least one aromatic ring, wherein the linkage can occur either over an aromatic or a hydrated ring and either directly or over a methylene group,

anellated bi- and tricyclic aromatic or partially hydrated heterocyclic ring systems with 8 to 16 ring atoms and at least one aromatic ring, wherein one to three ring atoms can be selected from N and/or S and/or O and the linkage can occur either over an aromatic ring or a hydrated ring and either directly or over a methylene group,

R<sup>14</sup> has the same meaning as R<sup>13</sup>, but is selected independently thereof;


R<sup>15</sup> is selected from  
hydrogen, hydroxy, methyl, benzyl, phenyl,

monocyclic aromatic five or six-membered heterocycles, which can contain one to three hetero-atoms from the group N and/or S and/or O and are either bound directly or over a methylene group,

annelated bi- and tricyclic aromatic or partially hydrated carbocyclic ring systems with 8 to 16 ring atoms and at least one aromatic ring, wherein the linkage can occur either over an aromatic or a hydrated ring and either directly or over a methylene group,

annelated bi- and tricyclic aromatic or partially hydrated heterocyclic ring systems with 8 to 16 ring atoms and at least one aromatic ring, wherein one to three ring atoms can be selected from N and/or S and/or O and the linkage can occur either over an aromatic ring or a hydrated ring and either directly or over a methylene group,

whereby G in the form of G<sup>1</sup> cannot have the meaning



in the case that the following substitutions simultaneously signify

R<sup>13</sup>            pyridyl or (optionally halogen-, alkyl-, alkoxy- or Trifluoromethyl- substituted) phenyl,

R<sup>14</sup> hydrogen or (phenyl optionally substituted with halogen-, alkyl-, alkoxy- or trifluoromethyl,

R<sup>15</sup> is hydrogen, and


A represents alkylene, optionally substituted ethenylene or butadienylene,

D alkylene or alkenylene as well as


E piperazine or homopiperazine and

s = 1;

G<sup>2</sup> is selected from



or



whereby r and s as well as the substituents R<sup>13</sup> to R<sup>15</sup> can have the above meaning, or the grouping



can also be a nitrogen heterocycle bound over the nitrogen atom selected from

saturated or unsaturated monocyclic, four to eight-membered heterocycles, which, aside from the essential nitrogen atom, can optionally still contain one or two further hetero-atoms selected from N and/or S and/or O, or

saturated or unsaturated bi- or tricyclic, anellated or bridged heterocycles with 8 to 16 ring atoms, that aside from the essential nitrogen atom, can optionally still

contain one or two further hetero-atoms that are selected from N and/or S and/or O;

G<sup>3</sup> has the meaning  $\text{—SO}_2\text{—(CH}_2\text{)}_r\text{—R}^{13}$  (G3)

wherein r and R<sup>13</sup> have the above definition,

G<sup>4</sup> has the meaning



whereby

Ar<sup>1</sup> and Ar<sup>2</sup> can be selected independently from each other from phenyl, pyridyl or naphthyl;

G<sup>5</sup> has the meaning



R<sup>16</sup> is selected from trifluoromethyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, C<sub>3</sub>-C<sub>6</sub>-alkenyloxy, and benzyloxy,

whereby aromatic ring systems in the substituents are R<sup>1</sup>, R<sup>2</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>13</sup>, R<sup>14</sup>, R<sup>15</sup>, R<sup>16</sup>, Ar<sup>1</sup> and Ar<sup>2</sup> and/or in the ring system  $\text{—NR}^{13}\text{R}^{15}$  can be substituted independently from each other by one to three of the same or different groups selected from

halogen, cyano, C<sub>1</sub>-C<sub>6</sub>-alkyl, trifluoromethyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl, phenyl, benzyl, hydroxy, C<sub>1</sub>-C<sub>6</sub>-hydroxyalkyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, C<sub>1</sub>-C<sub>6</sub>-alkoxy entirely or partially substituted by fluorine, benzyloxy, phenoxy, mercapto, C<sub>1</sub>-C<sub>6</sub>-alkylthio, carboxy, C<sub>2</sub>-C<sub>7</sub>-carboxyalkyl, C<sub>2</sub>-C<sub>7</sub>-carboxyalkenyl, C<sub>2</sub>-C<sub>7</sub>-alkoxycarbonyl, benzyloxycarbonyl, nitro, amino, mono-C<sub>1</sub>-C<sub>6</sub>-

alkylamino, di-(C<sub>1</sub>-C<sub>6</sub>-alkyl)-amino and, methylene dioxide for two adjacent residues on the aromatic ring, and whereby

alkyl-, alkenyl- and cycloalkyl residues in the groups G<sup>1</sup>, G<sup>2</sup> and G<sup>3</sup> can be substituted by one or two of the same or different groups which are selected from hydroxy, carboxy, C<sub>2</sub>-C<sub>7</sub>-alkoxycarbonyl, benzyloxycarbonyl, amino, mono-C<sub>1</sub>-C<sub>6</sub>-alkylamino and di-(C<sub>1</sub>-C<sub>6</sub>-alkyl-amino);

their cis- and trans-isomers, E- and Z-isomers, especially in case that A is a cyclopropane ring or D contains one or more double bonds, including the enantiomers, diastereomers and other isomers as well as their racemic or non-racemic mixtures and the corresponding endo- and exo-isomers for the case that the ring system E is bicyclic;

their tautomeres; as well as

their acid addition salts including their hydrates and solvates.

3. Compound according to the claims 1 and 2, wherein the

R<sup>1</sup> is selected from hydrogen, halogen, cyano, C<sub>1</sub>-C<sub>6</sub>-alkyl, trifluoromethyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl, C<sub>1</sub>-C<sub>6</sub>-hydroxyalkyl, hydroxy, C<sub>1</sub>-C<sub>4</sub>-alkoxy, benzyloxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>5</sub>-alkanoyloxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>2</sub>-C<sub>5</sub>-alkoxycarbonyl, aminocarbonyl, C<sub>2</sub>-C<sub>5</sub>-alkylaminocarbonyl, C<sub>3</sub>-C<sub>9</sub>-dialkylaminocarbonyl, carboxy, phenyl, phenoxy, phenylthio, pyridyloxy, and NR<sup>5</sup>R<sup>6</sup>, whereby

R<sup>5</sup> and R<sup>6</sup> are selected independently from each other from hydrogen and C<sub>1</sub>-C<sub>6</sub>-alkyl;



- R<sup>2</sup>** is selected from  
hydrogen, halogen, cyano, C<sub>1</sub>-C<sub>6</sub>-alkyl, trifluoromethyl,  
hydroxy, C<sub>1</sub>-C<sub>4</sub>-alkoxy;
- R<sup>3</sup>** is selected from  
hydrogen, halogen and C<sub>1</sub>-C<sub>6</sub>-alkyl;
- R<sup>4</sup>** is selected from  
hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-alkenyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl,  
hydroxy, C<sub>1</sub>-C<sub>6</sub>-alkoxy and benzyloxy;
- k** has the meaning 0 or 1,
- A** is selected from  
C<sub>1</sub>-C<sub>6</sub>-alkylene, optionally substituted one to three-fold  
by C<sub>1</sub>-C<sub>3</sub>-alkyl, hydroxy, fluorine, or phenyl,  
  
C<sub>2</sub>-C<sub>6</sub>-alkylene, in which a methylene unit is  
isosterically replaced by O, S, NR<sup>9</sup>, CO, SO or SO<sub>2</sub>,  
whereby, with the exception of CO, the isosteric  
substitution cannot be adjacent to the amide group and,  
in NR<sup>9</sup>, the residue R<sup>9</sup> is selected from hydrogen, C<sub>1</sub>-C<sub>6</sub>-  
alkyl, C<sub>1</sub>-C<sub>6</sub>-acyl or methane sulfonyl;  
  
1,2-cyclopropylene,  
  
C<sub>2</sub>-C<sub>6</sub>-Alkenylene, optionally substituted once to three-  
fold by C<sub>1</sub>-C<sub>3</sub>-alkyl, hydroxy, fluorine, cyano or phenyl,  
  
C<sub>4</sub>-C<sub>6</sub>-alkadienylene, optionally substituted once to two-  
fold by C<sub>1</sub>-C<sub>3</sub>-alkyl, fluorine, cyano or phenyl;  
  
1,3,5-hexatrienylene, optionally substituted by C<sub>1</sub>-C<sub>3</sub>-  
alkyl, fluorine, cyano, and

ethynylene

D is selected from  
C<sub>2</sub>-C<sub>10</sub>-alkylene, optionally substituted once or twice by  
C<sub>1</sub>-C<sub>3</sub>-alkyl or hydroxy;


C<sub>4</sub>-C<sub>10</sub>-alkenylene, optionally substituted once or twice  
by C<sub>1</sub>-C<sub>3</sub>-alkyl or hydroxy;

C<sub>4</sub>-C<sub>10</sub>-alkynylene, optionally substituted once or twice  
by C<sub>1</sub>-C<sub>3</sub>-alkyl or hydroxy; as well as

C<sub>2</sub>-C<sub>10</sub>-alkylene, C<sub>4</sub>-C<sub>10</sub>-alkenylene or C<sub>4</sub>-C<sub>10</sub>-alkynylene,  
in which one to three methylene units in each  
isosterically replaced by O, S, NR<sup>10</sup>, CO, SO or SO<sub>2</sub>,  
whereby

R<sup>10</sup> has the same meaning as R<sup>9</sup>, but is selected  
independently thereof;

E signifies



whereby

q has the meaning 1, 2 or 3;

R<sup>11</sup> is selected from  
hydrogen, C<sub>1</sub>-C<sub>3</sub>-alkyl, hydroxy, hydroxymethyl, carboxy,  
or C<sub>2</sub>-C<sub>7</sub>-alkoxycarbonyl and

R<sup>12</sup> is selected from  
hydrogen or an oxo group adjacent to a nitrogen atom or

R<sup>11</sup> and R<sup>12</sup>, optionally together, form a C<sub>1</sub>-C<sub>3</sub>-alkylene  
bridge under formation of a bicyclic ring system;

G is selected from G<sub>1</sub>, G<sub>2</sub>, G<sub>3</sub>, G<sub>4</sub> or G<sub>5</sub>, whereby

G<sub>1</sub> represents  $\text{---}(\text{CH}_2)_r\text{---}(\text{CR}^{14}\text{R}^{15})_s\text{---R}^{13}$  (G<sub>1</sub>)

r has the meaning 0 to 2,

s is 0 or 1 and

R<sup>13</sup> is selected from

hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-alkenyl, C<sub>3</sub>-C<sub>6</sub>-alkinyl, C<sub>3</sub>-  
C<sub>8</sub>-cycloalkyl; benzyl, phenyl;

monocyclic aromatic five or six-membered heterocycles,  
which can contain one to three hetero-atoms selected  
from the group N and/or S and/or O and are either bound  
directly or over a methylene group,

annelated bi- and tricyclic aromatic or partially  
hydrated carbocyclic ring systems with 8 to 16 ring  
atoms and at least one aromatic ring, wherein the  
linkage can occur either over an aromatic or a hydrated  
ring and either directly or over a methylene group,

annelated bi- and tricyclic aromatic or partially  
hydrated heterocyclic ring systems with 8 to 16 ring  
atoms and at least one aromatic ring, wherein one to  
three ring atoms can be selected from N and/or S and/or  
O and the linkage can occur either over an aromatic ring

or a hydrated ring and either directly or over a methylene group,

R<sup>14</sup> has the same meaning as R<sup>13</sup>, but is selected independently thereof;


R<sup>15</sup> is selected from  
hydrogen, hydroxy, methyl, benzyl, phenyl;

monocyclic aromatic five or six-membered heterocycles, which can contain one to three hetero-atoms selected from the group N and/or S and/or O and are either bound directly or over a methylene group,


annelated bi- and tricyclic aromatic or partially hydrated carbocyclic ring systems with 8 to 16 ring atoms and at least one aromatic ring, wherein the linkage can occur either over an aromatic or a hydrated ring and either directly or over a methylene group,

annelated bi- and tricyclic aromatic or partially hydrated heterocyclic ring systems with 8 to 16 ring atoms and at least one aromatic ring, wherein one to three ring atoms can be selected from N and/or S and/or O and the linkage can occur either over an aromatic ring or a hydrated ring and either directly or over a methylene group;

G<sup>2</sup> is selected from



or



whereby r and s as well as the substituents  $R^{13}$  to  $R^{15}$  can have the above meaning, or the grouping



can also be a nitrogen heterocycle bound over the nitrogen atom selected from

saturated or unsaturated monocyclic, four to eight-membered heterocycles, which, aside from the essential nitrogen atom, can optionally still contain one or two further hetero-atoms selected from N and/or S and/or O, or

saturated or unsaturated bi- or tricyclic, anellated or bridged heterocycles with 8 to 16 ring atoms, that aside from the essential nitrogen atom, can optionally still contain one or two further hetero-atoms that are selected from N and/or S and/or O;

$G^3$  has the meaning  $\text{---SO}_2\text{---(CH}_2)_r\text{---R}^{13}$  (G3)

wherein r and  $R^{13}$  have the above definition,

$G^4$  has the meaning



whereby

$\text{Ar}^1$  and  $\text{Ar}^2$  can be selected independently from each other from phenyl, pyridyl or naphthyl;

$G^5$  has the meaning



R<sup>16</sup> is selected from  
trifluoromethyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, C<sub>3</sub>-C<sub>6</sub>-alkenyloxy, and  
benzyloxy,

whereby aromatic ring systems in the substituents are R<sup>1</sup>,  
R<sup>2</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>13</sup>, R<sup>14</sup>, R<sup>15</sup>, R<sup>16</sup>, Ar<sup>1</sup> and Ar<sup>2</sup> and/or in the  
ring system — NR<sup>13</sup>R<sup>15</sup> can be substituted independently from  
each other by one to three of the same or different groups  
selected from  
halogen, cyano, C<sub>1</sub>-C<sub>6</sub>-alkyl, trifluoromethyl, C<sub>3</sub>-C<sub>8</sub>-  
cycloalkyl, phenyl, benzyl, hydroxy, C<sub>1</sub>-C<sub>6</sub>-hydroxyalkyl, C<sub>1</sub>-  
C<sub>6</sub>-alkoxy, C<sub>1</sub>-C<sub>6</sub>-alkoxy entirely or partially substituted by  
fluorine, benzyloxy, phenoxy, mercapto, C<sub>1</sub>-C<sub>6</sub>-alkylthio,  
carboxy, C<sub>2</sub>-C<sub>7</sub>-carboxyalkyl, C<sub>2</sub>-C<sub>7</sub>-carboxyalkenyl, C<sub>2</sub>-C<sub>7</sub>-  
alkoxycarbonyl, benzyloxycarbonyl, nitro, amino, mono-C<sub>1</sub>-C<sub>6</sub>-  
alkylamino, di-(C<sub>1</sub>-C<sub>6</sub>-alkyl)-amino and, methylene dioxide in  
the case of two adjacent residues on the aromatic ring,

whereby alkyl-, alkenyl- and cycloalkyl residues in the  
groups G<sup>1</sup>, G<sup>2</sup> and G<sup>3</sup> can be substituted by one or two of the  
same or different groups which are selected from  
hydroxy, carboxy, C<sub>2</sub>-C<sub>7</sub>-alkoxycarbonyl, benzyloxycarbonyl,  
amino, mono-C<sub>1</sub>-C<sub>6</sub>-alkylamino and di-(C<sub>1</sub>-C<sub>6</sub>-alkyl-amino).

4. Compounds according to claims 1-3, characterized in that

R<sup>1</sup> is selected from  
hydrogen, halogen, cyano, methyl, ethyl,  
trifluoromethyl, hydroxy, C<sub>1</sub>-C<sub>4</sub>-alkoxy, benzyloxy, C<sub>1</sub>-  
C<sub>5</sub>-alkanoyloxy, methylthio, ethylthio, methoxycarbonyl,  
tert-butoxycarbonyl aminocarbonyl, carboxy, phenoxy, and  
phenylthio

R<sup>2</sup> is selected from

R<sup>3</sup> is selected from  
hydrogen, halogen;

R<sup>4</sup> is selected from  
hydrogen, C<sub>1</sub>-C<sub>3</sub>-alkyl, allyl, hydroxy and, C<sub>1</sub>-C<sub>3</sub>-alkoxy;

k is 0 or 1,

A is selected from  
C<sub>1</sub>-C<sub>6</sub>-alkylene, optionally substituted once or twice by  
C<sub>1</sub>-C<sub>3</sub>-alkyl, hydroxy or fluorine;

C<sub>2</sub>-C<sub>6</sub>-alkylene, in which a methylene unit is  
isosterically replaced by O, S, NR<sup>9</sup>, CO, SO or SO<sub>2</sub>,  
whereby, with the exception of CO, the isosteric  
substitution cannot be adjacent to the amide group,

C<sub>2</sub>-C<sub>6</sub>-alkylene, optionally substituted once or twice by  
C<sub>1</sub>-C<sub>3</sub> alkyl, hydroxy and/or fluorine;

C<sub>4</sub>-C<sub>6</sub>-alkadienylene, optionally substituted by C<sub>1</sub>-C<sub>3</sub>-  
alkyl or one or two fluorine atoms;

1,3,5-hexatrienylene, optionally substituted by,  
fluorine;


D is selected from  
C<sub>2</sub>-C<sub>8</sub>-alkylene, optionally substituted once or twice by  
methyl or hydroxy;

C<sub>4</sub>-C<sub>8</sub>-alkenylene, optionally substituted once or twice  
by methyl or hydroxy;

C<sub>4</sub>-C<sub>8</sub>-alkynylene, optionally substituted once or twice by methyl or hydroxy; and

C<sub>2</sub>-C<sub>8</sub>-alkylene, C<sub>4</sub>-C<sub>8</sub>-alkenylene or C<sub>4</sub>-C<sub>8</sub>-alkynylene, wherein one to three methylene units are each isosterically replaced by O, S, NH, N(CH<sub>3</sub>) N(COCH<sub>3</sub>), N(SO<sub>2</sub>CH<sub>3</sub>), CO, SO or SO<sub>2</sub>, whereby

E has the meaning



whereby

q is 1 or 2;

R<sup>11</sup> is selected from hydrogen, C<sub>1</sub>-C<sub>3</sub>-alkyl, hydroxymethyl, or carboxy, and

R<sup>12</sup> is selected from hydrogen or an oxo group adjacent to a nitrogen atom

G is selected from G<sub>1</sub>, G<sub>2</sub>, G<sub>3</sub>, G<sub>4</sub> or G<sub>5</sub>, whereby

G<sub>1</sub>  $\text{---} (\text{CH}_2)_r \text{---} (\text{CR}^{14}\text{R}^{15})_s \text{---} \text{R}^{13}$  (G<sub>1</sub>)  
represents

r is 0 to 2 and,

s is 0 or 1; and

R<sup>13</sup> is selected from



hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl; benzyl, phenyl;

benzocyclobutyl, indanyl, indenyl, oxoindanyl, naphthyl, dihydronaphthyl, tetrahydronaphthyl, oxotetrahydronaphthyl, biphenylenyl, fluorenyl, oxofluorenyl, anthryl, dihydroanthryl, oxodihydroanthryl, dioxodihydroanthryl, phenanthryl, dihydrophenanthryl, oxodihydrophenanthryl, dibenzocycloheptenyl, oxodibenzocycloheptenyl, dihydrodibenzocycloheptenyl, oxodihydrodibenzocycloheptenyl, dihydrodibenzocyclooctenyl, tetrahydrodibenzocyclooctenyl or oxotetrahydrodibenzocyclooctenyl bound directly or over a methylene group;

furyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, iso-thiazolyl, pyrazolyl, imidazolyl, oxadiazolyl, thiadiazolyl, triazolyl, pyridyl, pyrazinyl, pyridazinyl, pyrimidinyl, triazinyl, imidazothiazolyl, benzofuryl, dihydrobenzofuryl, benzothienyl, dihydrobenzothienyl, indolyl, indolinyl, isoindolinyl, oxoindolinyl, dioxoindolinyl, benzoxazolyl, oxobenzoxazolyl, benzooisoxazolyl, oxobenzoisoxazolyl, benzothiazolyl, oxobenzthiazolyl, benzoisothiazolyl, oxobenzoisothiazolyl, benzoimidazolyl, oxobenzoimidazolyl, indazolyl, oxoindazolyl, benzofurazanyl, benzothiadiaazolyl, benzotriazolyl, oxazolopyridyl, oxodihydrooxazolopyridyl, thiazolopyridyl, oxodihydrothiazolopyridyl, isothiazolopyridyl, imidazopyridyl, oxodihydroimidazopyridyl, pyrazolopyridyl, oxodihydropyrazolopyridyl, thienopyrimidinyl, chromanyl, chromanonyl, benzopyranyl, chromonyl, quinoloyl, isoquinoloyl, dihydroquinolyl, oxodihydroquinolyl, tetrahydroquinolyl, oxotetrahydroquinolyl, benzodioxanyl, quinoxalyl, guinazolyl, naphthyridinyl, carbazolyl, tetrahydrocarbazolyl, oxotetrahydrocarbazolyl, pyridoindolyl, acridinyl, oxodihydroacridinyl, phenanthridinyl, dihydrophenanthridinyl,

03596001-031600

oxodihydrophenanthridinyl, dibenzoisoquinolinyl, dihydrodibenzoisoquinolinyl, oxodihydrodibenzoisoquinolinyl, phenothiazinyl, dihydrodibenzooxepinyl, oxodihydrodibenzooxepinyl, benzocycloheptathienyl, oxobenzocycloheptathienyl, dihydrothienbenzothiepinyl, oxodihydrothienbenzothiepinyl, dihydrodibenzothiepinyl, oxodihydrodibenzothiepinyl, octahydrodibenzothiepinyl, dibenzoazepinyl, dihydrodibenzoazepinyl, oxodihydrodibenzoazepinyl, octahydrodibenzoazepinyl, benzocycloheptapyridyl, oxobenzocycloheptapyridyl, pyridobenzoazepinyl, dihydropyridobenzoazepinyl, oxodihydropyridobenzoazepinyl, dihydropyridobenzo-diazepinyl, dihydrodibenzooxazepinyl, dihydropyridobenzooxepinyl, dihydropyridobenzo-oxazepinyl, oxodihydropyridobenzooxazepinyl, dihydrodibenzothiazepinyl, oxodihydrodibenzothiazepinyl, dihydropyridobenzothiazepinyl or oxodihydropyridobenzothiazepinyl bound directly or over a methylene group;

R<sup>14</sup> is synonymous with R<sup>13</sup> but is selected independent thereof;


R<sup>15</sup> is selected from

hydroxy, methyl, benzyl, phenyl, indanyl, indenyl, naphthyl, dihydronaphthyl, tetrahydronaphthyl, furyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyrazolyl, imidazolyl, oxadiazolyl, thiadiazolyl, triazolyl, pyridyl, pyrazinyl, pyridazinyl, pyrimidinyl, triazinyl, benzofuryl, benzothienyl, indolyl, indolinyl, benzooxazolyl, benzothiazolyl, benzoimidazolyl, chromanyl, quinolyl or tetrahydroquinolyl bound directly or over a methylene group;

G<sup>2</sup> is selected from



[f]azepine, (5H)-Dihydrobenzo[b]pyri-do[f]azepine, (11H)-Dihydrodibenzo[b,e]oxazepine, (11H)-dihydrodibenzo[b,e]thiazepine, (10H)-dihydrodibenzo[b,f]-oxazepine, (10H)-dihydrodibenzo[b,f]thiazepine, (5H)-tetra-hydrodibenzoazocine, (11H)-dihydrobenzo[e]pyrido[b]-1,4-diazepin-6-one or (11H)-dihydrobenzo[b]pyrido[e]-1,4-diazepin-5-one.



wherein  $r$  and  $R^{13}$  have the above definition,

$G^4$  has the meaning



whereby

$\text{Ar}^1$  and  $\text{Ar}^2$  are selected independently from each other from phenyl, pyridyl or naphthyl;

$G^5$  has the meaning



$R^{16}$  is selected from trifluoromethyl,  $C_1$ - $C_6$ -alkoxy,  $C_3$ - $C_6$ -alkenyloxy, and benzyloxy,

whereby aromatic ring systems in the substitutents can be substituted independently from each other by one to three of the same or different groups selected from halogen, cyano,  $C_1$ - $C_6$ -alkyl, trifluoromethyl,  $C_3$ - $C_8$ -cycloalkyl, phenyl, benzyl, hydroxy,  $C_1$ - $C_6$ -hydroxyalkyl,  $C_1$ - $C_6$ -alkoxy,  $C_1$ - $C_6$ -alkoxy entirely or partially substituted by fluorine, benzyloxy, phenoxy, mercapto,  $C_1$ - $C_6$ -alkylthio,

carboxy, C<sub>2</sub>-C<sub>7</sub>-carboxyalkyl, C<sub>2</sub>-C<sub>7</sub>-carboxyalkenyl, C<sub>2</sub>-C<sub>7</sub>-alkoxycarbonyl, benzyloxycarbonyl, nitro, amino, mono-C<sub>1</sub>-C<sub>6</sub>-alkylamino, di-(C<sub>1</sub>-C<sub>6</sub>-alkyl)-amino and, methylene dioxide in the case of two adjacent residues on the aromatic ring, and

whereby alkyl-, alkenyl- and cycloalkyl residues in the groups G<sup>1</sup>, G<sup>2</sup> and G<sup>3</sup> can be substituted by one or two of the same or different groups which are selected from hydroxy, carboxy, C<sub>2</sub>-C<sub>7</sub>-alkoxycarbonyl, benzyloxycarbonyl, amino, mono-C<sub>1</sub>-C<sub>6</sub>-alkylamino and di-(C<sub>1</sub>-C<sub>6</sub>-alkyl-amino).

5. Compounds according to claims 1-4, wherein

R<sup>1</sup> is selected from hydrogen, fluorine, chlorine, bromine, methyl, ethyl, trifluoromethyl, hydroxy, C<sub>1</sub>-C<sub>4</sub>-alkoxy, methylthio, ethylthio, carboxy and phenoxy;

R<sup>2</sup> is selected from hydrogen, chlorine and methyl;

R<sup>3</sup> is hydrogen;

R<sup>4</sup> is selected from hydrogen, C<sub>1</sub>-C<sub>3</sub>-alkyl and hydroxy,

k is 0

A is selected from C<sub>2</sub>-C<sub>6</sub>-alkylene, which is optionally substituted once or twice by hydroxy or fluorine;

C<sub>2</sub>-C<sub>6</sub>-alkylene, wherein a methylene unit is isosterically replaced by O, S or CO, whereby, with the exception of CO, the isosteric substitution cannot be adjacent to the amide group;

C<sub>2</sub>-C<sub>6</sub>-alkenylene which is optionally substituted by C<sub>1</sub>-C<sub>3</sub>-alkyl and/or fluorine;

C<sub>4</sub>-C<sub>6</sub>-alkadienylene;

- D is selected from C<sub>2</sub>-C<sub>8</sub>-alkylene which is optionally substituted by methyl or hydroxy;


C<sub>4</sub>-C<sub>8</sub>-alkenylene, which is optionally substituted by hydroxy;

C<sub>4</sub>-C<sub>8</sub>-alkinylene, which is optionally substituted by hydroxy;


C<sub>2</sub>-C<sub>8</sub>-alkylene, C<sub>4</sub>-C<sub>8</sub>-alkenylene, C<sub>4</sub>-C<sub>8</sub>-alkinylene wherein a methylene unit is respectively isosterically replaced by O, NH, N(CH<sub>3</sub>), CO or SO<sub>2</sub> or an ethylene group is isosterically replaced by a group NH-CO and/or CO-NH or a propylene group is isosterically replaced by a group NH-CO-O and/or O-CO-NH;

- E is selected from piperazine or hexahydro-1,4-diazepine (homopiperazine), wherein the ring can be optionally substituted by one or two methylene groups and/or by an oxo group adjacent to a nitrogen atom;


- G is selected from hydrogen, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl, methoxycarbonyl, tert-butoxycarbonyl, benzyloxycarbonyl, trifluoroacetyl, diphenylphosphinoyl, or a group




and



and



and



wherein

$r$  has the meaning 0 or 1

$s$  is 0 or 1,

$R^{13}$  is selected from hydrogen, methyl, benzyl, phenyl,

indanyl, indenyl, oxoindanyl, naphthyl, tetrahydronaphthyl, fluorenyl, oxofluorenyl, anthryl, dihydroanthryl, oxodihydroanthryl, dioxodihydroanthryl, phenanthryl, dihydrophenanthryl, oxydihydrophenanthryl, dibenzocycloheptenyl, dihydrodibenzocycloheptenyl, oxodihydrodibenzocycloheptyl bound directly or over a methylene group,

furyl, thienyl, oxazolyl, isooxazolyl, thiazolyl, imidazolyl, oxadiazolyl, pyridyl, pyrazinyl, pyrimidinyl, imidazothiazolyl, benzofuryl, benzothienyl, indolyl, indolinyl, oxoindolinyl, dioxoindolinyl, benzoxazolyl, oxobenzoxazolyl, benzothiazolyl, oxobenzthiazolyl, benzimidazolyl, oxobenzimidazolyl, indazolyl, benzofurazanyl, benzotriazolyl, oxazolo-pyridyl, oxodihydrooxazolopyridyl, thiazolopyridyl, oxodihydrothiazolopyridyl, imidazopyridyl, oxodihydroimidazopyridyl, chromanyl, chromanonyl, benzopyranyl, chromonyl, quinolyl, isoquinolyl, oxodihydroquinolyl, tetrahydroquinolyl, oxotetrahydroquinolyl, benzodioxanyl, quinazolyl,

carbazolyl, acridinyl, dihydroacridinyl, oxodihydroacridinyl, dihydrophenanthridinyl dihydrobenzoisoquinolinyl, phenothiazinyl, dihydrodibenzooxepinyl, benzocycloheptathienyl, dihydrothienobenzothiepinyl, dihydrodibenzothiepinyl, oxodihydrodibenzothiepinyl, dihydrodibenzoozepinyl, oxodihydrodibenzoozepinyl, octahydrodibenzoozepinyl, benzocycloheptapyridyl, dihydropyridobenzodiazepinyl, dihydrodibenzothiazepinyl bound directly or over a methylene group,

R<sup>14</sup> is selected from hydrogen, methyl, benzyl, phenyl;

R<sup>15</sup> is selected from hydrogen, hydroxy, methyl, benzyl, phenyl;

naphthyl, tetrahydronaphthyl, furyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridyl, benzofuryl, benzothienyl, indolyl, indolinyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, chromanyl, quinolyl or tetrahydroquinolyl bound directly or over a methylene group;

whereby the grouping -NR<sup>13</sup>R<sup>15</sup> represents a ring bound over the nitrogen atom of a residue from the series

pyrrolidine, piperidine, hexahydroazepine, piperazine, hexahydrodiazepine, morpholine, hexahydroxazepine, thiomorpholine, 7-aza-bicyclo[2,2.1]heptane, 2,5-diaza-bicyclo[2.2.1]heptane, indoline, isoindoline, (1H)-dihydroquinoline, (1H)-tetrahydroquinoline, (2H)-tetrahydroisoquinoline, (4H)-dihydrobenzoxazine, (4H)-dihydrobenzothiazine, (1H)-tetrahydrobenzo[b]azepine, (1H)-tetrahydrobenzo[d]azepine, (5H)-tetrahydrobenzo[b]oxazepine, (5H)-tetrahydrobenzo[b]thiazepine, (10H)-dihydroacridine, 1,2,3,4-tetrahydroacridanone, (10H)-dihydrophenanthridine, (1H)-dihydrobenzo[d,e]isoquinoline, (10H)-phenothiazine, (5H)-dibenzo[b,f]azepine, (5H)-dihydrodibenzo[b,f]azepine,



(5H)-dihydrodibenzo[c,e]azepine, (5H)-dihydrodibenzo-diazepine, (11H)-dihydrodibenzo[b,e]oxazepine (11H)-dihydrodibenzo[b,e]thiazepine, (5H)-dihydrobenzo[b]pyrido[3,2-f]azepine and (11H)-6-oxodihydrobenzo[e]pyrido[3,2-b][1,4]diazepine, and whereby

aromatic ring systems in the substituents can be substituted, independently of each other, by one to three of the same or different groups selected from halogen, cyano, C<sub>1</sub>-C<sub>6</sub>-alkyl, trifluoromethyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl, phenyl, benzyl, hydroxy, C<sub>1</sub>-C<sub>6</sub>-hydroxyalkyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, C<sub>1</sub>-C<sub>6</sub>-alkoxy which can be entirely or partially substituted by fluorine, benzyloxy, phenoxy, mercapto, C<sub>1</sub>-C<sub>6</sub>-alkylthio, carboxy, C<sub>2</sub>-C<sub>7</sub>-carboxyalkyl, C<sub>2</sub>-C<sub>7</sub>-carbocycloalkenyl, C<sub>2</sub>-C<sub>7</sub>-alkoxycarbonyl, benzyloxycarbonyl, nitro, amino, mono-C<sub>1</sub>-C<sub>6</sub>-alkylamino or di-(C<sub>1</sub>-C<sub>6</sub>-alkyl)-amino, and in the case of two adjacent residues on the aromatic ring, methylenedioxy, and

whereby alkyl, alkenyl and cycloalkyl residues in the group G can be substituted by one or two of the same or different groups which are selected from hydroxy, carboxy, C<sub>2</sub>-C<sub>7</sub>-alkoxycarbonyl, benzyloxycarbonyl, amino, mono-C<sub>1</sub>-C<sub>6</sub>-alkylamino and ki-(C<sub>1</sub>-C<sub>6</sub>-alkyl) amino.

6. Compounds according too the claims 1-5, wherein  
R<sup>1</sup> is selected from  
hydrogen, fluorine, methyl, trifluoromethyl ethylthio;

R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are each hydrogen;

k has the meaning 0,

A is selected from

ethylene, propylene or butylene which are each optionally substituted by hydroxy or one or two fluorine atoms; or  $\text{OCH}_2$ ,  $\text{SCH}_2$ ;

ethenylene, or 1,3-butadienylene;

- D is selected from  $\text{C}_2$ - $\text{C}_6$ -alkylene which is optionally substituted by hydroxy;

$\text{C}_4$ - $\text{C}_6$  alkenylene;

$\text{C}_4$ - $\text{C}_6$  alkynylene; or

$\text{C}_2$ - $\text{C}_6$  alkylene,  $\text{C}_4$ - $\text{C}_6$  alkenylene or  $\text{C}_4$ - $\text{C}_6$  alkynylene, wherein one or two methylene units is isosterically replaced by O, NH, CO or  $\text{SO}_2$ ;

- E is selected from piperazine or hexahydro-1,4-diazeazepine;

- G is selected from phenyl, benzyl, phenethyl, diphenylmethyl, naphthyl, tetrahydronaphthyl, naphthylmethyl, fluorenyl, fluorenylmethyl, anthrylmethyl, dihydrodibenzo-cycloheptenyl;

furylmethyl, thienylmethyl, thiazolylmethyl, pyridylmethyl, benzothienylmethyl, quinolylmethyl, phenylthienylmethyl, phenylpyridylmethyl, benzocycloheptapyridinyl, dihydrobenzocycloheptapyridinyl, dihydrodibenzooxepinyl, dihydrodibenzothiepinyl, dihydrodibenzoazepinyl, dihydrobenzopyridodiazepinyl;

formyl, acetyl, pivaloyl, phenylacetyl, diphenylacetyl,

diphenylpropionyl, naphthylacetyl, benzoyl, naphthoyl, oxofluorenylcarbonyl, oxodihydroanthrylcarbonyl, dioxodihydroanthrylcarbonyl,

furoyl, pyridylacetyl, pyridylcarbonyl, chromonylcarbonyl, quinolylylcarbonyl,

phenylaminocarbonyl, naphthylaminocarbonyl, tetrahydronaphthylaminocarbonyl, dibenzylaminocarbonyl, benzylphenylaminocarbonyl, diphenylaminocarbonyl, indolinylyl-N-carbonyl, isoindolin-N-carbonyl, tetrahydroquinolinylyl-N-carbonyl, carbazolylyl-N-carbonyl, tetrahydrobenzoazepinylyl-N-carbonyl, dihydrodibenzoazepin-N-carbonyl, dihydrobenzopyridoazepinylyl-N-carbonyl, oxodihydrobenzopyridoazepinylyl-N-carbonyl;

methanesulfonyl, toluenesulfonyl, naphthylsulfonyl, quinolinsulfonyl and

diphenylphosphinoyl,

wherein aromatic ring systems can be substituted independently of each other by one to three of the same or different groups which are selected from halogen, cyano, C<sub>1</sub>-C<sub>6</sub>-alkyl, trifluoromethyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl, phenyl, benzyl, hydroxy, C<sub>1</sub>-C<sub>6</sub>-hydroxyalkyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, C<sub>1</sub>-C<sub>6</sub>-alkoxy entirely or partially substituted by fluorine, benzyloxy, phenoxy, mercapto, C<sub>1</sub>-C<sub>6</sub>-alkylthio, carboxy, C<sub>2</sub>-C<sub>7</sub>-carboxyalkyl, C<sub>2</sub>-C<sub>7</sub>-carboxyalkenyl, C<sub>2</sub>-C<sub>7</sub>-alkoxycarbonyl, benzyloxycarbonyl, nitro, amino, mono-C<sub>1</sub>-C<sub>6</sub>-alkylamino or di-(C<sub>1</sub>-C<sub>6</sub>-alkyl)-amino, and in the case of two adjacent residues in the aromatic ring methylenedioxy, and whereby

alkyl, alkenyl and cycloalkyl residues in the group G can be substituted by one or two of the same or different groups which are selected from

hydroxy, carboxy, C<sub>2</sub>-C<sub>7</sub>-alkoxycarbonyl, benzyloxycarbonyl, amino, mono-C<sub>1</sub>-C<sub>6</sub>-alkylamino or di-(C<sub>1</sub>-C<sub>6</sub>-alkyl)-amino.

7. Compounds according to formula (I) according to claim 1 or 2, characterized in that they are present in the form of the following compounds:

N-{4-(4-diphenylmethylpiperazin-1-yl)-3-hydroxybutyl}-3-pyridin-3-yl-acrylamide;  
 N-{3-(4-diphenylmethylpiperazin-1-yl)-propoxy}-3-pyridin-3-yl-acrylamide;  
 N-{4-(4-diphenylmethylpiperazin-1-yl)-4-oxo-butyl}-3-pyridin-3-yl-acrylamide;  
 N-{3-(4-diphenylmethylpiperazin-1-sulfonyl)-propyl}-3-pyridin-3-yl-acrylamide;  
 N-{2-[2-(4-diphenylmethylpiperazin-1-yl)-ethoxy]-ethyl}-3-pyridin-3-yl-acrylamide;  
 N-{4-{4-[bis-(4-fluorophenyl)-methyl]-piperazin-1-yl}-but-2-in-yl}-3-pyridin-3-yl-acrylamide;  
 N-{4-[4-(4-carboxyphenyl-phenylmethyl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-acrylamide and  
 N-{4-{4-[(4-aminophenyl)-phenylmethyl]-piperazin-1-yl}-butyl}-3-pyridin-3-yl-acrylamide.

N-{4-[4-(9H-fluoren-9-yl)-piperazin-1-yl]-butyl}-2-(pyridin-3-yloxy)-acetamide;  
 N-{5-[4-(9H-fluoren-9-yl)-piperazin-1-yl]-pentyl}-3-pyridin-3-yl-acrylamide;  
 N-{6-[4-(9H-fluoren-9-yl)-piperazin-1-yl]-hexyl}-3-pyridin-3-yl-acrylamide;  
 3-pyridin-3-yl-N-{4-[4-(1,2,3,4-tetrahydronaphthalin-1-yl)-piperazin-1-yl]-butyl}-acrylamide;

3-pyridin-3-yl-N-{4-[4-(5,6,7,8-tetrahydronaphthalin-1-yl)-piperazin-1-yl]-butyl}-acrylamide and  
N-{4-[4-(naphthalin-1-yl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-acrylamide.

N-[4-(4-biphenyl-2-yl-piperazin-1-yl)-butyl]-3-pyridin-3-yl-propionamide;

N-[5-(4-biphenyl-2-yl-piperazin-1-yl)-pentyl]-3-pyridin-3-yl-acrylamide;

N-[6-(4-biphenyl-2-yl-piperazin-1-yl)-hexyl]-3-pyridin-3-yl-acrylamide;

N-[4-(4-biphenyl-2-yl-piperazin-1-yl)-butyl]-2-(pyridin-3-yloxy)-acetamide as well as

N-[4-(4-biphenyl-2-yl-piperazin-1-yl)-butyl]-5-(pyridin-3-yl)-penta-2,4-diensäureamide.

N-{4-[4-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-yl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-propionamide;

N-{5-[4-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-yl)-piperazin-1-yl]-pentyl}-3-pyridin-3-yl-acrylamide;

N-{6-[4-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-yl)-piperazin-1-yl]-hexyl}-3-pyridin-3-yl-acrylamide;

N-{4-[4-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-yl)-piperazin-1-yl]-butyl}-5-(pyridin-3-yl)-penta-2,4-diensäureamide;

N-{4-[4-(6,11-dihydro-dibenzo[b,e]oxepin-11-yl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-propionamide and

N-{2-[4-(6,11-dihydrodibenzo[b,e]thiepin-11-yl)-piperazin-1-yl]-ethyl}-3-pyridin-3-yl-acrylamide.

N-[4-(4-diphenylacetyl-piperazin-1-yl)-butyl]-3-pyridin-3-yl-acrylamide;

N-[4-(4-benzoylpiperazin-1-yl)-butyl]-3-pyridin-3-yl-acrylamide;

N-{4-[4-(2-aminobenzoyl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-acrylamide;

N-{4-[4-(4-carboxybenzoyl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-acrylamide;


00596001-334500

N-{4-[4-(biphenyl-2-carbonyl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-acrylamide;  
 N-{4-[4-(9-oxo-9H-fluoren-4-carbonyl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-acrylamide and  
 N-{4-[4-(furan-2-carbonyl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-acrylamide.

N-{4-[4-(naphthalin-1-yl-aminocarbonyl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-propionamide;  
 N-{4-[4-(diphenylaminocarbonyl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-acrylamide;  
 N-{4-[4-(naphthalin-2-sulfonyl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-acrylamide as well as  
 N-[4-(4-diphenylphosphinonyl-piperazin-1-yl)-butyl]-3-pyridin-3-yl-acrylamide.


N-[4-(4-biphenyl-2-yl-piperazin-1-yl)-butyl]-3-pyridin-3-yl-acrylamide;  
 N-{4-[4-(9H-fluoren-9-yl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-acrylamide and  
 N-{4-[4-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-yl)-piperazin-1-yl]-butyl}-3-pyridin-3-yl-acrylamide.

8. Method for the production of compounds according to claims 1-7 according to formula (I)




(I)

(A) Characterized in that  
 carboxylic acids of formula (II)



(II)

in which  $R^1$ ,  $R^2$ ,  $R^3$ , A and k have the meaning given above or their reactive derivatives, especially their acid chlorides or activated esters, are reacted, optionally in the presence of condensation agents, with compounds of formula (III)



(III)

wherein D, E, G and  $R^4$  are defined as in claim 1 in the form of the respective free base or the respective acid addition salt, preferably in one or more inert solvents, at a temperature between  $-40^\circ\text{C}$  and  $180^\circ\text{C}$  optionally in the presence of an auxiliary base or according to the method variant.

(B) The compounds according to the general formula (I), according to the claims 1-7 are produced in the manner that compounds of the formula (I), wherein G is hydrogen, are reacted as starting materials with a compound of formula (IV)



(IV)

in which G has the meanings given in the claims 1-7, with the exception of hydrogen, and L is a suitable nucleofuge or reactive group respectively, or according to method variant.

(B1) Compounds of formula (I), in which G, with the exception of hydrogen, has the meaning of  $G^1$  according to the definition in claim 1 can also be produced, aside from method

(A), by reacting compounds of formula (I), in which G is hydrogen, with a suitable alkylation agent and/or arylation agent of formula (IV), wherein G is an alkyl-, alkenyl-, alkynyl-, cycloalkyl-, aryl-, aralkyl-, heteroaryl- or heteroaralkyl residue according to definition and the leaving group L represents a reactive derivative of an alcohol, such as a halogen atom such as chlorine, bromine or iodine or a sulfonic acid ester a methanesulfonyloxy group, trifluoromethanesulfonyloxy-, ethanesulfonyloxy-, benzene-sulfonyloxy-, p-toluenesulfonyloxy-, p-bromobenzene-sulfonyloxy-, m-nitrobenzenesulfonyloxy group or a terminal epoxide group as a reactive group, wherein this reaction occurs in a suitable inert solvent at a temperature between 0°C and 180°C, depending on the reactivity of the educt, or according to method variant.

(B2) Compounds of formula (I), in which G represents an acyl residue, a carbamoyl residue, a sulfonyl residue or a phosphinoyl residue according to the definition according to claim 1 produced in a manner by reacting compounds of formula (I), wherein G has the meaning hydrogen, with a carboxylic acid, carbamic acid, sulfonic acid and/or phosphinic acid of formula (V),



(V)

wherein G is an acyl residue, carbamoyl residue, sulfonyl residue or phosphinoyl residue according to definition, or their derivatives capable of reaction, whereby the reaction of reactive derivatives with compounds (I), in which G is hydrogen, preferably occurs in the presence of auxiliary bases in solvents and under conditions as they are described in method (A); or according to method variant.



25



25




25

25



25



in which  $\text{R}^{13}$  has the meaning given in claim 1, in an absolute, inert solvent at a temperature from  $-20^{\circ}\text{C}$  to  $150^{\circ}\text{C}$ .

9. Compound or compound mixture according to one of claims 1 to 7 for use in a therapeutic method for treatment of the human or animal body or in a corresponding diagnosis method.

10. Compound or compound mixture according to claim 9 for use in a therapeutic or diagnostic method, characterized in that the therapeutic use is in connection with cancerostatic or cytostatic or immunosuppressive treatment or abnormal cell growth and/or preventing the formation of metastases and/or proliferation, optionally in connection with suitable pharmaceutically acceptable adjuvants and carriers and/or one or more further active ingredients.

11. Use of one or more compounds according to one of claims 1 to 7 for the production of a medicament for the treatment of the human or animal body in the medical indications named above in claim 10 including all compounds which are excluded according to definition in claim 1 and 2.

12. Medicament with an amount of 1 or more active ingredients according to claim 1 to 7 optionally in connection with a pharmaceutically acceptable carrier, next to toxicologically safe adjuvants, optionally in combination with other active ingredients.

13. A method for the production of a medicament according to claim 12, characterized in that one or more compounds according to one or more of claims 1 to 7 including all compounds which are excluded according to definition in claim 1 and 2 and the respective claims dependent thereon are

processed to finished medical forms with suitable pharmacologically acceptable carriers and adjuvants.

14. Medicament according to claim 12, characterized in that it is present in a solid, peroral administrable form as a tablet, capsule, coated tablet, or as a liquid, peroral administrable solution, suspension, effervescent tablet, in the form of tabs or sachets, optionally in sustained action, and/or in gastric fluid-resistant form.

15. Medicaments according to claim 12 or 14, characterized in that it is present in the form of a suitable injection or infusion preparation together with suitable pharmaceutically acceptable carriers and adjuvants, optionally in sustained action form and or as a parenteral depot medicinal form or implant or is used in the form of a concentrate, powder or lyophilisate and the parenteral dilution agent is optionally manufactured in the packaging separately therefrom, wherein the mixing of both compounds with each other or of the active ingredient with a common parenterally applicable dilution agent occurs immediately before use.

16. Medicament according to claim 12, characterized in that it is present in the form of an inhalation therapeutic agent, for example, in the form of a spray together with suitable pharmaceutically acceptable propellants, carriers and adjuvants.

17. Medicament according to claim 12, characterized in that it is present in the form of a transdermal therapeutic system for systemic treatment.

18. Medicament according to claim 12, characterized in that it is present in the form of a gastrointestinal therapeutic system (GITS) for systemic treatment.

19. Medicament according to claim 12, characterized in that it is present in the form of a salve, suspension, emulsion, a

00595001-051500

balm or plaster or in the form of an externally applicable solution.

20. Medicament according to claim 16 for administration by means of a controlled dosage aerosol or in the form of a dry powder dosage formulation.

21. Medicament according to claim 12, characterized in that it is present in the form of a rectal, genital, or transurethral administration emulsion, a solution, a liposomal solution, an implant, suppository or a capsule.

22. Medicament according to claim 12, characterized in that it is present in the form of a composition capable of being applied nasally, otologically or ophthalmologically.

23. Medicament according to one of the claims 12 or 14, characterized in that it is present in the form of a buccally applicable form.

24. Medicament according to one of the claims 12 and 14 to 16, characterized in that a dosage unit for single administration contains 0.001 or 0.01 to 2.0 mg or 0.1, 1, 2, 5, 10, 20, 25, 30, 50, 100, 200, 300, 500, 600, 800, 1000, 2000, 3000, 4000 or 5000 mg active ingredient according to the claims 1 to 7, 9 and 10.

25. Medicament according to claim 16, characterized in that the pharmaceutically acceptable carrier and/or diluent is a propellant aerosol.

26. Medicament according to claim 16 or 25, characterized in that the propellant aerosol is tetrafluoroethane and/or heptafluoropropane and/or propane, butane, or dimethyl ether or mixtures thereof.

27. Medicament according to one of the claims 16, 25 or 26, characterized in that the propellant aerosol contains surface active adjuvants.

28. Medicament according to one of the claims 12 or 16, characterized in that it contains glucose and/or lactose as a dry powder dosage.

29. Substance or substance mixture according to one of the claims 9 or 10, characterized in that the therapeutic use occurs with a further cytostatic agent or immunosuppressive agent.

30. Medicament according to one of the claims 12 and 14 to 28, characterized in that it is present in combination with a further cytostatic agent or immunosuppressive agent, optionally in the form of separate dosage units in the pharmaceutical package.

31. Use of one or more compounds according to the general formula (I) according to claims 1 to 6 and according to claim 7 including compounds which are excluded according to definition in claims 1 and 2 for cytostatic and/or cancerostatic or immunomodulatory and/or immunosuppressive treatment, optionally in combination with a further cytostatic agent or immunosuppressive agent and/or further medicaments suitable for these indications.

## ABSTRACT

The invention relates to new piperazinyl-substituted pyridylalkane, alkene, and alkine acid amides substituted with saturated or one or several-fold unsaturated hydrocarbon residue in the carboxylic acid group according to the general formula (I) as well as methods for the production of these compounds, medicaments containing these and their production as well as their therapeutic use, especially as cytostatic agents and immunosuppressive agents, for example in the treatment or prevention of various types of tumors and control of immune reactions such as autoimmune diseases.

